The Registered Nurse Population

Findings from the 2008 National Sample Survey of Registered Nurses

September 2010 U.S. Department of Health and Human Services Health Resources and Services Administration



The Registered Nurse Population

Findings from the 2008 National Sample Survey of Registered Nurses

September 2010

U.S. Department of Health and Human Services Health Resources and Services Administration

Preface

The U.S. Department of Health and Human Services (HHS), Health Resources and Services Administration (HRSA), is the principal Federal agency responsible for the evaluation and development of the nursing workforce in the United States. HRSA's Bureau of Health Professions (BHPr) works to appraise and improve the U.S. nursing workforce with the goal of ensuring there are enough qualified nurses to meet the Nation's health needs. To do this, BHPr reviews the supply, composition, and distribution of nurses on the national and State levels. To that end, BHPr commissioned the National Sample Survey of Registered Nurses (NSSRN).

This publication is the report of the ninth NSSRN representing data collected about registered nurses (RNs) holding active nursing licenses as of March 2008. As with previous iterations of the NSSRN, this report is available to all primary stakeholders involved in health care planning and evaluation as well as the public. The 2008 NSSRN public use files and code book are available to researchers and analysts and can be accessed at http://bhpr.hrsa.gov/healthworkforce/rnsurvey/default.htm.

The 2008 NSSRN was developed for the U.S. Department of Health and Human Services, Health Resources and Services Administration under contract with Westat. The study was overseen by BHPr and funded through the Division of Nursing.

The first national survey was conducted in 1977; HRSA has conducted this national survey every 4 years since 1980. Data from these periodic surveys provide the basis for evaluating trends and projection of the future supply of nursing resources at the national and State levels. The NSSRN is the cornerstone of nursing workforce data. It is the principal data source used for disseminating information to the Federal Government, researchers, and the public on the nursing workforce. It provides essential information for performing supply-demand projections of nursing requirements and foreseen shortages. Periodic monitoring of the number and characteristics of the registered nurse population is vital to effectively maintain an up-to-date picture of the RN population and to assess the future availability of this critical resource. For example, previous surveys were integral in identifying the shortages of RNs that occurred at the end of the 1980s and then reappeared around 2000.

The NSSRN is the Nation's most extensive and comprehensive source of statistical data on all individuals in the United States that hold active RN licenses, whether they are currently practicing

nursing or not. The survey assesses the number of RNs; their educational background; employment setting, position, and specialty areas; job satisfaction; and salaries. In addition, the survey also identifies the geographic distribution of nurses throughout the United States, as well as the personal composition of the U.S. nursing workforce in terms of gender, racial/ethnic background, age, and family structure.

Table of Contents

| <u>Chapter</u> | | Page |
|----------------|---|--|
| | Preface | |
| | Executive Summary | xxvii |
| | Introduction | XXXV |
| 1 | Survey Methods | 1-1 |
| | 1.1 Summary 1.2 2008 Questionnaire 1.3 Sample Design 1.4 Data Collection Methods 1.5 Sample Performance 1.6 Imputation 1.7 Sample Weighting 1.7.1 Overview of the Weighting Process | 1-3 1-4 1-5 1-6 1-7 1-9 |
| | 1.7.2 Implications of Weighting for NSSRN Data Users | 1-9 |
| 2 | Education and Licensing of Registered Nurses | 2-1 |
| | 2.1 Introduction 2.2 Initial Nursing Education 2.3 Age at Graduation 2.4 Sources of Financing 2.5 Education and Experience Prior to Initial Nursing | 2-1 2-2 |
| | Education 2.6 Nursing Licensure 2.7 Degrees Obtained After Initial Nursing Education 2.8 Currently Enrolled Students 2.9 Emergency Preparedness 2.10 Certifications | 2-6 2-7 2-11 2-12 |
| 3 | The Employment of Registered Nurses | 3-1 |
| | 3.1 Introduction | 3-1 3-3 3-5 |

| | 3.5.1 | Employment Setting Trends | 3-11 |
|------|----------------|--|------|
| | 3.5.2 | Employment Settings and Age | 3-11 |
| | 3.5.3 | Employment Settings and Education | 3-13 |
| 3.6 | Job Titl | es | 3-14 |
| | 3.6.1 | Job Titles and Employment Settings | 3-15 |
| | 3.6.2 | Job Titles and Experience | 3-16 |
| | 3.6.3 | Job Titles and Education | 3-16 |
| 3.7 | Second | ary Nursing Jobs | 3-17 |
| 3.8 | | of Registered Nurses | 3-19 |
| | 3.8.1 3.8.2 | Hours and Weeks Worked and Age Group Hours and Weeks Worked and Highest | 3-24 |
| | | Education | 3-24 |
| | 3.8.3 | On-Call Work | 3-25 |
| | 3.8.4 | Overtime Work | 3-25 |
| | 3.8.5 | Hours Worked and Employment Setting | 3-26 |
| | 3.8.6 | Hours Worked and Job Title | 3-28 |
| 3.9 | Employ | ment Settings and Level of Care | 3-29 |
| | 3.9.1 | Education and Areas of Clinical Specialty | 3-31 |
| 3.10 | Compo | sition of Nursing Work | 3-33 |
| | 3.10.1 | Composition of Work and Job Title | 3-34 |
| | 3.10.2 | Composition of Work and Education | 3-34 |
| 3.11 | Tempor | rary and Travel Work | 3-36 |
| 3.12 | Crossin | g State Lines and Compact States | 3-37 |
| 3.13 | Unioniz | zation in Principal Nursing Positions | 3-37 |
| 3.14 | Earning | gs of Registered Nurses | 3-38 |
| | 3.14.1 | Earnings and Job Titles | 3-40 |
| | 3.14.2 | Earnings and Employment Setting | 3-41 |
| | 3.14.3 | Earnings From Secondary Jobs | 3-41 |
| | 3.14.4 | Earnings and Experience | 3-42 |
| | 3.14.5 | Earnings and Education | 3-42 |
| 3.15 | Job Sati | sfaction | 3-43 |
| | 3.15.1 | Job Satisfaction and Employment Setting | 3-45 |
| | 3.15.2 | Job Satisfaction and Job Titles | 3-45 |

| | | 3.15.3 3.15.4 | Job Satisfaction and Age Job Satisfaction and Education | 3-48 3-48 |
|---|-------|------------------|--|--------------|
| | | | | |
| | 3.16 | | inges of Registered Nurses | 3-51 |
| | 3.17 | | or Future Employment | 3-53 |
| | 3.18 | Non-N | ursing Jobs | 3-54 |
| 4 | Nurse | Faculty | | 4-1 |
| | 4.1 | Introdu | ction | 4-1 |
| | 4.2 | Numbe | r of Nurses Working in Faculty Roles | 4-2 |
| | 4.3 | • | and Age | 4-3 |
| | 4.4 | • | and Education | 4-4 |
| | 4.5 | - | and Hours of Work | 4-5 |
| | 4.6 | Faculty | Earnings and Satisfaction | 4-6 |
| 5 | Advan | iced Pract | tice Registered Nurses | 5-1 |
| | 5.1 | Introdu | ction | 5-1 |
| | 5.2 | Nurse I | Practitioners | 5-1 |
| | | 5.2.1 | Number of Nurse Practitioners | 5-3 |
| | | 5.2.2 | Nurse Practitioners and Education | 5-5 |
| | | 5.2.3 | Nurse Practitioners and Employment | 5-7 |
| | | 5.2.4 | Nurse Practitioner Earnings and Job Satisfaction | 5-10 |
| | 5.3 | Nurse N | Midwives | 5-13 |
| | | 5.3.1 | Number of Nurse Midwives | 5-14 |
| | | 5.3.2 | Nurse Midwives and Education | 5-15 |
| | | 5.3.3 | Nurse Midwives and Employment | 5-17 |
| | | 5.3.4 | Nurse Midwife Earnings and Job Satisfaction | 5-19 |
| | 5.4 | Nurse A | Anesthetists | 5-20 |
| | | 5.4.1 | Number of Nurse Anesthetists | 5-21 |
| | | 5.4.2 | Nurse Anesthetists and Education | 5-22 |
| | | 5.4.3 | Nurse Anesthetists and Employment | 5-24 |
| | | 5.4.4 | Nurse Anesthetist Earnings and Job Satisfaction | 5-25 |
| | 5.5 | Clinical | Nurse Specialists | 5-26 |
| | | 5.5.1 | Number of Clinical Nurse Specialists | 5-26 |
| | | 5.5.2 | Clinical Nurse Specialists and Education | 5-27 |
| | | 5.5.3 | Clinical Nurse Specialists and Employment | 5-30 |

| 6 | Nurs | | |
|---|--------|---|----|
| | 1 NULS | es Who Do Not Work in Nursing | 6 |
| | 6.1 | Introduction | 6 |
| | 6.2 | Demographics of Nurses Not Working in Nursing | 6 |
| | | 6.2.1 Nurses Not Working in Nursing and Age | 6 |
| | | 6.2.2 Presence of Children | 6 |
| | 6.3 | Length of Time Away From Nursing Work | 6 |
| | 6.4 | Reasons for Not Working in Nursing | 6 |
| | 6.5 | Plans Regarding Nursing Work | 6 |
| 7 | Dem | ographics of the Registered Nurse Workforce | 7 |
| | 7.1 | Introduction | 7 |
| | 7.2 | Registered Nurses and Age | 7 |
| | 7.3 | Registered Nurses and Gender | 7 |
| | 7.4 | Registered Nurses and Race and Ethnicity | 7 |
| | 7.5 | Registered Nurses and Language Competency | 7 |
| | 7.6 | Registered Nurses and Marital Status and Dependents | 7 |
| | 7.7 | Registered Nurses and Household Income | 7 |
| | 7.8 | Geography, Nurse Demographics, and Employment | 7 |
| 8 | Intern | nationally Educated Nurses (IENs) | 8 |
| | 8.1 | Introduction | 8 |
| | 8.2 | IEN Licensure and National Origin | 8 |
| | 8.3 | IEN Demographic Characteristics | 8 |
| | 8.4 | IEN Employment | 8 |
| 9 | Recen | nt Registered Nurse Graduates | 9 |
| | 9.1 | Introduction | 9 |
| | 9.2 | Demographic Characteristics of Recent Registered Nurse Graduates | 9 |
| | 9.3 | Recent Registered Nurse Graduates and Education | 9. |
| | 9.4 | Recent Registered Nurse Graduates and Diversity | 9. |
| | 9.5 | Family Situation of Recent Nurse Graduates | 9. |
| | 9.6 | Recent Registered Nurse Graduates and Employment | 9. |
| | 9.7 | Recent Registered Nurse Graduates and Job Changes | 9 |

<u>Page</u>

| | 9.8 | Recent Registered Nurse Graduates and Secondary Nursing Jobs |
|----|---------|---|
| | 9.9 | Recent Registered Nurse Graduates and Earnings9-139-14 |
| 10 | Registe | ered Nurses Nearing Retirement 10-1 |
| | 10.1 | Introduction |
| | 10.2 | Demographic Characteristics of Registered Nurses |
| | | Nearing Retirement |
| | 10.3 | Employment Characteristics of Older Registered Nurses10-2 |
| | | 10.3.1 Employment Positions and Settings |
| | | 10.3.2 Hours Worked by Older Registered Nurses |
| | | 10.3.3 Secondary Nursing Employment |
| | | 10.3.4Earnings of Older Registered Nurses10-6 |
| | 10.4 | Job Changes and Intentions Regarding Nursing Work 10-6 |
| | 10.5 | Older Registered Nurses Not Working in Nursing 10-8 |

Appendixes

| А | Data | Fables | | A-1 |
|---|-------|-------------------------|--|----------------------|
| В | Surve | y Method | ology | B-1 |
| | B.1 | Samplir | ng | B-1 |
| | | B.1.1 B.1.2 | Sample Selection Final Sample Allocation | B-1 B-3 |
| | B.2 | Sample | Performance | B-3 |
| | | B.2.1 | Vermont Sample Performance | B-11 |
| | B.3 | Imputa | tion | B-13 |
| | | B.3.1 B.3.2 B.3.3 | Imputation Methods Procedures Results and Implications on Variance Estimates | B-14 B-15 B-16 |
| | B.4 | Weighti | ing and Estimation | B-17 |
| | | B.4.1 B.4.2 | Sample Weighting Estimation, Variance Estimation, and Sampling | B-18 |
| | | | and Nonsampling Errors | B-24 |

Appendixes

<u>Page</u>

| | B.5 | Nonres | Nonresponse Bias Analysis | |
|---|-------|----------------|---|--------------|
| | | B.5.1 | Underrepresentation of Racial and Ethnic Minorities | B-29 |
| | | B.5.2 B.5.3 | Underrepresentation of Younger Nurses Comparative Analyses | B-30 B-31 |
| | Refer | | | B-41 |
| С | NSSI | N Quest | ionnaire | C-1 |

Tables

| 1-1 | Summary of 2008 NSSRN sample performance | 1-7 |
|-----|---|------|
| 2-1 | Degrees prior to initial nursing education, by survey year, 2000-2008 | 2-5 |
| 2-2 | Education and health occupation experience prior to initial nursing education, by initial nursing education | 2-6 |
| 2-3 | Highest nursing degree, by employment status | 2-7 |
| 2-4 | Master's and doctorate degrees as highest nursing or nursing- related education preparation, 2000-2008 | 2-9 |
| 2-5 | Average years between completion of initial and highest nursing degrees | 2-11 |
| 2-6 | Enrollment in nursing and nursing-related education programs | 2-12 |
| 2-7 | Registered nurse certifications | 2-13 |
| 3-1 | Employment in hospital units, by type of hospital | 3-10 |
| 3-2 | Employment settings, by age group | 3-13 |
| 3-3 | Job titles of registered nurses | 3-15 |
| 3-4 | Job titles, by experience in nursing | 3-16 |
| 3-5 | Job titles, by highest nursing or nursing-related education | 3-17 |

<u>Tables</u>

| 3-6 | Employment arrangements in secondary nursing positions | 3-19 |
|------|---|------|
| 3-7 | Employment settings of secondary nursing positions | 3-19 |
| 3-8 | Average hours and weeks worked in secondary nursing positions | 3-23 |
| 3-9 | Average hours per year worked in all nursing jobs, by number of nursing jobs | 3-23 |
| 3-10 | Average hours worked per year in all nursing positions, by age group | 3-24 |
| 3-11 | Average hours worked per year in all nursing positions, by highest nursing and nursing-related education | 3-24 |
| 3-12 | Average hours on standby or worked from on-call per week in the principal nursing position | 3-25 |
| 3-13 | Average hours per week in principal nursing position, by full- time/part-time status and employment setting | 3-27 |
| 3-14 | Average hours per week in principal nursing position, by full- time/part-time status and job title | 3-28 |
| 3-15 | Levels of care in nursing work for hospital and nonhospital settings | 3-29 |
| 3-16 | Primary patient age group cared for in principal nursing position, for hospital and nonhospital settings | 3-30 |
| 3-17 | Primary clinical specialty in principal nursing position for hospital and nonhospital settings | 3-31 |
| 3-18 | Primary clinical specialty in principal nursing position, by highest nursing and nursing-related education | 3-32 |
| 3-19 | Average percent time spent on various activities, by highest nursing or nursing-related education | 3-35 |
| 3-20 | Employment by temporary employment services (TES), for primary and secondary nursing positions, by principal nursing position in a hospital | 3-36 |
| 3-21 | Employment settings of secondary employment through a nontravel temporary employment service | 3-36 |

<u>Tables</u>

| 3-22 | Licensure in a Compact State, by where registered nurses live and work | 3-37 |
|------|--|------|
| 3-23 | Unionization, by employment setting | 3-38 |
| 3-24 | Average annual earnings from all nursing positions, by total number of hours worked per year in all nursing positions | 3-39 |
| 3-25 | Average annual earnings, by job title of principal nursing position, for registered nurses working full-time | 3-40 |
| 3-26 | Average annual earnings, by employment setting of principal nursing position, for registered nurses working full-time | 3-41 |
| 3-27 | Average annual earnings for registered nurses working full-time in their principal nursing position, by years since initial nursing education | 3-42 |
| 3-28 | Average annual earnings of registered nurses working full-time in their principal nursing position, by highest nursing or nursing- related education and job title | 3-43 |
| 3-29 | Employment status in 2008, compared with status in 2007 | 3-52 |
| 3-30 | Employment changes among registered nurses employed in nursing in both 2007 and 2008 | 3-53 |
| 3-31 | Employment in non-nursing occupations, by nursing employment status | 3-57 |
| 4-1 | Registered nurses employed as faculty in their principal nursing position, by type of nursing education program | 4-2 |
| 4-2 | Registered nurses employed as faculty in their secondary nursing position(s), by job title of principal nursing position | 4-2 |
| 4-3 | Months worked per year and hours worked per week, by registered nurses who are faculty in their principal nursing position | 4-6 |
| 4-4 | Average annual earnings of registered nurses who work full-time as faculty in their principal nursing position | 4-7 |
| 5-1 | Certification and State Board recognition of nurse practitioners employed in nursing, and whether certification or recognition is required for their job | 5-6 |

Tables

| 5-2 | Primary clinical specialties of employed nurse practitioners with job title of nurse practitioner | 5-9 |
|-----|---|------|
| 5-3 | Employed nurse practitioners whose primary level of care is ambulatory or primary care, by job title5 | 5-9 |
| 5-4 | Certification and State Board recognition of nurse midwives employed in nursing, and whether certification or recognition is required for their job | 5-17 |
| 5-5 | Clinical specialties of registered nurses prepared as nurse midwives, for selected job titles in the hospital setting | 5-19 |
| 5-6 | Certification and State Board recognition of nurse anesthetists employed in nursing, and whether certification or recognition is required for their job | 5-23 |
| 5-7 | Certification and State Board recognition of clinical nurse specialists employed in nursing, and whether certification or recognition is required for their job | 5-29 |
| 6-1 | Intentions of nurses who are not working in nursing, overall, and by marital status | 6-6 |
| 7-1 | Median age of registered nurses | 7-2 |
| 7-2 | Gender of employed registered nurses, by year of initial registered nurse license | 7-3 |
| 7-3 | Highest education of registered nurses working in nursing, by gender | 7-3 |
| 7-4 | Employment settings, by gender | 7-4 |
| 7-5 | Median current age and age at licensure of registered nurses working in nursing, by race/ethnicity | 7-8 |
| 7-6 | Initial nursing education of registered nurses employed in nursing, by race/ethnicity | 7-8 |
| 7-7 | Highest education of registered nurses employed in nursing, by race/ethnicity | 7-9 |
| 7-8 | Advanced practice preparation of registered nurses, by race/ethnicity | 7-9 |

<u>Tables</u>

| 7-9 | Employment setting of registered nurses, by race/ethnicity | 7-10 |
|------|--|------|
| 7-10 | Job titles of registered nurses, by race/ethnicity | 7-10 |
| 7-11 | Marital status and presence of children or adult dependents at home | 7-11 |
| 7-12 | Employment of registered nurses, by presence of children or adult dependants in the home | 7-12 |
| 7-13 | Annual household income of registered nurses, by employment status of the nurse | 7-12 |
| 8-1 | Internationally educated nurses in the United States | 8-2 |
| 8-2 | Years between initial nursing education and first U.S. licensure | 8-3 |
| 8-3 | Median current age and age at licensure of registered nurses employed in nursing, by international education status | 8-3 |
| 8-4 | Men among U.S. and internationally educated employed nurses, by year of licensure | 8-4 |
| 8-5 | Registered nurses working in nursing with at least a bachelor's degree, by location of education | 8-4 |
| 8-6 | Registered nurses working full-time, by location of education and year of licensure | 8-5 |
| 8-7 | Average hours worked per year and per week in principal nursing position, by full-time and part-time status and location of education | 8-5 |
| 8-8 | Number and percent of nurses employed in a hospital setting for their principal nursing position, by location of initial education and year of licensure | 8-6 |
| 8-9 | Registered nurses employed in ambulatory care and academic education, by location of initial education | 8-7 |
| 8-10 | Registered nurses employed as staff nurses, by location of initial education and year of licensure | 8-7 |
| 8-11 | Employment intentions, by location of initial education | 8-7 |

<u>Tables</u>

| 9-1 | Estimated population of registered nurses, by graduation cohort | 9-2 |
|------|--|------|
| 9-2 | Initial education of registered nurses, by graduation cohort | 9-2 |
| 9-3 | Location of initial education of nurses, by graduation cohort | 9-3 |
| 9-4 | Initial nursing education, by graduation cohort and location of education | 9-3 |
| 9-5 | Average age at graduation and age distribution at graduation, by graduation cohort | 9-3 |
| 9-6 | Registered nurses with postsecondary education degrees before starting initial nursing education program | 9-4 |
| 9-7 | Fields of study for registered nurses who had postsecondary degrees before initial nursing education, by graduation cohort | 9-4 |
| 9-8 | Employment in a health occupation, by registered nurses before initial nursing education, by graduation cohort | 9-5 |
| 9-9 | Sources of financing for initial nursing education, by graduation cohort | 9-6 |
| 9-10 | Highest level of nursing or nursing-related education for recent registered nurse graduates, by initial nursing education (2001– 2004) | 9-6 |
| 9-11 | Gender of registered nurses, by graduation cohort | 9-7 |
| 9-12 | Racial/ethnic composition of registered nurse population, by graduation cohort | 9-7 |
| 9-13 | Marital status and presence of children, by graduation cohort | 9-8 |
| 9-14 | Employment of registered nurses, by graduation cohort | 9-9 |
| 9-15 | Employment of new registered nurse graduates, by presence of children | 9-9 |
| 9-16 | Employment setting of principal nursing position, by graduation cohort, 2008 and 2004 | 9-9 |
| 9-17 | Job titles of registered nurses, by graduation cohort | 9-10 |

<u>Tables</u>

| 9-18 | Average hours worked per year and per week by registered nurses, by full-time/part-time status and graduation cohort | 9-10 |
|-------|--|------|
| 9-19 | Satisfaction with principal nursing position, by graduation cohort | 9-11 |
| 9-20 | Intentions regarding nursing employment, by graduation cohort | 9-11 |
| 9-21 | Job changes between March 2007 and March 2008, for registered nurses employed in 2008, by graduation cohort | 9-12 |
| 9-22 | Reasons for job changes, by graduation cohort | 9-12 |
| 9-23 | Employment setting changes for registered nurses who graduated 2001–2008 | 9-13 |
| 9-24 | Secondary nursing positions, by graduation cohort | 9-13 |
| 9-25 | Average earnings from principal nursing position, and from all nursing positions, by graduation cohort | 9-14 |
| 10-1 | Demographic and educational characteristics of registered nurses, by age | 10-2 |
| 10-2 | Employment settings of registered nurses in their principal nursing position, by age group | 10-4 |
| 10-3 | Job titles of registered nurses in their principal nursing position, by age group | 10-4 |
| 10-4 | Job satisfaction, by principal employment setting | 10-5 |
| 10-5 | Average hours worked per week for registered nurses in their principal nursing position, by age | 10-5 |
| 10-6 | Percent of employed registered nurses with secondary nursing positions, by age | 10-0 |
| 10-7 | Average earnings from primary nursing employment, by age | 10-0 |
| 10-8 | Registered nurses employed in nursing in 2007 who changed position or employer, by age | 10-7 |
| 10-9 | Reasons for job changes, by age group | 10-7 |
| 10-10 | Intentions to change employers or leave the nursing profession, by age group | 10-8 |

<u>Tables</u>

| 10-11 | Length of time since last employed in nursing, for registered nurses not employed in nursing in 2008, by age group | 1(|
|------------|--|----|
| 10-12 | Reasons for not working in nursing, by age group | 1(|
| 10-13 | Percent of registered nurses not employed in nursing who work in non-nursing positions, by age group | 1(|
| Appendix A | A Tables | |
| 1 | Gender, racial/ethnic background, and age group, by employment status: 2008 | А |
| 2 | Year of graduation from initial nursing education and the average age at graduation, by type of initial nurse education: 2008 | А |
| 3 | Financial resources used for tuition and fees for initial nursing education, by type of initial nursing education: 2008 | А |
| 4 | Characteristics of registered nurses with a degree before entering initial nursing education, by type of initial nursing education: 2008 | А |
| 5 | Health occupations prior to initial nursing education, by type of initial nursing education: 2008 | А |
| 6 | Registered nurses who were ever licensed as practical/vocational nurse, by type of initial nursing education: 2008 | А |
| 7 | State of location of registered nurses as of March 2008 and State of graduation, by type of initial nursing education and number of years since graduation: 2008 | А |
| 8 | Highest nursing or nursing-related education, by initial nursing education: 2008 | А |
| 9 | Primary focus of post-nursing master's and doctorate degrees: 2008 | А |
| 10 | Current enrollment in nursing or nursing-related education programs, by employment status and student status: 2008 | А |
| 11 | National nursing certifications for registered nurses: 2008 | А |
| 12 | Employment setting in principal nursing position: 2008 | А |

| Appendix | <u>A Tables</u> |
|----------|--|
| 13 | Employment setting in principal nursing position, by age group: 2008 |
| 14 | Employment setting in principal nursing position, by highest nursing or nursing-related education: 2008 |
| 15 | Job title in principal nursing position: 2008 |
| 16 | Job title in principal nursing position, by hospital and nonhospital setting: 2008 |
| 17 | Employment setting in principal nursing position, by selected job titles: 2008 |
| 18 | Job title in principal nursing position, by highest nursing or nursing-related education: 2008 |
| 19 | Employment status and average total earnings for registered nurses with one and more than one nursing position: 2008 |
| 20 | Average annual hours worked and employment setting in principal nursing position: 2008 |
| 21 | Average hours per week and employment setting for registered nurses working full-time in their principal nursing position: 2008 |
| 22 | Average hours per week and employment setting for registered nurses working part-time in their principal nursing position: 2008 |
| 23 | Level of care by type of clinical specialty in principal nursing position: 2008 |
| 24 | Clinical specialty by patient population in principal nursing position for registered nurses who provide patient care: 2008 |
| 25 | Level of care by clinical specialty in principal nursing position for registered nurses who provide patient care in hospital settings: 2008 |
| 26 | Clinical specialty by patient population in principal nursing position for registered nurses who provide patient care in hospital settings: 2008 |
| 27 | Percent of time in each functional area during usual work week in principal nursing position: 2008 |

| <u>Appendix</u> | Appendix A Tables | |
|-----------------|--|------|
| 28 | Average percent of time in each functional area during usual work week in principal nursing position, by highest nursing or nursing-related education: 2008 | A-29 |
| 29 | Employment setting in principal nursing position, by employment relationship: 2008 | A-30 |
| 30 | Average annual earnings for registered nurses working full-time in their principal nursing position, by job title: 2008 | A-31 |
| 31 | Average annual earnings for registered nurses working full-time in their principal nursing position, by employment setting: 2008 | A-32 |
| 32 | Average annual earnings for registered nurses working full-time in their principal nursing position, by job title and highest nursing or nursing-related education: 2008 | A-33 |
| 33 | Job satisfaction, by employment setting in principal nursing position: 2008 | A-34 |
| 34 | Job satisfaction, by job title in principal nursing position: 2008 | A-35 |
| 35 | Job satisfaction of registered nurses employed in nursing, by age group: 2008 | A-36 |
| 36 | Job satisfaction of registered nurses employed in nursing, by highest nursing or nursing-related education: 2008 | A-37 |
| 37 | Employment status in 2007, by employment status in 2008: 2008 | A-38 |
| 38 | Employment setting in 2007, by employment setting in 2008: 2008 | A-39 |
| 39 | Reasons for employer or position change among registered nurses employed in nursing in 2007 and 2008: 2008 | A-40 |
| 40 | Reasons for registered nurses to have occupation other than nursing: 2008 | A-41 |
| 41 | Type of employment of registered nurses who are in non-nursing occupations and not employed in nursing: 2008 | A-42 |
| 42 | Detailed type of employment of registered nurses who are in non-nursing occupations and not employed in nursing: 2008 | A-43 |

Page

43 Job title of registered nurses who are in non-nursing occupations and not employed in nursing: 2008..... A-44 44 Advanced practice registered nurses, by employment status: 2008..... A-45 45 Advanced practice registered nurses and job title, by national certification and State Board recognition: 2008..... A-46 Advanced practice study focus: 2008..... 46 A-47 47 Certifications in advanced practice registered nursing: 2008 A-48 48 Age distribution of registered nurses not employed in nursing, by length of time not working in nursing: 2008..... A-49 49 Age and presence of children, by employment status in principal nursing position: 2008..... A-50 50 Resident State in 2007 and 2008, by age group: 2008..... A-51 51 Registered nurses not employed in nursing and length of time since last worked as a nurse, by whether or not registered nurse was seeking nursing position or had other occupation outside of nursing: 2008..... A-52 52 Registered nurses not employed in nursing and actively seeking nursing employment, by type of employment sought and number of months looking: 2008..... A-53 53 Age and caring for adults at home or others elsewhere, by employment status in principal nursing position: 2008 A-54 54 Marital status and presence of children, by employment status in principal nursing position: 2008 A-55 55 Marital status and caring for adults at home or others elsewhere, by employment status in principal nursing position: 2008 A-56 56 Household income and marital status, by employment status in principal nursing position: 2008 A-57 57 Registered nurses in each State and geographic area, by activity status: 2008 A-58 58 State and geographic area of registered nurses, by employment status in principal nursing position: 2008..... A-60

Appendix A Tables

Appendix A Tables

<u>Page</u>

| 59 | Employment setting of principal nursing position, by geographic area: 2008 A-62 |
|----|--|
| 60 | Average annual earnings for registered nurses working full-time as staff nurses in their principal nursing position, by geographical area: 2008 A-63 |
| 61 | Registered nurses employed in nursing in each State and geographic area, by highest nursing or nursing-related education: 2008 A-64 |
| 62 | Age distribution and average age of registered nurses, by geographic area for all registered nurses and registered nurses employed in nursing: 2008 |
| 63 | Racial/ethnic background of registered nurses, by geographic area: 2008 A-66 |
| 64 | Geographic location of registered nurses, by employment status: 2008 |
| 65 | Predominant countries where internationally educated nurses received initial nursing education, by employment status: 2008 A-68 |
| 66 | Predominant States employing internationally educated nurses (IENs): 2008 A-69 |
| 67 | Employment setting of internationally educated nurses (IENs), by selected job titles: 2008 A-70 |

Appendix B Tables

| B1-1 | Sample information for States where the strata were based only on Alaska, Hawaii, and New Mexico (sorted by frame size) | B-4 |
|------|--|------|
| B1-2 | Sample information for States where strata were based only on Alaska, Hawaii, and New Mexico (sorted alphabetically) | B-5 |
| B1-3 | Sampling States where strata are based on RNs licensed in neighboring States as well | B-6 |
| B2-1 | Response rates, unweighted and weighted by State and for the United States as a whole | B-12 |

Appendix B Tables

<u>Page</u>

| B3-1 | Distribution of imputation rates | B-17 |
|------|--|------|
| B4-1 | Standard errors for select estimates | B-27 |
| B5-1 | Evaluating potential for reduction in bias: Comparing final 2008 estimates with 2008 estimates using a weight that does not account for differential response rates by age | B-34 |
| B5-2 | Comparing 2008 NSSRN and 2008 American Community Survey (ACS) estimates on racial/ethnic background, and age group | B-38 |

Figures

| 1-1 | The nursing workforce of the United States, 1980-2008 | xxxviii |
|-----|---|---------|
| 2-1 | Distribution of registered nurses according to initial nursing education, 1980-2008 | 2-2 |
| 2-2 | Average age at graduation from initial nursing education, by type of program | 2-3 |
| 2-3 | Sources of financing for initial education for all U.S. registered nurses | 2-4 |
| 2-4 | Highest nursing or nursing-related education preparation, 1980- 2008 | 2-8 |
| 2-5 | Highest education of registered nurses, by initial nursing education | 2-10 |
| 3-1 | The nursing population and workforce in the United States, 1980-2008 | 3-2 |
| 3-2 | Employment status of registered nurses | 3-3 |
| 3-3 | Age distribution of the registered nurse population, 1980-2008 | 3-4 |
| 3-4 | Employment of registered nurses, by age group | 3-5 |
| 3-5 | Employment rates, by initial nursing education | 3-6 |
| 3-6 | Employment rates, by highest nursing or nursing-related education | 3-7 |
| 3-7 | Employment settings of registered nurses | 3-9 |

Page

3-10

Types of hospitals in which registered nurses employed in hospital settings work.....

Figures

3-8

| 3-9 | Employment settings of registered nurses, 1980-2008 | 3-12 |
|------|---|------|
| 3-10 | Employment settings, by highest nursing or nursing-related education | 3-14 |
| 3-11 | Registered nurses holding multiple positions, by primary employment status | 3-18 |
| 3-12 | Months worked per year in principal nursing position | 3-21 |
| 3-13 | Hours worked per year in all nursing positions | 3-22 |
| 3-14 | Hours worked per week in principal nursing position | 3-23 |
| 3-15 | Time spent by registered nurses in various activities | 3-33 |
| 3-16 | Time spent by registered nurses in various activities, by job title | 3-35 |
| 3-17 | Actual and inflation-adjusted "real" average earnings for full-time registered nurses in their principal nursing position | 3-39 |
| 3-18 | Job satisfaction of registered nurses | 3-44 |
| 3-19 | Job satisfaction, by employment setting of principal nursing position | 3-46 |
| 3-20 | Job satisfaction, by job title in principal nursing position | 3-47 |
| 3-21 | Job satisfaction of registered nurses employed in nursing, by age | 3-49 |
| 3-22 | Job satisfaction of registered nurses employed in nursing, by highest nursing or nursing-related education | 3-50 |
| 3-23 | Comparison of employment status of registered nurse population in 2007 and 2008 | 3-52 |
| 3-24 | Intentions to stay in nursing over the next 3 years, by age group | 3-54 |
| 3-25 | Non-nursing positions among registered nurses not employed in nursing, by years since last worked in nursing | 3-56 |
| 3-26 | Employment settings of registered nurses employed outside of nursing | 3-57 |

Figures

| 3-27 | Job titles of registered nurses employed outside of nursing | 3-58 |
|------|--|------|
| 4-1 | Age distribution of registered nurses who work as faculty | 4-3 |
| 4-2 | Highest education of registered nurses who work as faculty | 4-4 |
| 4-3 | Highest education of registered nurses who work as faculty in their principal position, by type of program in which they work | 4-5 |
| 4-4 | Satisfaction of registered nurses who work as faculty in their principal nursing position | 4-7 |
| 5-1 | Registered nurses prepared for advanced practice, 2004-2008 | 5-2 |
| 5-2 | Registered nurses prepared in advanced practice nursing | 5-4 |
| 5-3 | Age distribution of registered nurses prepared as nurse practitioners | 5-5 |
| 5-4 | Nurse practitioner preparation, by highest nursing or nursing- related degree | 5-6 |
| 5-5 | Hours worked per week by employed nurse practitioners in their principal nursing position | 5-7 |
| 5-6 | Job titles of nurse practitioners | 5-8 |
| 5-7 | Employment settings in which nurse practitioners work | 5-11 |
| 5-8 | Percent of nurse practitioners whose primary level of care is ambulatory or primary care, by employment setting | 5-12 |
| 5-9 | Nurse practitioner satisfaction with principal position | 5-13 |
| 5-10 | Age distribution of registered nurses prepared as nurse midwives | 5-15 |
| 5-11 | Nurse midwife preparation, by highest nursing or nursing-related degree | 5-16 |
| 5-12 | Hours worked per week by employed nurse midwives in their principal nursing position | 5-17 |
| 5-13 | Job titles of nurse midwives | 5-18 |
| 5-14 | Employment settings in which nurse midwives work | 5-19 |

| <u>Figures</u> | | <u>Page</u> |
|----------------|---|-------------|
| 5-15 | Nurse midwife satisfaction with principal position | 5-20 |
| 5-16 | Age distribution of registered nurses prepared as nurse anesthetists | 5-22 |
| 5-17 | Nurse anesthetist preparation, by highest nursing or nursing- related degree | 5-23 |
| 5-18 | Hours worked per week by employed nurse anesthetists in their principal nursing position | 5-24 |
| 5-19 | Nurse anesthetists satisfaction with principal position | 5-25 |
| 5-20 | Age distribution of registered nurses prepared as clinical nurse specialists | 5-28 |
| 5-21 | Clinical nurse specialist preparation, by highest nursing or nursing-related degree | 5-29 |
| 5-22 | Hours worked per week by employed clinical nurse specialists in their principal nursing position | 5-30 |
| 5-23 | Job titles of employed nurses prepared as clinical nurse specialists | 5-31 |
| 5-24 | Job titles of employed nurses dual-prepared as clinical nurse specialists and nurse practitioners | 5-32 |
| 5-25 | Employment settings in which clinical nurse specialists work | 5-33 |
| 5-26 | Clinical nurse specialist satisfaction with principal position | 5-34 |
| 6-1 | Age distribution of nurses who are not working in nursing | 6-2 |
| 6-2 | Percentage of nurses not working in nursing, by age and presence of children | 6-3 |
| 6-3 | Length of time since nonworking nurses last worked in nursing, by age | 6-4 |
| 6-4 | Reasons nurses are not working in nursing, by age group | 6-5 |
| 6-5 | Percent of registered nurses seeking nursing employment, by number of years since last worked in nursing and age | 6-7 |
| 7-1 | Percent of registered nurses working in nursing, by age cohort, 1988-2008 | 7-2 |

Page

| 7-2 | Percent of men employed in nursing, by job title | 7-4 |
|------|---|-----|
| 7-3 | Percent of registered nurses who are White or non-White | 7-5 |
| 7-4 | Racial/ethnic distribution, by year of graduation from initial nursing education | 7-6 |
| 7-5 | Distribution of registered nurses and the U.S. population, by racial/ethnic background | 7-7 |
| 7-6 | Distribution of employment settings, by geographic region | 7-1 |
| 7-7 | Median earnings for staff and management/administration registered nurses, by geographic region | 7-1 |
| 7-8 | Percent of employed registered nurses with a bachelor's degree and higher, by region | 7-1 |
| 7-9 | Percent of employed registered nurses in racial/ethnic minority groups, by geographic region | 7-1 |
| 8-1 | Countries of origin of internationally educated nurses in the United States | 8-2 |
| 8-2 | Secondary nursing positions held by IENs working full-time in their principal nursing position | 8-0 |
| 10-1 | Percent of employed registered nurses working full-time and part-time, by age group | 10 |

For citation please use: U.S. Department of Health and Human Services, Health Resources and Services Administration (2010). *The Registered Nurse Population: Findings from the 2008 National Sample Survey of Registered Nurses.*

For an electronic copy of the report and further information: <u>http://bhpr.hrsa.gov/healthworkforce/rnsurvey</u>

Figures

Executive Summary

Purpose of the National Sample Survey of Registered Nurses

The National Sample Survey of Registered Nurses (NSSRN), which has been conducted since 1977, is the largest survey of registered nurses (RNs) in the United States. Nurses with active RN licenses are asked to report on their education, employment, intentions regarding nursing work, and demographics. These data are used by policymakers and numerous stakeholders, both domestic and international, to assess the status of and trends in the U.S. nursing workforce.

The Size of the Registered Nurse Workforce

There were an estimated 3,063,162 licensed registered nurses living in the United States, as of March 2008. This is an increase of 5.3 percent from March 2004, representing a net growth of 153,806 RNs. An estimated 444,668 RNs received their first U.S. license from 2004 through 2008, and thus approximately 291,000 RNs allowed their U.S. licenses to lapse, possibly indicating that the substantial retirements that have been anticipated may have begun.

Initial Education of Registered Nurses

The most commonly reported initial nursing education of RNs in the United States is the Associate Degree in Nursing (ADN), representing 45.4 percent of nurses. Bachelor's or graduate degrees were received by 34.2 percent of RNs, and 20.4 percent received their initial education in hospital-based diploma programs. The average age of RNs who graduated from their initial nursing education program is rising. More than 21 percent of RNs earned an academic degree prior to their initial nursing degree. Nearly two-thirds of RNs reported working in a health occupation prior to their initial nursing education.

Executive Summary

Highest Education Preparation of Registered Nurses

After initial licensure, RNs can obtain additional academic degrees in nursing or nursing-related fields. In 2008, half of the RN population had a bachelor's or higher degree in nursing or a nursing-related field, while the other half's highest education level was a diploma or an ADN.

Employment of Registered Nurses

In 2008, an estimated 2,596,399 RNs were employed in nursing, representing 84.8 percent of licensed RNs. This was the highest rate of nursing employment since the NSSRN commenced in 1977. There also has been the first increase in full-time employment since 1996, rising from 58.4 percent of RNs in 2004 to 63.2 percent in 2008. Among nurses under 50 years old, 90 percent or more are employed in nursing positions; this percentage drops to less than half of RNs over age 65.

Hospitals remain the most common employment setting for RNs in the United States, increasing from 57.4 percent in 2004 to 62.2 percent of employed RNs in 2008. The increase in this percentage is the first increase since 1984. While nearly 90 percent of RNs under 25 years old work in hospitals, less than 53 percent of RNs age 55 and older work in hospitals.

Fewer than half of nurses with master's degrees work in hospitals; more than 18 percent are in ambulatory care settings and nearly 12 percent are in academic education.

The most common job title of RNs in the United States is "staff nurse," or its equivalent (66.3 percent). Between 2004 and 2008, the proportion of staff RNs increased by 2.2 percent. Just under 20 percent of RNs with graduate degrees are staff RNs, compared with 72.8 percent of those without a graduate degree. The next most common job title in 2008 included management and administration titles (12.5 percent).

Many registered nurses hold more than one job in nursing. Overall, about 12 percent of RNs who have a full-time primary nursing position and 14 percent of those with part-time primary positions have additional nursing positions.

More than half of RNs work at least 40 hours per week in their principal nursing position, and another 24.2 percent work 32 to 39 hours per week. A total of 19.1 percent of RNs report that they

worked on-call or could have been called to work (on "standby") during a typical work week in their principal nursing positions. Among RNs employed in nursing positions, 27.5 percent report that they worked overtime averaging 7.5 hours per week and received pay for such work.

Earnings and Satisfaction of Registered Nurses

Average annual earnings for RNs employed full-time in 2008 were \$66,973, rising 15.9 percent since the 2004 average of \$57,785. When annual earnings are adjusted for inflation to 1980 dollars, earnings in 2008 increased only 1.7 percent from average inflation-adjusted 2004 earnings. The highest earnings were reported by nurse anesthetists (NAs), who averaged \$135,776 per year. Staff nurses earned \$61,706 per year, on average. Registered nurses with graduate degrees earned an average of at least \$20,000 more than did RNs with other levels of education.

In 2008, 29.3 percent of RNs reported that they were extremely satisfied and a further 51.8 percent reported being moderately satisfied with their principal nursing position. This compares with 2004 rates of satisfaction of 27.5 percent and 50.5 percent, respectively. Only 11.1 percent were dissatisfied in 2008, as compared with 13.8 percent in 2004. The highest rates of being moderately or extremely satisfied were reported by registered nurses working in academic education programs (86.6 percent), while the lowest rate of being either moderately or extremely satisfied was reported by nurses in nursing homes/extended care (74.5 percent).

Job Changes and Future Employment Plans

Nearly 80 percent of RNs were employed in nursing in both 2007 and 2008, while 12.3 percent were not employed in nursing in either year. Nearly 3 percent were employed in nursing in 2007, but not in 2008, and conversely, 2.4 percent returned to employment in 2008. Of those who were employed in 2007 but not in 2008, 27.3 percent (24,430) reported that they stopped working due to retirement. Among nurses who worked full-time in 2007, 11.6 percent changed to a different employer by 2008. Another 6.5 percent of these RNs worked with the same employer, but held a different position. More than 73 percent of RNs reported that they changed positions or employers due at least in part to workplace issues, such as lack of good management or inadequate staffing. Personal career reasons, such as interest in another position or improved pay or benefits, were a factor in job

changes for 37.5 percent of nurses. Nearly 30 percent of nurses changed jobs at least in part due to personal family reasons.

The percentage of employed RNs who intended to leave nursing within 3 years is quite small, at 3 percent or fewer, among nurses under 55 years old. Among RNs 55 and older who worked in nursing in 2008, 12.5 percent intended to leave the nursing profession within 3 years and another 8.9 percent intended to leave their current nursing jobs and were unsure if they will remain in nursing afterward.

Nurse Faculty

There were an estimated 31,056 RNs working as faculty in their principal nursing position in 2008. The majority (60 percent) of these worked in programs that provide education at the Bachelor's of Science in Nursing (BSN) level or a higher level. Another 38.3 percent worked in ADN education programs. An additional 34,666 RNs worked in academic education settings as a secondary nursing position.

In 2008, more than 40 percent of RNs who worked as faculty in their principal nursing position were between the ages of 50 and 59, and more than 19 percent of RNs whose principal position is as faculty were 60 years or older. Nursing faculty received somewhat lower annual earnings than RNs in general from their principal full-time nursing position. Earnings from principal nursing positions for RNs who work full-time averaged \$66,973 in 2008, while earnings for faculty RNs were \$63,985 on average. Nearly 90 percent of faculty reported that they are moderately or extremely satisfied with their position, as compared with 81.1 percent of all working RNs. The satisfaction of faculty is similar to that of nurses with graduate degrees, of whom 87.3 percent were moderately or extremely satisfied.

Nurse Practitioners

In 2008, an estimated 158,348 nurses had preparation as nurse practitioners (NPs). NPs represent the largest group of advanced practiced registered nurses (APRNs). About 35 percent of NPs were under 45 years old. Nearly 85 percent of NPs reported that they held a master's degree and 3.9 percent reported holding a doctorate degree. Of all NPs, 10.3 percent also were prepared as clinical nurse specialists (CNSs). More than 93 percent of NPs employed in nursing reported having either national certification or State Board recognition.

Most NPs were employed in nursing (89.2 percent), and 69.6 percent of these NPs had a principal position job title of NP. Virtually all NPs who reported NP as their job title had national certification or State Board recognition. Some NPs (11.4 percent) reported that their job title was staff nurse or an equivalent; others worked in instruction or had a management or administration title. More than 38 percent of NPs were employed in hospital settings, including primary care clinics located in or owned by a hospital. Another 35.3 percent worked in ambulatory care settings outside hospitals.

Nurses prepared as NPs in 2008 earned an average of \$73,776 per year in their principal nursing position. More than 87 percent of NPs were moderately or extremely satisfied with their principal position.

Nurse Midwives

Nurse midwives (NMs) were the smallest of the APRN groups, with an estimated 18,492 nurses prepared as NMs. More than 54 percent were 50 years or older, and half (50.3 percent) reported having a graduate degree. More than 91 percent of NMs employed in nursing had national certification or recognition from their State Board of Nursing.

The majority (84.3 percent) of NMs were employed in nursing positions in 2008, but only 41.7 percent of employed NMs reported a job title of nurse midwife. All of these NMs also had State Board recognition or national certification in this field. The majority (57.9 percent) of NMs worked in hospital settings, and 25.1 percent worked in ambulatory care.

Nurses prepared as NMs earned an average of \$69,222 per year. They had a high level of satisfaction with their principal nursing position, with 86.2 percent being moderately or extremely satisfied.

Executive Summary

Nurse Anesthetists

NAs were the third largest group of APRNs, with an estimated 34,821 nurses prepared as anesthetists in 2008. NAs were also the youngest group of APRNs: nearly 40 percent were under 45 years old. Nearly 40 percent of NAs were men. In 2008, 65.4 percent of NAs held a master's or doctorate degree as their highest nursing or nursing-related degree, and almost all had national certification or State Board recognition (99.3 percent).

In 2008, 31,868 NAs were employed in nursing positions (91.5 percent). Registered nurses prepared as NAs earned an average of \$135,776 per year, making them the highest-paid APRN group. They were the most satisfied of the APRN groups, with 93.5 percent reporting moderate or extreme satisfaction with their principal position.

Clinical Nurse Specialists

In 2008, there were an estimated 59,242 nurses prepared as CNSs. CNSs were the second largest group of APRNs. There was a marked decline of 18.3 percent in the number of CNSs between 2004 and 2008. CNSs were older than other APRNs in 2008, with 63.6 percent over 50 years old. The vast majority of CNSs (92.8 percent) reported at least a master's degree as their highest preparation, with 7.2 percent completing a doctorate degree for their CNS preparation. More than 50 percent had national certification or recognition from their State Board of Nursing. CNSs had the lowest rate of employment in nursing of all the advanced specialties, with 84 percent working in nursing. Only 18.8 percent of CNSs reported their job title as clinical nurse specialist. Other common job titles among nurses prepared as CNSs are instruction (21.1 percent), and management or administration (17.8 percent). Nearly half of nurses prepared as CNSs worked in hospitals (49.4 percent).

Some nurses prepared as CNSs also had preparation in other advanced specialties. More than 27 percent of CNSs also had preparation as NPs. Among such dual-prepared RNs the most common job title was nurse practitioner (54.8 percent).

Nurses prepared as CNSs in 2008 earned an average of \$74,918 per year in their principal nursing position. CNSs were very satisfied with their principal positions, with 91 percent being moderately or extremely satisfied.

Nurses Who Do Not Work in Nursing

An estimated 466,564 RNs were not employed in nursing in 2008. Nearly one-third of RNs who were not working in nursing were 65 years of age or older; almost half were 60 years of age or older. The decision to work in nursing is more closely tied to age of registered nurses' children than presence of them; RNs whose children were all under age 6 were less likely to work in nursing than were RNs whose children were all at least 6 years old.

Overall, about half of RNs who were not working in nursing in 2008 have been out of nursing work for 4 or fewer years. Among nurses who were under 50 years old, 57.5 percent have been out of nursing for less than 4 years. Reasons for not working in nursing are often associated with life stages, such as the need to care for children or retirement. However, 41 percent of RNs under 50 years old reported that they were not working in nursing due to workplace problems such as burnout, stressful work environment, and poor management. This is of concern. Among nurses 50 years and older, personal and family reasons (41.5 percent), workplace reasons (35.5 percent), and personal career factors (32.4 percent) also were important. Among RNs under 50 years old who were not working in nursing, only about 15 percent did not intend to return to nursing work.

Demographics and Diversity

For the first time in the past three decades, the aging of the nursing workforce slowed. In 1988, half of the working population of RNs was less than 38 years of age, but in 2004 that figure rose to 46 years and remained steady at 46 years of age in 2008. This slowdown in the aging trend resulted from an increase in employed RNs less than 30 years of age—the first increase seen in this age group since the inaugural NSSRN in 1977.

Men comprised a growing percent of the RNs in 2008, making up 6.2 percent of employed RNs who were licensed before 2000 and 9.6 percent of those licensed in 2000 or more recently. The employment profile of men who work in nursing differs from that of women. More men reported principal employment in a hospital and work as NAs.

Significant gains in racial/ethnic minority representation among RNs have been made over time. The population of RNs from minority racial/ethnic groups increased from 333,368 in 2000 to 513,860 in 2008, with the growth occurring primarily among recently graduated RNs. However, the racial and ethnic distribution of the RN population is substantially different from that of the U.S. population. Approximately 65 percent of the U.S. population is White, non-Hispanic, while 83.2 percent of RNs are White, non-Hispanic.

Internationally Educated Nurses

Internationally educated nurses (IENs) are RNs whose initial nursing education took place outside of the United States or in the U.S. territories (formerly termed foreign-trained or foreign-educated nurses). They represent a larger percentage of the U.S. nursing workforce in recent years, comprising 5.1 percent of RNs licensed before 2004, compared with 8.1 percent since then. The Philippines continued to dominate as the source country of the IEN workforce (50 percent), followed by Canada at nearly 12 percent. India supplied 9.6 percent and the United Kingdom provided 6 percent, with the contributions from India accelerating and surpassing those from the United Kingdom among recent licensees. Approximately one-quarter of IENs lived in California in 2008, with New York, Florida, and Texas each home to 10 to 12 percent of IENs.

Most IENs (88.3 percent) were employed in nursing in 2008, a somewhat higher percentage than RNs in general (84.8 percent). IENs were more likely than their U.S.-educated counterparts to work full-time. A greater percentage (17.2 percent) of internationally educated RNs hold secondary nursing positions than do RNs in general. Approximately three-quarters of IENs licensed since 2004 are employed in hospital settings. Most IENs licensed since 2004 (94.1 percent) have staff nurse job titles.

Recent Nurse Graduates

In this report, recent graduates are defined as those RNs who completed their initial nursing education after 2000 (from 2001 to 2008). The most recent graduates are those who completed their initial nursing education after 2004 (2005 to 2008). Recent graduates comprised nearly 20 percent of all RNs in the United States in 2008 and about 23 percent of employed RNs.

About 39 percent of recent graduates entered the profession with a BSN; only 3 percent had a diploma nursing education. RNs who graduated before 2001 completed their nursing programs at an average age of 26.7 years, while the average age of the recent graduates was a little over 30 years.

RNs among the most recent graduate cohort were more likely than RNs who graduated in any previous cohort to have received a postsecondary degree prior to entering nursing; 31.7 percent of graduates from 2001 to 2004 had a prior postsecondary degree, compared with 37.1 percent of graduates from 2005 to 2008.

More than 10 percent of RNs who completed their initial nursing education between 2001 and 2004 later completed additional degrees. Bachelor's degrees were completed by nearly 10 percent of graduates from diploma or ADN programs; master's or doctorates were completed by more than 10 percent of BSN graduates and almost 2 percent of diploma or ADN graduates.

The vast majority of recent graduates were employed in nursing positions in 2008 (94.5 percent of 2001–2004 graduates and 96.6 percent of 2005–2008 graduates), and 83.8 percent of the most recent graduates worked full-time.

Hospitals employed more than 83 percent of the most recent graduates and more than 75 percent of RNs who graduated in 2001 to 2004. The most common job title held by all working RNs was staff nurse or an equivalent title, with 92.2 percent of the most recent graduates and 80.7 percent of graduates from 2001 to 2004 having this title.

Recent graduates were less satisfied with their principal nursing positions than were RNs who graduated before 2001.

In addition, recent graduates change jobs somewhat more frequently than RNs who graduated prior to 2001. Of all employed RNs, 14.9 percent of graduates from 2001 to 2004 had changed employers for their principal nursing position in the previous year and 7.5 percent had changed positions but were with the same employer. Among the 2005 to 2008 graduates, the percentages for job changing were similar. The reasons for job changes varied, but those cited most frequently were related to working conditions. Recent graduates were more likely to report that they changed jobs for career reasons than were pre-2001 graduates. Recent graduates also were more likely than pre-2001 graduates to report that personal or family reasons drove their job change.

Nearly 40 percent of recent graduates planned to leave their current jobs within 3 years, compared with 27.1 percent of pre-2001 graduates. Nonetheless, the vast majority of recent graduates intended to remain in the nursing profession, with only 3.3 percent of those who intended to leave their current job stating that they did not plan to stay in nursing.

Average earnings from all nursing positions for the most recent graduates were \$52,994. The total nursing earnings of the most recent graduates were lower on average than the average total earnings of RNs who graduated from 2001 to 2004 (\$57,489) and lower than the average total earnings for RNs graduating before 2001 (\$62,903).

Nurses Nearing Retirement

Older RNs over age 50 comprised 44.7 percent of the total RN population in 2008, compared with 33 percent in 2000. The percentage of older RNs working in nursing in 2008 declines steadily from 87.5 percent for those 50–54 years of age to 33.3 percent for those aged 75 and older. RNs aged 50–54 in 2008 who were employed full-time worked an average of 43.7 hours per week, slightly more than the average for full-time nurses under age 50. Beginning at age 60, the hours worked by part-time nurses declines steadily with age. Older RNs are also less likely than younger RNs to have secondary positions in nursing.

While a smaller percentage of older nurses worked in hospital positions in 2008, higher percentages are employed in nursing homes/extended care, academic education programs, and home health settings. The majority of RNs in 2008 remained employed with the same employer, and older RNs were more likely to be in the same position and with the same employer year to year when compared with younger nurses. Registered nurses 50 to 59 years old were less likely to report an intention to leave their current nursing position within 3 years than were RNs under 50 years old. As RNs grow older, and especially for RNs older than age 60, retirement becomes an increasingly dominant reason for employment changes.

Introduction

Conducted since 1977, the National Sample Survey of Registered Nurses (NSSRN) is the largest survey of the Registered Nurses (RNs) workforce in the United States. A sample of nurses with active RN licenses from every State are asked to report on their education, employment, intentions regarding nursing work, and demographics. The resulting data are used by policymakers and numerous stakeholders, both domestic and international, to assess status and trends in the U.S. nursing workforce. The data are also used to evaluate the responsiveness of the workforce to Federal and State programs seeking to ensure workforce capacity for providing essential health care services and identifying areas for further research. These data illustrate education and skills of the RN workforce, portray the diversity of RNs, illustrate the U.S. employment of internationally educated nurses, depict the factors that affect RNs' decisions to work in the nursing profession, and support efforts to forecast the future supply of and demand for RNs. After the first survey was conducted in 1977, the NSSRN has been conducted every 4 years since 1980. This report provides findings from the survey conducted in 2008.

The Registered Nurse Workforce in the United States

There were an estimated 3,063,162 licensed RNs living in the United States, as of March 2008 (Figure 1-1 and Appendix A, Table 1).¹ This is an increase of 5.3 percent from March 2004, representing a net growth of 153,806 RNs.² Based on data from the results of the 2008 survey, an estimated 444,668 RNs received their first U.S. license from 2004 through 2008, and thus approximately 291,000 RNs had a lapsed RN license during this time period. These data indicate that the long anticipated and substantial retirements from the nursing field may have begun. Thus, even as new graduates enter the field, overall nursing workforce growth may be affected by RNs leaving the profession.

¹ The standard error for the estimate of the total number of RNs licensed and living in the United States is 4,045.8. The 95 percent confidence interval has a lower limit of 3,055,232 and an upper limit of 3,071,092. Please see Chapter 1 and Appendix B for more details on weighting and estimation.

² There were an additional estimated 11,903 internationally educated RNs holding active U.S. licenses who were not living in the United States as of March 2008. RNs living outside of the Unites States are not included in the analyses of the U.S. nursing workforce for the purposes of this report.




Figure 1-1.The nursing workforce of the United States, 1980-2008

Source: 1980-2008 National Sample Survey of Registered Nurses

1.1 Summary

The 2008 National Sample Survey of Registered Nurses (NSSRN) was based on a randomly selected sample of individual RNs from each State's listing of licensed RNs. The sampling rate differed by State. Sampling for all previous administrations of NSSRN was based on a randomly selected sample of alphabetic clusters of names in each State using a "nested alpha segment design." When nested alpha-segment design was used for the first administration of the NSSRN in 1977, available data sources for identifying eligible RN listings could be on paper listings, index cards, electronic files, or other media. The alpha-segment design provided an innovative way of determining an RN's probability of selection for the survey, in spite of the data source limitations, especially for those RNs licensed in multiple States, without having to identify a given RN on all the lists on which that individual appeared.

Technological advances in computer hardware and software since earlier NSSRN surveys have made identifying multiple appearances of the same RN on different lists feasible. The sample design for the 2008 NSSRN takes advantage of such technological advances. In so doing, the precision of the national estimates has improved. The 2008 NSSRN included a sample of 55,151 RN records. From this sample, 33,549 surveys were completed, yielding an unweighted response rate¹ of 62.41 percent.

Data were collected between July 2008 and March 2009 using mailed paper surveys, electronic surveys on the World Wide Web (the Web), and direct telephone interviews. Each survey was carefully reviewed for missing or inaccurate data. When critical data were found to be missing, incomplete, or insufficient, professional telephone interviewers called the respondent to ask about the data item(s) in question. When complete data could not be obtained from the sampled RN, appropriate statistical imputation methods were used to reduce nonresponse error. The 2008 NSSRN imputation protocol used regression modeling to generate likely responses for missing values for individual questions within completed cases. Imputation of this type had not been

¹ The American Association for Public Opinion Research Response Rate #3 calculation was used; it is defined as the number of completed surveys divided by the number of returned surveys (complete plus partial) plus an estimate of the number of eligible cases among those who did not respond and whose eligibility could not be ascertained.

undertaken for any previous administration of the NSSRN but is a powerful statistical method using defined protocols.

The establishment of sample weights for the participating RNs was the final step in preparing the data for analysis and reporting. The weighting process for the 2008 NSSRN had several steps, which included identifying the initial probability of selection for each sampled record; adjusting for survey nonresponse; and accounting for multiple chances of selection for RNs who appeared on the listings of more than one State. In addition, a weight calibration procedure known as "raking" was employed to achieve consistency between the counts appearing on each State listing and the estimates of the number of RNs currently licensed in each State, based on the survey weight assignments. Weighted responses were then used to calculate national estimates.

Also different from prior NSSRN survey analyses, the nonresponse adjustment procedure used for the 2008 NSSRN used weighting to adjust for differences in rates of response to the survey by different age groups. Response rates by age group varied; generally older RNs participated at a higher rate than younger RNs. Without a weighted adjustment for age group response rates, older RNs would be overrepresented in population estimates, leading to a disproportionately large effect on survey estimates associated with age. Thus, adjusting sample weights to reflect age group participation rate served to reduce the potential for bias in some NSSRN estimates.

This chapter provides an overview of the survey methods used in the 2008 NSSRN process. Appendix B provides a more detailed description of the sample allocation, sample performance, imputation, weighing methods, and comparisons with previous survey sampling and weighting techniques. NSSRN data users who are interested in comparing the 2008 NSSRN findings with findings from previous NSSRN administrations should pay particular attention to the discussions of imputation and weighting here and in Appendix B.

The data reported in tables in the body of the report may differ slightly from data in the correspondingly cited appendix. This may be due to several factors such as the rounding of aggregate data or the context of the discussion where there is some difference in the population of interest.

1.2 2008 Questionnaire

The content and structure of the 2008 questionnaire were very similar to the questionnaire used for the 2004 NSSRN administration. A copy of the 2008 questionnaire is included in Appendix C. The following were key questionnaire topics:

- *Eligibility* for the survey;
- *Education*, including initial nursing education, upgrade education post licensure, and education preparation for advanced practice nursing;
- Principal nursing employment, including geographic location, employment setting, job title, hours worked, level of care, patient population, clinical specialty, earnings, and job/employment intentions;
- Secondary employment, including employment settings, hours worked, and earnings;
- Nurses not working in nursing, including their intention to return to nursing and reasons for not working in nursing;
- Employment outside nursing, including area/field of work, hours worked, and earnings;
- *Prior nursing employment,* including reasons for job change, geographic location of prior employment, and prior employment setting;
- *General information* about the responding RN, including job satisfaction, geographic location of current residence and residence 1 year ago, race and ethnicity, language fluency, number of dependents in the household, and household earnings; and
- Licensing and certification information.

Some revisions to the 2008 questionnaire addressed new or emerging topics of interest and improved data quality. Additionally, some questions were converted from "check one" to "check all that apply" to better capture the richness and breadth of information about RNs. Of particular note are the survey additions or revisions in the following areas:

- Employment setting (questions 23 and 57);
- Job title (question 24);
- Level of care (question 29a);
- Patient population served by the RN (question 29b);
- Clinical specialty (question 29c);

- Hours worked per week in their principal nursing position (question 27) and in any secondary nursing positions (question 36); and
- Distribution of time across job functions (question 28).

1.3 Sample Design

From the inception of the NSSRN in 1977 through the 2004 administration, the sample of RNs was identified using a nested cluster design in which RNs were clustered alphabetically by last name: This last name cluster design is referred to as the "alpha-segment design." Although the alpha-segment design produced good estimates of the nursing workforce over the years, it has some shortcomings. First, because RNs of the same racial/ethnic background may have similar alphabetically clustered names, the alpha-segment design may have resulted in larger than desired variation for some estimates, such as racial and ethnic composition of the RN workforce. Second, constructing an alpha-segment frame and implementing the sample design were tedious and time-consuming tasks. Third, sorting to achieve implicit stratification was not feasible. Lastly, the alpha-segment design made adjustments for differential nonresponse a bit more cumbersome.

The 2008 NSSRN sample design is based on independent systematic random samples selected from State-based strata, with equal probability of selection within each stratum. An equal probability systematic random sample can be thought of as randomly selecting a record from the beginning of a list with probability 1/n and then every nth record after that until all records on the list have had a chance of being selected. This kind of design was not feasible in 1976, but technological advances and the advent of electronic recordkeeping have largely eliminated the barriers. The 2008 NSSRN sample design was straightforward to implement and eliminated the clustering that could contribute undesirable variability to survey estimates, particularly for survey results associated with race/ethnicity.

The sample design was developed to address a number of issues:

- A given RN may be licensed in more than one State.
- Unduplication was carried out to eliminate additional records for a single nurse, such as a record associated with advanced practice nursing when an RN already had a basic RN license record. Therefore, for the purposes of sampling, an individual RN in a given State is associated with exactly one RN license.

- Some States (22 at the time of the 2008 administration) participate in what is known as the Nurse Licensure Compact, which affects who may appear on a given State's listing of licensed RNs. An RN licensed in one Compact State would not have a license in other Compact States, but could work as an RN in any of the Compact States.
- Some areas of the country experience significant interstate commuting of RNs, with RNs who live in one State working in another, nearby State or participate as a travel nurse for a short period of time.

Each State (and the District of Columbia) maintains a list of RNs actively licensed in that State. These State lists were used to draw a probability sample of RNs for the NSSRN from among those RNs who were licensed as of the end of 2007/early 2008. RNs appearing in the listings of licensed RNs for more than one State had multiple chances of selection for the 2008 NSSRN. To avoid duplication, multi-State strata were formed for several groups of States where interstate commuting was expected to be fairly sizeable. Probability matching was used to form such strata so that only a single record for those RNs appears in the listing for the multi-State stratum.

After these adjustments, a sample of 55,151 licensed RNs was drawn for the 2008 NSSRN. This sample size was expected to yield sufficient completed surveys to support stable State, regional, and national estimates. Data were collected on the RN population who were actively licensed as of March 10, 2008. Appendix B provides a detailed description of the sample strata, determination of the probabilities of selection, and allocation of the 55,151 RNs across the 50 States and the District of Columbia.

1.4 Data Collection Methods

The 2008 NSSRN employed a multimode approach to data collection. Data were collected from July 2008 through March 2009. Using contact information contained in State licensing records, sampled RNs were first notified of the study via a letter mailed through the U.S. Postal Service (USPS). The letter invited RNs to complete a survey via the Web. Shortly after the Web invitation was sent, paper surveys were also sent via USPS. Finally, RNs who did not respond were contacted by telephone and expedited mail. The different steps of data collection were as follows:

• **First Contact.** A survey notification letter was sent via USPS to all RNs in the sample. The first contact mailing included a description of the study and instructions for logging on and completing the survey via the Web. No paper questionnaire was included in this first contact mailing.

- Second Contact. Two weeks after the first contact mailing, a paper questionnaire was sent via USPS to all RNs who had not completed their survey via the Web. The second contact mailing included a description of the study, a second invitation to complete the survey via the Web, and Web log-on instructions. A paper questionnaire and a postage-paid return envelope for the questionnaire were included to increase response from those RNs without Web access or who preferred paper questionnaires.
- Third Contact. Approximately 3 weeks after the second contact mailing, post cards were sent to all sampled RNs reminding them to complete their survey either on paper or on the Web. A note of thanks was also included on the post card for RNs who had already completed their surveys.
- Fourth Contact. A second copy of the paper questionnaire was sent to nonrespondents 3 weeks after the third contact. The fourth contact mailing was sent only to nonrespondents and included a letter requesting completion of the survey, Web log-on instructions, a paper questionnaire, and a postage-paid return envelope for the questionnaire.
- Fifth Contact. After 3 weeks had passed following the fourth contact, nonresponding RNs were contacted. RNs who failed to respond after the first four contact attempts were split into two groups for follow-up. RNs whose telephone numbers were available from licensing lists were contacted for follow-up by professional telephone interviewers. Services that provide matching and telephone number searches were also used to find a telephone number for a telephone follow-up with nonresponding RNs. If the telephone number search was unsuccessful, RNs who did not respond were sent a final package via FedEx that included a letter requesting completion of the survey, Web log-on instructions, a paper questionnaire, and a postage-paid return envelope for the questionnaire.

When a survey was received that contained responses to all questions deemed necessary to meet the requirement for a completed survey, the survey was noted as "complete" and the survey data were added into the database of 2008 NSSRN complete surveys. Data collection for the 2008 NSSRN ended on March 31, 2009.

1.5 Sample Performance

Response rates are often used to help assess the quality or unbiasedness of survey results. The 2008 NSSRN unweighted response rate was 62.41² percent with a weighted response rate of 61.48 percent.

² The American Association for Public Opinion Research Response Rate #3 calculation was used. It is defined as the number of completed surveys divided by the number of returned surveys (complete plus partial) plus an estimate of the number of eligible cases among those who did not respond and whose eligibility could not be ascertained.

Weighted response rates take into account sample design and probabilities of selection, adjusting for the fact that some States and strata are represented disproportionately in the sample. The weighted response rate for this survey can be interpreted as an estimate of the expected propensity to respond of an RN randomly selected from among all RNs currently licensed in the Nation.

Table 1-1 shows the overall performance of the sample. Table B2-1 in Appendix B provides a distribution of response rates by State.

| Sample disposition | Number in sample |
|---------------------------|------------------|
| Completed surveys | 33,549 |
| Over the Web | 13,066 |
| On paper | 14,897 |
| Over the telephone | 5,586 |
| Ineligible for the survey | 870 |
| Nonresponse | 14,160 |
| Refusals | 6,572 |
| Total sample | 55,151 |

Table 1-1. Summary of 2008 NSSRN sample performance

Source: 2008 National Sample Survey of Registered Nurses

1.6 Imputation

There are two types of nonresponse in surveys: unit nonresponse and item nonresponse. Unit nonresponse occurs when a sampled RN does not complete the survey at all. Unit nonresponse was reflected in the calculation of the response rate and in weighting through nonresponse adjustment.

Item nonresponse occurs when an RN responded to the survey but failed to provide answers to one or more individual questions appropriate for that individual. This results in a missing value for one or more survey items. When items were missing values, the items were filled in from the State sample frame files (licensure lists) when possible. The variables usually available on the sample frames included demographic characteristics (date of birth, race/ethnicity, and gender), home address, and first RN license date. Not all States provided these variables, and data for them could be very sparse even when the variable was included in the State frame.

If any important item nonresponse existed after review of the State sample files, respondents were called and queried about the missing data item(s), a process referred to as data retrieval. Some items deemed important and worthy of initiating data retrieval were year of licensure, employment setting

for principal nursing position, job title, number of hours worked, and other data elements central to the analysis of the 2008 NSSRN data. As a last resort, when item nonresponse persisted after review of the sample frame file and data retrieval, statistical imputation was used to assign values for missing data.

The 2008 NSSRN imputation protocol relied on regression modeling to generate likely responses where there were missing values for individual questions within completed cases. Imputation statistical procedures identify a value that not only would probably be appropriate for the respondent but would also be in keeping with the distribution of values for that variable. For example, the survey items "Nurse's year of graduation from initial nursing education" and "Year of first licensure" are highly correlated. If the year of licensure is missing from an RN's record but not the year of graduation, the missing year of licensure value can be imputed using regression modeling. The year of licensure is the dependent variable in this case, and the year of graduation is the independent (or predictor) variable. The regression model is established using respondent cases in which both the year of licensure and year of graduation are known.

There is a component of variation associated with the generation of imputed values that is generally not reflected in survey estimates of variance. As a result, imputation can result in an understatement of the variance in survey estimates. The degree to which estimates of variability have been understated is generally considered negligible if the imputation rate is low (less than 10 percent). However, if the imputation rate is more than 20 percent, the degree of underestimation may be nontrivial.

Eighty percent of the imputed variables for the 2008 NSSRN have imputation rates below 10 percent, 19 percent are borderline (imputation rate of 10-19 percent), and three variables (1 percent of all survey variables) have an imputation rate greater than 20 percent.³ The three variables are question 11B—total number of hours provided for emergency preparedness training in the past year (imputation rate: 46 percent); question 13D_A—year of receiving an (upgrade) Associate Degree in Nursing (ADN) (imputation rate: 36 percent); and question 41—years until planned return to paid work in nursing (imputation rate: 21 percent). When working with these three variables, NSSRN data users should be aware of the potential underestimation of the variance and bias in the point estimates.

³ Imputation rates were computed as the ratio of the number of imputed cases of that variable to the total number of applicable cases in the data set.

Further information on imputed variables and data file flags for imputed variables are included in the 2008 NSSRN codebook.

1.7 Sample Weighting

1.7.1 Overview of the Weighting Process

Data weighting was one of the final steps in preparing the data for analysis and reporting. The 2008 NSSRN weighting process consisted of several steps, as follows:

- Determination of the initially assigned weights at the sampled license level;
- Adjustment for survey nonresponse by age group (for strata where age was available);
- Assignment of factors to account for multiple opportunities for selection (determined as part of the matching effort to determine the number of sample frames on which an RN appeared);
- Establishing an initial weight at the RN level; and
- Calibration of the RN level weights (via raking) to achieve consistency with State and national totals.

A detailed discussion of each of these steps in the weighting processes is found in Appendix B. The remainder of this overview discussion focuses on some key issues related to sample weighting for the 2008 NSSRN.

1.7.2 Implications of Weighting for NSSRN Data Users

Sampling error can cause sample estimates to differ from true population parameter values. Sampling error is the consequence of not measuring every unit in the population. It can be reduced with larger sample sizes, with efficiently designed sampling strategies, or with a combination of both.

In the 2008 NSSRN, both strategies were employed. For example, higher sampling rates were used for smaller States to permit more accurate State-level estimation for these States. However, this served to reduce the precision associated with national estimates compared with what would have

been achieved with a simple random sample of the same size but with no variation by State in sampling rates. The factor by which the variance is reduced is known as a "design effect."

In the 2008 NSSRN, sampling rates also varied between strata within a State for RNs who were either:

- Residents of Alaska, New Mexico, or Hawaii but were on another State's listing of RNs with currently active licenses; or
- Members of the groups of States for which the design used multi-State strata.

Replicate weights were computed for use in estimating the variability of 2008 NSSRN estimates. Users of the 2008 NSSRN data as well as past NSSRN data are encouraged to use these weights to assess the precision of sample estimates as well as to compare estimates either within the 2008 NSSRN database or between estimates from various years.

Another important difference in the weighting methods for 2008 was the adjustment of differential nonresponse by age group. Earlier administrations of the NSSRN may have underestimated the representation of younger RNs due to undercoverage on State sampling frames. In the 2000 and 2004 NSSRN administrations, poststratification of the weights of the youngest RNs (those under the age of 26) was done to adjust the weights. However, this group represented only a small portion of licensed RNs, and the overall impact of this procedure did not address any issues that may have existed for other age categories.

While undercoverage of newly licensed RNs may still be an issue, another source of bias associated with younger RNs is that response propensities by age showed that younger RNs responded at lower rates than older RNs in the 2008 NSSRN.

In administrations prior to the 2008 NSSRN, the weight adjustment processes were global in nature. Generally speaking, poststratification was used to achieve consistency between responding RNs associated with a given State of licensure and the corresponding control total, licensed RNs in the State (who were currently active). The control total took into account RNs who were also licensed in what were characterized as "higher priority" States. Thus, if older RNs had higher response rates in a State, as they often did, they would be represented at a disproportionately high rate among the population of currently active RNs licensed in the State. This is a potential source of bias for survey estimates that are correlated with age. In addition to the general age distribution of the RN

population, other variables that could be affected include those related to education and employment.

In summary, the 2008 NSSRN final weighting processes included a poststratification adjustment of eligible and ineligible respondents by age group to corresponding age group totals for each State frame where age was provided by the State licensing Boards (44 of 51 State Boards).

Assessing the Impact of Accounting for Differential Nonresponse by Age

To assess the extent to which survey estimates may have changed after accounting for differential nonresponse by age group, a separate set of weights was developed using a global approach to nonresponse adjustment. That is, a poststratification of all responding eligible and ineligible RNs to a single State control total for each State was undertaken. This approach thus served to simulate the approach to nonresponse adjustment used in past NSSRN administrations. This weight is referred to here as the "surrogate" weight.

A series of statistics were run to compare the estimate based on the final weight to the estimate based on the surrogate weight. Looking at some key data elements from the 2008 NSSRN, the comparison of the final weight to the surrogate weight suggests that including a weight adjustment reflecting differential response propensities by age had an important impact on the 2008 NSSRN survey estimate. Estimates where sizeable effect is in evidence are: initial nursing education, RNs with a bachelor's degree; RNs employed in nursing; white, non-Hispanic RNs; RNs employed in a hospital setting; and most of the age group and graduation year categories.

These findings regarding the effects of nonresponse adjustment by age group should be taken into account when comparing estimates from the 2008 NSSRN to those from previous NSSRN administrations. Differences between the 2008 NSSRN estimates and those from prior years for estimates correlated with age may be due to an improvement in the weighting process rather than a significant change in the RN population over time.

Appendix B provides a more detailed discussion of the weighting process and nonresponse adjustment by age.

This page deliberately blank.

2.1 Introduction

RNs are prepared in several types of education programs. The most frequent initial education program is the Associate Degree in Nursing (ADN), with 45.4 percent of U.S. RNs prepared in ADN programs (Figure 2-1). Some RNs pursue their initial registered nurse education after completing postsecondary education in other fields or working in another occupation. After completing an approved initial education program, a registered nurse candidate takes the National Council Licensure Examination (NCLEX), which is created and administered by the National Council of State Boards of Nursing (NCSBN), in order to receive a State license. RNs subsequently can pursue additional nursing education, including graduate-level degrees and education in Advanced Practice fields such as nurse practitioner or nurse anesthetist. Some RNs seek higher degrees in non-nursing fields (such as education and business) that are related to their nursing career.

2.2 Initial Nursing Education

Initial RN education occurs through a variety of routes. ADN programs generally require 2 years of nursing coursework after completion of prerequisites that include basic science. Bachelor's degree programs typically span 4 years of full-time education. Diploma programs, which are traditionally based in hospitals and last 3 years, were once the most common form of nursing education in the United States. The number of these programs has steadily declined since the 1970s and many now offer an Associate of Science Degree (AS) through local community colleges. A small number of RNs qualify for their licenses based on military training or other licensing credentials. There are a few initial RN education programs that confer a master's degree or a doctorate but most are specifically designed for students who have obtained a bachelor's or higher degree in another field prior to pursuing nursing education.

The most commonly reported route for initial nursing education of RNs in the United States is the ADN program (Figure 2-1).




Figure 2-1. Distribution of registered nurses according to initial nursing education, 1980-2008*

Education and Licensing of Registered Nurses

* The counts for all initial degrees may not add to the total registered nurse estimates for each survey due to incomplete information provided by respondents and the effect of rounding. Only those who provided education preparation information are included in the calculations used for this figure.

**For 2008 initial education, there were 13,325 registered nurses with master's degrees (0.4%) and 954 registered nurses with doctorate degrees (0.03%).


Source: 1980-2008 National Sample Survey of Registered Nurses


Since 1980, there has been a steady shift away from RN education in diploma programs to ADN and bachelor's programs although diploma education still accounts for the initial education of 20.4 percent of RNs in 2008. However, only 3.1 percent of RNs who graduated after 2004 reported a diploma as initial nursing education. Fully 34.2 percent of RNs in 2008 were initially prepared at a bachelor's or higher degree level, almost twice as many as in 1980.

2.3 Age at Graduation

Average age at graduation has been comparatively stable since 1994, with the exception of diplomaeducated RNs, for whom the average age has continued to rise. The average age at graduation of

RNs who graduated from their initial nursing education program after 2004 is 31 years, notably higher than the average age of 24 years for RNs who graduated before 1985 (Figure 2-2). The increase in average age at graduation occurred primarily among RNs who graduated between 1985 and 1994. Registered nurses whose initial nursing education is a bachelor's degree or higher tend to be 5 years younger on average than either ADN or diploma graduates.





Source: 2008 National Sample Survey of Registered Nurses

2.4 Sources of Financing

The most common source of educational financing used by RNs for their initial education was from family resources (42.7 percent), with health care-related employment earnings (30.5 percent), and federally assisted loans (29.3 percent) comprising the next highest sources of nursing education

payment sources (Figure 2-3 and Appendix A, Table 3). Details about the sources of financing for recently graduated RNs are presented in Chapter 9.

Registered nurses who completed ADN programs were most likely to have received some financial support for their education from earnings from their health care-related employment (Appendix A, Table 3). RNs prepared in ADN programs, are older than RNs prepared in Bachelor of Science in Nursing (BSN) programs, and may have continued their employment while pursuing their RN education.




Figure 2-3. Sources of financing for initial education for all U.S. registered nurses*

*Entries for sources of financing add to more than 100 percent because more than one source of financing may have been used by an individual RN.

Source: 2008 National Sample Survey of Registered Nurses

2.5 Education and Experience Prior to Initial Nursing Education

Among the 21.7 percent of the RN population in 2008 who had an academic degree prior to their initial nursing degree, 47.8 percent had an ADN, 52.8 percent a bachelor's, and 7.2 percent had a master's or higher degree¹ (Table 2-1 and Appendix A, Table 4). More than 40 percent of all degrees obtained by RNs prior to initial nursing preparation were in a health-related field.

A growing percentage of RNs are earning academic degrees prior to their initial nursing education. From 2000 to 2008, the percentage with prior postsecondary degrees rose from 13.3 percent to 21.7 percent. This trend is discussed in more detail in Chapter 9.

Table 2-1.Degrees prior to initial nursing education, by survey year, 2000-2008

| | 2000 | 2004 | 2008 |
|--|-----------|-----------|-----------|
| | (percent) | (percent) | (percent) |
| Nurses with degrees prior to initial nursing education | 13.3 | 16.2 | 21.7 |

Source: 2000-2008 National Sample Survey of Registered Nurses

More than two-thirds of RNs (67.2 percent) reported working in a health occupation prior to their initial nursing education. Of these, the majority (68.6 percent) worked as aides or nursing assistants (Table 2-2 and Appendix A, Table 5). A higher percentage of registered nurses whose initial RN education was an ADN had previously worked in a health occupation than RNs with other initial education. The length of time in which RNs worked in these other health occupations and the extent to which this work experience might be important to RNs' careers is unknown.

Among all RNs, 17.9 percent were licensed at one time as a licensed practical or vocational nurse (LPN or LVN) (Table 2-2 and Appendix A, Table 5). Registered nurses whose initial RN education was in an ADN program were more likely to have previously held an LPN or LVN license. In some States, partial completion of an RN education program provides the qualifications to take the exam for LPN/LVN licensure so that RN students may take the exam and work prior to completion of the RN program. Other ADN programs provide flexible fast track education for LPN or LVNs.

¹ The number of degrees adds to more than the total because more than one degree may have been named by an RN.



| | | Initial nursing education | | | |
|---|--------------------|---------------------------|------------------------|---------------------------------------|--|
| | Total (percent) | Diploma (percent) | Associate (percent) | Bachelor's and higher (percent) | |
| All nurses | | | | | |
| Nurses with prior academic degrees | 21.7 | 9.8 | 24.6 | 25.0 | |
| Nurses with prior health occupations | 67.2 | 49.1 | 76.3 | 66.2 | |
| Nurses with prior LPN license Of those with prior health occupations | 17.9 | 7.1 | 30.0 | 8.3 | |
| Nursing aide or assistant | 68.6 | 77.1 | 61.3 | 76.1 | |

Table 2-2.Education and health occupation experience prior to initial nursing education, by
initial nursing education

Source: 2008 National Sample Survey of Registered Nurses

2.6 Nursing Licensure

The majority of RNs who earn their initial nursing education in the United States obtain their first license in the same calendar year they graduate from their initial nursing education program and most RNs are first licensed in the State in which they were educated. Internationally educated nurses have different patterns of graduation and U.S. licensure, as described in Chapter 8.

About 1 percent of RNs obtained their first U.S. nursing license before the calendar year in which they graduated from their initial nursing education program. With the introduction of more nursing master's programs, some State licensing boards permit students who have completed a specific subset of courses and training to take the licensing examination and receive an RN license. Registered nurses are then expected to complete the remaining requirements for their degree, although in some States completion of the degree is not required to retain an RN license. A small number of RNs reported their initial nursing degree or credential as "other" and described such a situation.

More than 90 percent of RNs receive their first license in the State where they were educated. Over time, however, although some RNs may leave their State of initial education, approximately 80 percent remain in the same State at least 5 years after graduation. By 16 to 25 years after graduation however, 30 percent reside in a different State (Appendix A, Table 7). RNs whose initial nursing education is a bachelor's or higher degree are the least likely group of RNs to reside in the State in which they were educated, with 34.8 percent overall living in a different State than that in which they were educated (Appendix A, Table 7). Some RNs may live in a State bordering that in which they were educated and could be commuting to that State.

RNs who work in a State that participates in the Nurse Licensure Compact (the Compact) can work in multiple States even if they hold only one RN license. States in the Compact agree to recognize licenses from other States in the Compact rather than require another RN State license. The effect of the Compact on employment is discussed in Chapter 3.

2.7 Degrees Obtained After Initial Nursing Education

The NSSRN collects data on all academic degrees received after initial RN education; this includes both nursing degrees and non-nursing degrees. Table 2-3 presents the highest nursing degrees of RNs. Among all licensed RNs, 47.2 percent have earned at least a bachelor's degree in nursing.

| | Total (number) | Total (percent) | Working in nursing (number) | Working in nursing (percent) | Not working in nursing (number) | Not working in nursing (percent) |
|--------------------------|-------------------|--------------------|-----------------------------------|------------------------------------|---------------------------------------|--|
| Total | 3,063,162 | 100 | 2,596,599 | 100 | 466,564 | 100 |
| Diploma | 474,043 | 15.5 | 345397 | 13.3 | 128,646 | 27.6 |
| Associate | 1,140,465 | 37.2 | 1,002,682 | 38.6 | 137,783 | 29.5 |
| Bachelor's and higher | 1,444,528 | 47.2 | 1,245,180 | 48.0 | 199,348 | 42.7 |

| Table 2-3. | Highest nursing degree, by employment status |
|------------|--|
| | |

Source: 2008 National Sample Survey of Registered Nurses

Many RNs hold additional degrees that are not specifically nursing degrees but which are related to their nursing employment. If an RN respondent indicated that a non-nursing degree was related to his or her career in nursing, the degree is described as a "nursing-related" degree. Nursing-related degrees include public health, health administration, social work, education, and other fields. For much of this report, nursing and nursing-related degrees are presented together to provide a measure of the level of education from which RNs draw to complete their jobs.

In 2008, half of the RN population had a bachelor's or higher degree in nursing or a nursing-related field (Figure 2-4 and Appendix A, Table 8). In 2008, 32 percent of RNs with a bachelor's or higher degree reported that their initial RN education was a diploma or an ADN, demonstrating that many RNs pursue additional degrees after completion of their initial RN education. Other data reveal that RNs with graduate degrees have increased significantly in the last 20 years (Figure 2-4).




Figure 2-4. Highest nursing or nursing-related education preparation, 1980-2008*

*The totals in each bar may not equal the estimated numbers for registered nurses in each survey year due to incomplete information provided by respondents and the effect of rounding. Only those who provided nursing education preparation information are included in the calculations used for this figure.

Source: 1980-2008 National Sample Survey of Registered Nurses

- Registered nurses with graduate degrees comprised 13.2 percent of all licensed RNs in 2008, twice the proportion in 1988.
- There has been marked growth in the numbers of RNs with a master's degree in nursing or a related field in less than 10 years rising from 257,812 in 2000 to 375,794 in 2008, a 45.8 percent increase (Table 2-4).
- There were an estimated 28,369 RNs with a doctorate degree in nursing or a nursing-related field in 2008, which is an increase of 64.4 percent since 2000.

Although many RNs pursue additional degrees after their initial RN education, many did not obtain further academic degrees post-licensure (Figure 2-5 and Appendix A, Table 8).

- Among the 623,548 RNs whose initial nursing education culminated in a diploma, 31.6 percent obtained additional degrees after licensure. The degree most commonly obtained was a BSN (81,689, or 13.1 percent).
- Among the 1,388,884 RNs whose initial nursing education culminated in an ADN, 20.8 percent obtained additional degrees after licensure. BSNs were received by 12.1 percent.
- Among the 1,044,306 RNs whose initial nursing education culminated in a bachelor's degree or higher, 21.6 percent held advanced degrees in 2008.

Table 2-4.Master's and doctorate degrees as highest nursing or nursing-related education
preparation, 2000-2008

| . . | Estimate | Estimate | Estimate |
|--|----------|----------|----------|
| Degree | 2000 | 2004 | 2008 |
| Master's | 257,812 | 350,801 | 375,794 |
| MSN nursing degree | 202,639 | 256,415 | 290,084 |
| Nursing-related master's degree | 55,173 | 94,386 | 85,709 |
| Percent of master's that are nursing (MSN) | 78.6% | 73.1% | 77.2% |
| Doctorate | 17,256 | 26,100 | 28,369 |
| Doctorate in nursing | 8,435 | 11,548 | 13,140 |
| Nursing-related doctorate degree | 8,821 | 14,552 | 15,229 |
| Percent of doctorates that are nursing | 48.9% | 44.2% | 46.3% |

Source: 2000-2008 National Sample Survey of Registered Nurses




Figure 2-5. Highest education of registered nurses, by initial nursing education

Source: 2008 National Sample Survey of Registered Nurses

There can be considerable time between the initial nursing degree and subsequent degrees (Table 2-5).

- Among diploma-educated nurses who continued their education, there was an average 10.5 year gap between initial RN education and attaining a bachelor's degree. Those whose highest education was a doctorate had a gap of more than 15 years between the diploma and the final doctorate in nursing.
- Nurses whose initial RN education was an ADN averaged 7.5 years between their ADN and attaining a BSN degree.
- Nurses whose initial RN education was a BSN took an average of 8.2 years to reach a master's degree as their highest degree in nursing; they took an average of 12.4 years to obtain a doctorate if that was their highest degree in nursing.

| Initial nursing education | Bachelor's | Master's | Doctorate |
|---------------------------|------------|----------|-----------|
| Diploma | 10.5 | 13.9 | 15.6 |
| Associate | 7.5 | 11.5 | 12.5 |
| Bachelor's | - | 8.2 | 12.4 |

Table 2-5.Average years between completion of initial and highest nursing degrees*

* Average years between diploma and ADN not calculated due to larger than average rates of missing data.

- Too few cases to report estimated percent (fewer than 30 respondents).

Source: 2008 National Sample Survey of Registered Nurses

Post-licensure graduate degree programs have a variety of foci. Among RNs who obtained a nursing or nursing-related master's degree or doctorate, 46.6 percent reported the focus of those degrees to be clinical practice; 19.2 percent had a focus of administration, business, or management; and 13.3 percent reported a focus of education. Education was the most common focus of doctoral studies (please see Appendix A, Table 9 for similar data).

Only 2.2 percent of RNs obtained an undergraduate or graduate level non-nursing degree that they reported was not related to their career in nursing. The most common focus of these degrees was humanities; liberal arts or social sciences; and administration, business, or management.

Education programs are increasingly being offered through distance-based learning. The flexibility offered by distance-based education permits working professionals and those in nonurban centers to pursue degree programs while continuing to work without having to travel to an educational institution. Just over 15 percent of RNs who completed any additional nursing or nursing-related degree program reported that the program was primarily undertaken through distance-based learning, with more than 50 percent of coursework through correspondence or online. Those who completed doctorate degrees had the highest rates of participation in distance-based programs (16.3 percent). The continued prevalence of distance and online education may influence further growth in RNs continuing education beyond their initial preparation program.

2.8 Currently Enrolled Students

Only 6.6 percent of all RNs were enrolled in March 2008 as students in nursing and nursing-related academic education programs. Of these, 44.5 percent were pursuing a bachelor's degree and 46.3 were pursuing a master's (Table 2-6). Among RNs who were students in 2008, 74.2 percent were working in nursing full-time and 18.7 percent were employed in nursing part-time. Only 7.1 percent were not employed in nursing while they were pursuing additional education (Appendix A, Table 10).



| Degree pursued | Number enrolled in 2008 | Percent of those enrolled | Percent with more than 75 percent distance- based |
|---------------------------|-------------------------|------------------------------|---|
| Associate | 3,151 | 1.5 | - |
| Bachelor's | 90,553 | 44.5 | 49.1 |
| Master's | 94,205 | 46.3 | 41.5 |
| Doctorate | 13,479 | 6.6 | 35.0 |
| Post-master's certificate | 2,065 | 1.0 | _ |
| Total | 203,453 | 100.0 | 44.3 |

Table 2-6. Enrollment in nursing and nursing-related education programs

- Too few cases to report estimated percent (fewer than 30 respondents).

Source: 2008 National Sample Survey of Registered Nurses

Distance-based programs are more common for programs offering a post-licensure bachelor's degree (49.1 percent) than for other post-licensure degrees (Table 2-6). Full-time RNs are more likely than part-time RNs or those not employed in nursing to take more than 75 percent of their coursework in a distance based format. Overall, RNs working full-time in nursing are more likely than any other group of RNs (48.2 percent) to take more than 75 percent of their coursework as distance learning (Appendix A, Table 10).

2.9 Emergency Preparedness

When emergencies and disasters occur, RNs are often required to assume responsibilities for managing and caring for overwhelmingly large numbers of patients with unfamiliar injuries, sometimes in environmental conditions that are extreme. More than half of working RNs (53.8 percent) indicated that they neither received nor provided emergency training in the previous year. Of those who had received or provided training, the most commonly reported training was for infectious disease incidents (71.1 percent) or natural disasters (70.0 percent). Only 31.3 percent of RNs reported receiving or providing training for nuclear or radiological incidents. Of the 46.2 percent of RNs who received training in some form of emergency preparedness training, the training they received averaged 10.5 hours in the year.

Few registered nurses reported feeling well prepared for an emergency. Of those RNs who have received or provided emergency preparedness training, only 10.5 percent feel very prepared, while 44.1 percent feel somewhat or not at all prepared. Nonetheless, 94.9 percent of RNs claim to have some understanding or a full understanding of the disaster plan at their place of work.

2.10 Certifications

After receiving initial RN education and licensure, registered nurses demonstrate knowledge and skill for a population or in a patient care specialty by obtaining national standardized certifications from a variety of organizations. Certifications can be obtained in dozens of fields, ranging from clinical specialties such as oncology, pediatrics, and critical care, to functional specialties such as administration, case management, or legal consulting. A myriad of organizations offer certifications, which are usually conferred only after completing a specified education or training program, a period of clinical practice and/or passing a national test. Advanced practice RNs, such as nurse practitioners, can and often must obtain certification in their advanced area. This is discussed further in Chapter 5.

More than 35 percent of RNs report some type of certification (Table 2-7 and Appendix A, Table 11). The most frequently held certification is life support/resuscitation. More than 31 percent of RNs have life support or resuscitation certification (BLS, ALS, BCLS, CPR, NRP, and others). The most common areas for clinical or managerial certification are critical care, maternal/neonate, oncology, case management, general surgery, and medical/surgical areas. In the 2008 NSSRN, some RNs report certifications that are not specific to nursing but may be relevant to their nursing careers, such as massage therapy.

Table 2-7. Registered nurse certifications*

| | Total | Total |
|--|-----------|-----------|
| Type of certification | (number) | (percent) |
| Total registered nurses with certifications | 1,094,838 | 35.7 |
| | | |
| Life support/resuscitation (BLS, ALS, BCLS, CPR, NRP, and others) | 956,472 | 31.2 |
| Trauma nursing/emergency medicine (TNCC, ATCN, ATN, EMT, ENPC, and others) | 125,008 | 4.1 |
| Critical care | 58,320 | 1.9 |
| Maternal/neonate | 43,123 | 1.4 |
| Oncology | 36,175 | 1.2 |
| Case management | 31,106 | 1.0 |
| General surgery | 28,599 | 0.9 |
| Medical/surgical | 26,594 | 0.9 |

*Table excludes advanced practice RN certifications, which are reported in Chapter 5.

**Entries for types of certification do not add to the total because an individual registered nurse may have reported more than one type of certification.

Source: 2008 National Sample Survey of Registered Nurses

This page deliberately blank.



3.1 Introduction

The RN workforce consists of nurses who have valid, active RN licenses and are employed in a nursing position. The vast majority of RNs are employed in nursing. Their attachment to the labor market is high, and in general RNs are satisfied with their work. Although RNs work in a variety of settings and roles, more than half are employed in hospitals and 66.3 percent are in staff nurse roles.

3.2 Nursing Workforce Size

In 2008, an estimated 3,063,162 RNs were in the United States (Figure 3-1 and Appendix A, Table 1), and 84.8 percent of them were employed in nursing positions (2,596,599). This was the highest rate of nursing employment since the NSSRN commenced in 1977 (Figure 3-2). Full-time employment has also increased, rising from 58.4 percent of RNs in 2004 to 63.2 percent in 2008. This is the first increase in the percentage of RNs working full-time since 1996.

The number of employed RNs per 100,000 population is a ratio used to measure whether the RN workforce is growing more or less rapidly than the U.S. population as a whole. In 2008, there were an estimated 854 employed RNs per 100,000 population, an increase from 825 per 100,000 population in 2004 (Appendix A, Table 57). This measure varies widely among the States. The lowest numbers of employed RNs per 100,000 population were in Utah (598), Nevada (618), and California (638), while the largest numbers were in the District of Columbia (1,868), South Dakota (1,333), and North Dakota (1,273).




Figure 3-1. The nursing population and workforce in the United States, 1980-2008

Source: 1980-2008 National Sample Survey of Registered Nurses




Figure 3-2. Employment status of registered nurses*

*The totals of full-time and part-time employment may not add to the estimate of all RNs employed in nursing due to incomplete information provided by respondents on employment status and the effect of rounding.

Source: 1980-2008 National Sample Survey of Registered Nurses

3.3 Employment and Age

The age distribution of the RN population has been rising for more than two decades (Figure 3-3). In recent years this aging trend has slowed. RNs are less likely to work in nursing positions as they age (Figure 3-4). Among RNs under 50 years old, 90 percent or more are employed in nursing positions. The percentages of RNs working in nursing drops in each age group older than 50, decreasing from 87.5 percent of RNs aged 50-54 to 85.2 percent of RNs aged 55 to 59 years and to less than half of RNs over age 65. More than 70 percent of RNs under 30 years old work full-time. The percentage of RNs aged 30 to 59 years and working full-time ranges between 65 percent and 70 percent. The percent of RNs who work full-time drops rapidly after age 60 though more than 30 percent of RNs over the age of 70 continue to work in nursing in either full-time or part-time positions.




Figure 3-3. Age distribution of the registered nurse population, 1980-2008

Source: 1980-2008 National Sample Survey of Registered Nurses




Figure 3-4. Employment of registered nurses, by age group*


* The total percent of registered nurses across age groups may not equal the estimated total of all RNs due to the effect of rounding. Source: 2008 National Sample Survey of Registered Nurses


3.4 Employment and Education

Employment rates of RNs vary both by initial and highest education. More than 87 percent of nurses whose initial RN education was an ADN, bachelor's, or master's degree report that they are employed in nursing positions, as compared with less than 75 percent of those whose initial

3

education was a nursing diploma (Figure 3-5 and Appendix A, Table 8). This is consistent with the older average age of diploma graduates, discussed in Chapter 7. Further, RNs whose initial education was in an ADN program have a higher rate of full-time employment than do those who entered the profession with a bachelor's or master's degree.





*Percents may not add to 100 due to the effect of rounding. Only those who provided education preparation information are included in the calculations used for this figure.

Source: 2008 National Sample Survey of Registered Nurses

Figure 3-6 illustrates that the relationship between highest education and employment is similar to the relationship for initial education. About 26 percent of RNs for whom diploma-education was their highest education no longer work in nursing, as compared with about 12 percent, 14 percent, and 17 percent for RNs whose highest education is an ADN, bachelor's, and master's/doctorate, respectively. RNs with a bachelor's or master's/doctorate degree are somewhat less likely to work

full-time than RNs with an ADN. Overall, more than 75 percent of RNs, regardless of current education level, work in nursing.




Figure 3-6. Employment rates, by highest nursing or nursing-related education*

*Only those who provided education preparation information are included in the calculations used for this figure.

Source: 2008 National Sample Survey of Registered Nurses

3.5 Employment Settings

RNs work in a variety of settings and have many different roles and responsibilities in each of these settings. For this report, employment settings are defined using industry categories in labor market analyses by the U.S. Bureau of Labor Statistics. For the first time, in the 2008 NSSRN, nursing home

and skilled nursing units that are based within hospitals are grouped as part of the hospital setting. In the 2004 NSSRN, these settings were grouped with nursing homes and extended care facilities. Therefore, in the 2008 NSSRN the number of RNs working in hospital settings versus nursing home and skilled care facilities are not completely comparable to previous surveys.

Hospitals remain the most common employment setting for RNs in the United States, with 62.2 percent of employed RNs reporting that they worked in hospitals in 2008 (Figure 3-7).¹ Ambulatory care is the next most common setting, with 10.5 percent of employed RNs. Public and community health settings, which include schools and public health departments, account for the worksite of almost 8 percent of RNs, while 6.4 percent work in home health settings. Another 5.3 percent work in nursing homes and extended care facilities. Only 3.8 percent work in academic education. Therefore, almost 30 percent of RNs worked in nonacute patient care settings; more than 90 percent of RNs work in patient care settings of some sort.

Among RNs who work in hospitals, the majority (82.6 percent) worked in community hospitals. Another 7.8 percent were employed in specialty hospitals (Figure 3-8).

- More than 76 percent of RNs employed in non-Federal specialty hospitals worked in inpatient units (Table 3-1).
- Among RNs who worked in community hospitals, 11.7 percent were employed in outpatient clinics or medical practices.
- Only 1.5 percent of RNs who worked in hospital settings were employed in a long-term hospital, and 2 percent worked in a psychiatric hospital (Appendix A, Table 12).²
- Slightly more than 3 percent of RNs who worked in hospital settings were employed in Federal hospitals, such as the U.S. Department of Veterans Affairs (VA), military facilities, or Indian Health Services hospitals.

¹ For the 2008 NSSRN, as with previous administrations, the list of employment settings from which RNs could select was revised. Despite these revisions, the major employment categories are sufficiently distinct to support comparisons over time.

² RNs who worked in long-term hospitals are not grouped with nurses employed in other long-term or skilled nursing facilities in this report.




Figure 3-7. Employment settings of registered nurses*

*Percents may not add to 100 due to the effect of rounding. Only RNs who provided setting information are included in the calculations used for this figure.

**Public/community health includes school and occupational health.

***Ambulatory care includes medical/physician practices, health centers and clinics, and other types of nonhospital clinical settings.

 $\star\star\star\star$ Other includes insurance, benefits, and utilization review.




Figure 3-8. Types of hospitals in which registered nurses employed in hospital settings work*

*Percents may not add to 100 due to the effect of rounding. Only registered nurses who provided hospital subset information are included in the calculations used for this figure.

Source: 2008 National Sample Survey of Registered Nurses

| | Community hospital (percent) | Specialty hospital (percent) | Long-term hospital (percent) | Psychiatric hospital (percent) | Federal hospital (percent) | Other hospital (percent) | Total (percent) |
|-------------------|------------------------------------|------------------------------------|------------------------------------|--------------------------------------|----------------------------------|--------------------------------|--------------------|
| Inpatient | 78.1 | 76.5 | 80.6 | 82.1 | 52.9 | 61.7 | 76.8 |
| Nursing home unit | 0.0 | 0.0 | 9.2 | 1.6 | 3.5 | 1.1 | 0.7 |
| Outpatient | 11.7 | 15.6 | 0.0 | 7.1 | 26.8 | 14.3 | 12.3 |
| Other | 9.8 | 8.0 | 10.2 | 9.1 | 16.8 | 22.9 | 10.2 |
| Total | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 |

Table 3-1. Employment in hospital units, by type of hospital*

*Percents may not add to 100 due to the effect of rounding. Only registered nurses who provided hospital subset information are included in the calculations used for this table.

3.5.1 Employment Setting Trends

The employment settings of RNs have changed over the decades (Figure 3-9). In 2004, 57.4 percent of RNs worked in hospitals. The increase in the percentage of RNs working in hospitals between 2004 and 2008 is the first increase since 1984. The 17.7 percent growth in the number of RNs employed in hospitals is consistent with American Hospital Association (AHA) data. The AHA reported that between 2004 and 2008, RN full-time equivalent (FTE) employment increased 16.6 percent. The AHA also reported that RN FTE percent of total hospital employment rose from 25.4 to 27.³ Hospitals are employing more RNs than in 2004.

There also was a 68 percent increase from 2004 to 2008 in the percentage of RNs working in home health services from 3.8 percent in 2004 to 6.4 percent in 2008. Between 2004 and 2008, little else changed in the percentage of RNs employed in other settings.

3.5.2 Employment Settings and Age

Registered nurses' employment settings change with nurses' ages (Table 3-2 and Appendix A, Table 13). Nearly 85 percent of RNs under 30 years old work in hospitals, but this percentage declines steadily with age; less than 50 percent of RNs age 55 and older work in hospitals. Hospitals have historically been viewed as offering a good initial career development opportunity for a new RN because new RNs can gain hands-on experience with patients with varied and complex needs while having the support of other experienced clinicians inherent in the hospital setting. Nonhospital settings such as home health, public and community health, nursing homes and extended care, and academic education are employment settings for increasing percentages of the older RN age groups. It is not surprising that RNs working in academic education are older because RNs need to complete graduate education before moving into faculty positions.

³ Data retrieved from American Hospital Association, Trendwatch Chartbook 2009, available at <u>http://www.aha.org/aha/research-and-trends/chartbook/index.html</u>. This publication lists non-Federal resources in order to provide additional information to consumers. The views and contents in these resources have not been formally approved by the U.S. Department of Health and Human Services (HHS). Listing these resources is not an endorsement by HHS or its components.




Figure 3-9. Employment settings of registered nurses, 1980-2008*

*The total percent by setting may not equal the estimated total of all registered nurses due to incomplete information provided by respondents and the effect of rounding.

**Public/community health includes school health, occupational health, and home health.

***Other includes insurance, benefits, and utilization review.

| Table 3-2. Em | ployment | setting | s, by age | e group* | | | | | |
|---------------------------|----------|---------------------|-----------|----------|-------|-------|-------|-------|--------|
| | | Age group (percent) | | | | | | | |
| | Under | 30- | | | | | | | 65 and |
| Employment setting | 30 | 34 | 35-39 | 40-44 | 45-49 | 50-54 | 55-59 | 60-64 | over |
| Hospital | 84.5 | 73.5 | 68.3 | 64.7 | 61.3 | 55.9 | 53.0 | 46.5 | 35.5 |
| Nursing home/ | | | | | | | | | |
| extended care | 2.1 | 3.4 | 4.7 | 4.5 | 5.6 | 5.9 | 6.5 | 7.6 | 10.7 |
| Academic education | 1.7 | 1.8 | 2.4 | 2.7 | 3.8 | 4.5 | 5.1 | 6.6 | 10.5 |
| Home health | 2.6 | 4.3 | 6.6 | 6.7 | 6.7 | 7.2 | 7.7 | 7.8 | 10.5 |
| Public/community | | | | | | | | | |
| health** | 2.7 | 4.3 | 4.1 | 7.4 | 8.3 | 9.9 | 9.9 | 13.7 | 14.8 |
| Ambulatory care | 5.8 | 10.6 | 9.7 | 10.5 | 10.7 | 11.5 | 12.6 | 12.1 | 11.3 |
| Other*** | 0.8 | 2.1 | 4.1 | 3.6 | 3.5 | 5.1 | 5.2 | 5.8 | 6.8 |

*Percents may not add to 100 due to the effect of rounding. Only registered nurses who provided setting information are included in the calculations used for this table.

**Public/community health includes school and occupational health.

***Other includes insurance, benefits, and utilization review.

Source: 2008 National Sample Survey of Registered Nurses

3.5.3 Employment Settings and Education

The highest level of nursing or nursing-related education varies with employment setting (Figure 3-10 and Appendix A, Table 14). A larger percentage of RNs whose highest education is a bachelor's degree are employed in hospitals as compared to RNs with an ADN or diploma. Fewer than half of RNs with master's or doctorate degrees work in hospitals; more than 18.6 percent are in ambulatory care settings, and nearly 12 percent are in academic education. The relatively high percentage in ambulatory care reflects the large number of advanced practice registered nurses with master's degrees who work in ambulatory settings as discussed in Chapter 5. Among diploma-educated RNs, 14 percent work in ambulatory care settings, and more than 7 percent work in each of long-term care and home health. These results are consistent with the fact that diploma-educated RNs are older than other RNs, on average, and that older RNs are more likely to work in nonhospital settings.




Figure 3-10. Employment settings, by highest nursing or nursing-related education*

*Percents may not add to100 because of rounding. Only those who provided education preparation information are included in the calculations for this figure.

Source: 2008 National Sample Survey of Registered Nurses

3.6 Job Titles

The specific roles of RNs are defined by their job titles within each employment setting. RNs have a variety of roles and job titles. Different settings and employers often use different terminology to describe the same type of work performed by RNs. For example, some hospitals use the title "direct care nurse," while others use the title "staff nurse" even when the work of RNs with these titles is essentially the same. Job titles can indicate the clinical area or expertise of an RN, but even within a clinical area a variety of job titles can be found. Moreover, some RNs work in several roles within

the same facility. For the purposes of the 2008 NSSRN, RNs were asked to report a single job title corresponding to their position.

The most common job title of RNs in the United States is "staff nurse" or an equivalent title (Table 3-3). The percentage of RNs reporting this title increased from 64.1 percent in 2004 to 66.3 percent in 2008. Job titles of RNs in 2008 also included management and administration titles (12.5 percent) and patient coordinator (5.4 percent). The number responding with advanced practice titles (nurse practitioner, clinical nurse specialist, nurse anesthetist, and nurse midwife) were relatively unchanged since 2004.

Table 3-3. Job titles of registered nurses*

| | 2004 | 2008 |
|---------------------------|-----------|-----------|
| Job title | (percent) | (percent) |
| Staff nurse | 64.1 | 66.3 |
| First-line management | 6.2 | 4.1 |
| Middle management | 5.9 | 6.0 |
| Senior management | 2.5 | 2.4 |
| Patient coordinator | 5.5 | 5.4 |
| Nurse practitioner | 3.5 | 3.8 |
| Clinical nurse specialist | 1.2 | 0.9 |
| Nurse anesthetist | 1.1 | 1.1 |
| Nurse midwife | 0.3 | 0.3 |
| Instruction | 2.6 | 3.7 |
| Patient educator | 0.3 | 0.7 |
| Consultant | 1.5 | 0.9 |
| Researcher | 0.8 | 0.7 |
| Informatics | 0.4 | 0.3 |
| Survey/auditor/regulator | 0.5 | 0.4 |
| Other** | 3.5 | 3.0 |
| Total | 100.0 | 100.0 |

*Only those who provided job title information are included in the calculations used for this table. The response categories for job title were slightly different across the 2004 and 2008 survey administrations which may account for some of the change in distribution.

**Other includes registered nurses who reported they had no job title.

Source: 2004 and 2008 National Sample Survey of Registered Nurses

3.6.1 Job Titles and Employment Settings

Within hospitals, RNs are most likely to be a staff nurse or have an equivalent title (Appendix A, Table 16). Within nursing homes, 40.5 percent of RNs have management or administration job titles. These titles also are common among RNs in home health settings (20.9 percent). Patient

coordinator job titles are frequently reported in insurance claims/utilization review (40.1 percent) and home health (19.6 percent) employment settings.

3.6.2 Job Titles and Experience

Job titles of RNs change with experience and age, much as do the settings in which they work. A higher percentage of recently graduated RNs have staff nurse job titles, and the percentage of RNs with this title declines with increasing years of experience. The percentage of RNs with advanced practice titles is relatively similar for RNs with 6 or more years of experience (Table 3-4). After 10 years of experience, RNs tend to move from staff nursing to management and other positions, though even after 40 years of experience, 50 percent of RNs reported being staff nurses.

| | | Years since graduation from initial nursing education | | | | | | | |
|--|------------------------|---|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|-----------------------------|
| Job title | 5 or fewer years | 6 to 10 years | 11 to 15 years | 16 to 20 years | 21 to 25 years | 26 to 30 years | 31 to 35 years | 36 to 40 years | More than 40 years |
| Staff nurse | 88.4 | 72.2 | 65.5 | 63.2 | 57.3 | 55.4 | 53.8 | 55.2 | 50.1 |
| First-line management Middle/senior | 2.1 | 4.0 | 4.6 | 5.0 | 4.8 | 5.6 | 4.7 | 3.5 | 2.8 |
| management | 2.2 | 6.1 | 8.9 | 9.4 | 11.1 | 11.9 | 12.0 | 11.8 | 11.8 |
| Advance practice | | | | | | | | | |
| registered nurse | 2.0 | 6.2 | 5.9 | 5.9 | 8.4 | 8.4 | 8.9 | 6.3 | 6.3 |
| Instruction | 0.8 | 3.0 | 3.6 | 3.9 | 4.7 | 4.5 | 5.4 | 5.5 | 8.0 |
| Other job title | 4.6 | 8.5 | 11.5 | 12.5 | 13.6 | 14.1 | 15.2 | 17.7 | 21.0 |

Table 3-4. Job titles, by experience in nursing*

*Percents may not add to 100 because of rounding. Only those who provided job title information are included in the calculations used for this table.

Source: 2008 National Sample Survey of Registered Nurses

3.6.3 Job Titles and Education

Nursing and nursing-related education can support registered nurses' career goals and education is associated with RN job titles. More than 69 percent of RNs whose highest education is a nursing diploma, ADN, or bachelor's degree have staff nurse job titles, whereas less than 20 percent of those with graduate degrees hold a staff nurse job title (Table 3-5 and Appendix A, Table 18). Approximately 70 percent of RNs with graduate degrees have advance practice, management, and instruction titles.

| | Highest nursing or nursing-related education prepara (percent) | | | | |
|--|---|-----------|------------|-----------|--|
| | | Associate | Bachelor's | Master's/ | |
| Job title | Diploma | degree | degree | doctorate | |
| Staff nurse | 69.7 | 76.4 | 71.4 | 19.4 | |
| First-line management | 4.6 | 4.1 | 4.2 | 3.4 | |
| Middle/senior management | 7.7 | 7.1 | 7.5 | 15.5 | |
| Certified registered nurse anesthetist | 1.0 | 0.1 | 0.5 | 6.1 | |
| Clinical nurse specialist | 0.5 | 0.5 | 0.5 | 3.2 | |
| Certified nurse-midwife | 0.1 | 0.0 | 0.1 | 1.5 | |
| Nurse practitioner | 0.9 | 0.3 | 0.4 | 26.3 | |
| Patient educator | 0.7 | 0.6 | 0.8 | 0.8 | |
| Instruction | 1.7 | 1.5 | 3.1 | 13.6 | |
| Patient coordinator | 7.2 | 5.2 | 5.9 | 3.1 | |
| Informatics nurse | 0.3 | 0.3 | 0.4 | 0.5 | |
| Consultant | 1.1 | 0.5 | 0.9 | 1.8 | |
| Researcher | 0.6 | 0.5 | 0.9 | 0.8 | |
| Surveyor/auditor/regulator | 0.5 | 0.4 | 0.4 | 0.4 | |
| Other | 3.5 | 2.5 | 3.1 | 3.4 | |

Table 3-5. Job titles, by highest nursing or nursing-related education*

*Percents may not add to 100 because of rounding. Excludes registered nurses for whom education preparation was not known and registered nurses for whom job title was not known.

Source: 2008 National Sample Survey of Registered Nurses

3.7 Secondary Nursing Jobs

Overall, 327,259 RNs have more than one nursing position (Appendix A, Table 19), slightly less than the 352,027 who held more than one nursing position in 2004. More than 12 percent of RNs who have a full-time principal nursing position have reported at least one other nursing job, and 14 percent of those with part-time principal positions have reported more than one nursing job (Figure 3-11).

RNs may hold their secondary positions as employees of health care facilities or temporary placement agencies, or as consultants. Some RNs work "per diem" for hospitals, nursing homes, home care agencies, and outpatient facilities as facility employees or through agencies. Per diem employment arrangements allow RNs to work shifts for specific employers on an as needed basis, often without employee benefits but at a slightly higher pay rate than scheduled employees, based on employers' requests and the nurses' willingness to fill these requests. Per diem employment is often possible for all levels of RN clinicians including both staff nurses and advanced practice registered nurses.




Figure 3-11. Registered nurses holding multiple positions, by primary employment status

Source: 2008 National Sample Survey of Registered Nurses

More than half of RNs who hold secondary nursing positions report that, for at least one of their secondary positions, they are regular employees of some firm (Table 3-6). About 36 percent of RNs with secondary nursing jobs report that at least one of these secondary positions is per diem or they are self-employed. Another 8.4 percent of full-time RNs and 4.9 percent of part-time RNs say their secondary nursing jobs involved employment through a temporary placement agency that does not require them to travel long distances for assignments. Agency employment may afford the same flexibility as do per diem arrangements, allowing RNs to work a schedule of their own choosing.

| | Principal nursing position (percent) | | | | |
|----------------------------|--------------------------------------|-----------|---------------------------------------|--|--|
| Secondary nursing position | Full-time | Part-time | All nurses with multiple positions | | |
| Regular employee | 57.1 | 62.6 | 58.6 | | |
| Nontravel agency | 8.4 | 4.9 | 7.4 | | |
| Travel agency | 1.8 | 1.3 | 1.7 | | |
| Self-employed or per-diem | 36.7 | 36.3 | 36.6 | | |

Table 3-6. Employment arrangements in secondary nursing positions*

*Percents will not add to 100 because RNs could select more than one employment arrangement.

Source: 2008 National Sample Survey of Registered Nurses

Hospitals have the largest percent of secondary nursing positions, with 43.3 percent of RNs who have secondary nursing employment working in this setting (Table 3-7). Ambulatory care secondary positions are reported by 13.5 percent of RNs who have secondary employment, while home health is reported by 11.4 percent. Academic education is another setting for secondary employment (10.6 percent); employment in academic education is discussed in more detail in Chapter 4.

Table 3-7. Employment settings of secondary nursing positions*

| Employment setting | Percent |
|------------------------------------|---------|
| Hospital | 43.3 |
| Nursing home/extended care | 9.4 |
| Academic education program | 10.6 |
| Home health | 11.4 |
| Public/community health | 7.2 |
| School health services | 4.4 |
| Occupational health | 1.7 |
| Ambulatory care | 13.5 |
| Insurance claims/benefits | 1.5 |
| Telehealth/telenursing/call center | 0.8 |
| Military/reserves/national guard | 1.0 |
| Medical legal | 0.7 |
| Consultant/contractor | 0.9 |
| Other | 1.4 |
| | |

*Percents will not add to 100 because RNs could select more than one secondary employment setting.

Source: 2008 National Sample Survey of Registered Nurses

3.8 Supply of Registered Nurses

The amount of hours that RNs work per week and per year are measures of the supply of nursing services available for patient care, management, and other services. In the 2008 NSSRN, RNs were asked to report the number of months they work per year in their principal nursing position and the

number of weeks they work per year in their secondary nursing positions. RNs also were asked about the number of hours per week they work in each of their nursing positions.

The vast majority of RNs (92.7 percent) report that they work at least 11 months per year in their principal nursing position (Figure 3-12). This level of employment is equivalent to full-year employment for most positions. Another 5 percent of RNs work 9 to 10 months per year, which is full-year for the academic or school calendar. The remaining 2.3 percent work 8 or fewer months per year.

More than half of RNs work at least 2,000 hours per year in their nursing positions (54.2 percent).⁴ Only 7.6 percent of employed RNs work fewer than 1,000 hours per year in nursing (Figure 3-13). More than half of RNs work at least 40 hours per week in their principal nursing position (Figure 3-14), and another 24.2 percent work 32 to 39 hours per week. Some hospitals use 12-hour shifts for their schedules and consider three 12-hour shifts per week (36 hours) a full-time position. In this arrangement, RNs who choose to work one extra shift per week will normally work 48 hours. Only 4.5 percent of RNs work fewer than 16 hours per week in their principal nursing position; this is consistent with the small percentage of RNs that work fewer than 1,000 hours per year. RNs work an average of 1,944 hours per year in their principal nursing position (Appendix A, Table 20). Almost 17 percent of RNs work more than 2,500 hours per year.

RNs who hold one additional nursing position work an average of 29.6 weeks per year and 15.2 hours per week in their second job (Table 3-8). These data indicate that many RNs work only part of the year and part of the week in their second positions. RNs who work in third jobs average only 23.1 weeks per year and 11.7 hours per week, while those who work four and more jobs work even fewer hours in these additional jobs.

RNs employed in multiple nursing positions work more hours per year than do those with only one nursing position (Table 3-9). However, RNs employed in four or more positions work fewer hours per year on average than do RNs with three positions, suggesting that RNs with more positions may work several part-time positions.

⁴ A full-time position of 40 hours per week for 52 weeks is equal to 2080 hours per year. A registered nurse who works 48-50 weeks per year, allowing for vacation time, at 40 hours per week would work from 1920–2000 hours per year.




Figure 3-12. Months worked per year in principal nursing position

Source: 2008 National Sample Survey of Registered Nurses




Figure 3-13. Hours worked per year in all nursing positions

Source: 2008 National Sample Survey of Registered Nurses




Figure 3-14. Hours worked per week in principal nursing position

Source: 2008 National Sample Survey of Registered Nurses

Table 3-8. Average hours and weeks worked in secondary nursing positions

| | Weeks per year | Hours per week |
|-----------------------------|----------------|----------------|
| Second nursing job | 29.6 | 15.2 |
| Third nursing Job | 23.1 | 11.7 |
| Fourth or more nursing jobs | 11.9 | 7.8 |

Source: 2008 National Sample Survey of Registered Nurses

Table 3-9.Average hours per year worked in all nursing jobs, by number of nursing jobs

| | Average hours worked per year |
|---------------------------|-------------------------------|
| One nursing job | 1,945.4 |
| Two nursing jobs | 2,349.9 |
| Three nursing jobs | 2,859.7 |
| Four or more nursing jobs | 2,519.4 |

3.8.1 Hours and Weeks Worked and Age Group

RNs tend to work fewer hours as they grow older, in addition to being less likely to work in nursing (Table 3-10). RNs through age 60 worked, on average, more than 2,000 hours per year. After age 65, the average number of hours per year drops rapidly. Still, RNs who are 65 years or older do work an average of 1,393 hours of nursing work per year.

| Age group | Average hours worked per year |
|-------------|-------------------------------|
| Under 30 | 2,020.2 |
| 30-34 | 2,005.7 |
| 35-39 | 2,007.2 |
| 40-44 | 2,003.5 |
| 45-49 | 2,095.0 |
| 50-54 | 2,092.0 |
| 55-59 | 2,047.0 |
| 60-64 | 1,826.9 |
| 65 and over | 1,392.7 |

 Table 3-10.
 Average hours worked per year in all nursing positions, by age group

Source: 2008 National Sample Survey of Registered Nurses

3.8.2 Hours and Weeks Worked and Highest Education

The number of hours per year that RNs work in nursing positions is least for RNs whose highest nursing or nursing-related education is a diploma (Table 3-11). This is not surprising, since the average age of diploma-educated RNs is higher than for RNs with other types of education and older RNs work less hours. RNs with master's and doctorate degrees report the most hours worked per year. These are RNs who are also likely to be managers, faculty, or advanced practice registered nurses.

Table 3-11.Average hours worked per year in all nursing positions, by highest nursing and
nursing-related education

| Degree | Average hours worked per year |
|-----------------------|-------------------------------|
| Diploma | 1,864.9 |
| Associate degree | 2,047.1 |
| Bachelor's degree | 1,966.6 |
| Master's or doctorate | 2,122.7 |

3.8.3 On-Call Work

Some RNs work in an "on-call" arrangement for at least some of their total hours. In this type of arrangement, an RN is not physically present at an employer's facility (or at a patient's home in the case of hospice or home care services) but is available to be on-call to work during a prescheduled time. For example, a surgical nurse or nurse anesthetist might be on-call to assist in a surgery between 8 p.m. and 6 a.m. or a nurse midwife may be on-call for deliveries at night or on weekends. Often advanced practice primary care nurses take telephone calls for patient care issues after normal hours. RNs can be paid additionally for these on-call hours usually at a lower rate than is paid for regular work. However, if the RN is called in to work, the RN is paid the normal hourly rate often with an added premium for working after normal scheduled hours. Sometimes, if an RN works in a salaried position, the RN may be on-call but may not receive additional hourly standby pay.

In their principal nursing positions, 19.1 percent of RNs report that they worked on-call or were on standby in a typical work week. Most RNs who reported any on-call activity indicated that they were on standby or worked 20 or fewer hours in a typical week (Table 3-12). RNs who had any on-call activity indicated that they were called to work an average of 12.5 hours per week. They were on standby without being called to work an average of 23.1 hours per week. An average of 14 percent of on-call hours were paid at a standby rate.

| Hours | Percent |
|------------------|---------|
| 0 hours | 80.9 |
| 1 to 20 hours | 12.2 |
| 21 to 40 hours | 3.8 |
| 41 to 60 hours | 1.4 |
| 61 or more hours | 1.6 |
| Total | 100.0 |

Table 3-12.Average hours on standby or worked from on-call per week in the principal nursing position

Source: 2008 National Sample Survey of Registered Nurses

3.8.4 Overtime Work

Typically, RNs paid on an hourly wage basis may be paid for overtime work beyond 40 hours a week and may be compensated at a premium rate. Salaried RNs normally do not receive additional payment for overtime work but may sometimes take paid extra time off later to compensate for working extra hours. While overtime work is ideally a match between an employer's need for additional effort and the nurse's willingness to provide additional hours, sometimes RNs are required to work overtime ("mandatory overtime"). Mandatory overtime may sometimes be voluntarily accepted as part of the employment arrangement, rather than being totally involuntary. Moreover, in States or facilities where RNs are unionized, overtime may be subject to union contract.

The majority of RNs do not work overtime. Among RNs employed full-time in nursing positions, 27.5 percent report that they worked overtime and received pay for such work (Appendix A, Table 21). Mandatory overtime was reported by 7 percent of RNs. RNs work an average of 7.5 overtime hours per week and averaged 6.6 hours per week of mandatory overtime.

3.8.5 Hours Worked and Employment Setting

The number of hours RNs work per week and per year varies across employment settings (Table 3-13 and Appendix A, Tables 20, 21, and 22).

- RNs whose principal nursing position is in a school health setting reported the fewest hours per year, with an average of 1,443 hours per year; this low work rate is likely related to their work schedules being based on an academic calendar and the holidays, vacation periods, and other time off related to the school calendar. They also reported the fewest hours worked per week, averaging 39 hours for full-time RNs.
- The greatest number of average hours RNs worked per year in their principal nursing position (2,107 hours) was in nursing homes and extended care facilities.
- Among RNs working full-time, the greatest number of hours per week was reported by RNs whose principal position was in a nursing home (45.8), followed by those in home health (45.3).
- For RNs working part-time, the greatest number of hours per week was reported by RNs whose principal position was in a hospital (25.2), followed by home health (24.4).
- The average number of hours worked per week in principal nursing positions was more than 40 for full-time RNs in all settings, except those in school health.

| Employment setting | Mean hours/week full-time nurses | Mean hours/week part-time nurses |
|---------------------------------------|-------------------------------------|-------------------------------------|
| All settings | 42.9 | 24.1 |
| Hospital | 42.6 | 25.2 |
| Nursing home/extended care | 45.8 | 22.4 |
| Academic education | 44.0 | 22.8 |
| Home health | 45.3 | 24.4 |
| Public/community health | 42.7 | 21.6 |
| School health | 39.0 | 18.9 |
| Occupational health | 42.4 | 22.5 |
| Ambulatory care | 42.5 | 22.3 |
| Insurance/benefits/utilization review | 43.1 | 21.0 |
| Other | 45.1 | 23.0 |

Table 3-13.Average hours per week in principal nursing position, by full-time/part-time status
and employment setting

Source: 2008 National Sample Survey of Registered Nurses

Overtime work was more common among RNs who worked in hospital settings than among those in other settings; 33.8 percent of full-time hospital RNs reported that they worked paid overtime, with an average of 7.5 hours per week (Appendix A, Table 21). Mandatory overtime was reported by 8.2 percent of these RNs, averaging 6.4 hours per week. Among part-time RNs who worked in hospital settings, 19.9 percent worked paid overtime, with an average of 5.5 hours per week. Only 3.7 percent reported any mandatory overtime, with those who worked mandatory overtime averaging 4.5 hours per week (Appendix A, Table 22). On-call work was most prevalent in nonhospital work settings.

- The settings in which the fewest full-time RNs reported that they worked paid overtime were school health and insurance claims. In all other settings, at least 10 percent of RNs worked some paid overtime.
- After hospital-based RNs, full-time home health RNs were the second most likely group to report that they worked mandatory overtime (7.9 percent).

Full-time RNs in home health settings reported the highest rate of on-call hours (30.1 percent), with the second-highest percentage being among those in nursing homes and extended care facilities (15.9 percent) (Appendix A, Table 21).

- Full-time RNs in home health settings also had the highest rate of standby on-call hours, with 25.5 percent listing that they had some standby hours.
- Full-time RNs who worked in ambulatory care settings and had some standby hours (6.9 percent) reported an average of 40.9 standby hours per week. Less than half of these RNs were paid at a standby rate.

• Full-time home health RNs had the highest percentage of RNs whose standby hours were paid.

3.8.6 Hours Worked and Job Title

The number of hours that RNs work per week, which includes overtime and on-call hours worked, is associated with job title (Table 3-14).

| Job title | Mean hours/week full-time nurses | Mean hours/week part-time nurses |
|----------------------------|-------------------------------------|-------------------------------------|
| Staff nurse | 41.6 | 24.4 |
| First-line management | 46.3 | 24.2 |
| Middle/senior management | 48.1 | 25.3 |
| Consultant | 46.2 | 21.7 |
| Instruction | 43.8 | 21.1 |
| Nurse practitioner | 44.2 | 24.3 |
| Nurse midwife | 48.7 | 23.1 |
| Clinical nurse specialist | 43.9 | 25.8 |
| Nurse anesthetist | 45.1 | 26.4 |
| Researcher | 42.6 | 23.9 |
| Informatics | 45.6 | 21.7 |
| Surveyor/auditor/regulator | 42.2 | 24.7 |
| Patient coordinator | 43.2 | 24.6 |
| Patient educator | 40.0 | 21.3 |
| Other | 42.5 | 20.6 |

Table 3-14.Average hours per week in principal nursing position, by full-time/part-time status
and job title*

*Excludes registered nurses for whom job title was not known.

- Full-time staff nurses work an average of 41.6 hours per week, which is less than the average for all job titles except patient educators.
- The greatest number of hours per week is worked by full-time RNs with nurse midwife job titles (48.7 hours), followed by those with middle or senior management job titles (48.1 hours).
- Among part-time RNs, those with the greatest number of average hours per week are nurse anesthetists (26.4 hours), followed by clinical nurse specialists (25.8 hours), and nurses in middle or senior management positions (25.3 hours).

3.9 Employment Settings and Level of Care

RNs work in a variety of clinical settings, providing care at different levels of acuity or patient need. Tables 3-15, 3-16, and 3-17 present data on the level of care, types of patients, and clinical specialties of nurses. (Further detail can be found in Appendix A, Tables 23, 24, 25, and 26.)

| Level of care | Hospital setting (number) | Hospital setting (percent) | Nonhospital setting (number) | Nonhospital setting (percent) |
|--|---------------------------------|----------------------------------|------------------------------------|-------------------------------------|
| General or specialty inpatient | 595,742 | 37.9 | 101,768 | 10.8 |
| Critical/intensive care | 328,932 | 20.9 | 15,265 | 1.6 |
| Step down/transitional/progressive/telemetry | 187,973 | 12.0 | 8,707 | 0.9 |
| Subacute care | 44,131 | 2.8 | 25,258 | 2.7 |
| Emergency | 174,635 | 11.1 | 24,194 | 2.6 |
| Urgent care | 22,717 | 1.4 | 26,190 | 2.8 |
| Rehabilitation | 45,278 | 2.9 | 39,569 | 4.2 |
| Long-term care/nursing home | 17,610 | 1.1 | 134,362 | 14.2 |
| Surgery | 304,117 | 19.3 | 70,671 | 7.5 |
| Ambulatory care | 136,200 | 8.7 | 221,145 | 23.5 |
| Home health | 3,969 | 0.3 | 151,332 | 16.0 |
| Public/community health | 8,042 | 0.5 | 116,892 | 12.4 |
| Education | 61,002 | 3.9 | 117,291 | 12.4 |
| Business/administration/review | 93,509 | 5.9 | 133,281 | 14.1 |
| Other** | 43,548 | 2.8 | 30,819 | 3.3 |
| Total number of registered nurses | 1,571,861 | | 943,026 | |

| Table 3-15. | Levels of care in nursing work for hospital and nonhospital settings* |
|-------------|---|
|-------------|---|

*Only registered nurses who provided setting and level of care information are included in the calculations used for this table. Percents add to more than 100 as RNs may have reported more than one level of care.

**Other level of care/type of work includes ancillary care, research, informatics, flight nurse, and other.

Source: 2008 National Sample Survey of Registered Nurses

Among RNs who worked in hospital settings, the largest percentage (37.9 percent) worked in general or specialty inpatient environments, followed by 20.9 percent of RNs who worked in critical or intensive care, and 19.3 percent who worked in surgical services.

Among RNs in nonhospital settings, the most common nonhospital level of care is ambulatory care (23.5 percent), followed by home health (16 percent of nonhospital RNs). Nearly 11 percent of RNs who do not work in hospitals report that they spend the majority of their time in a general or specialty inpatient setting. The largest percentage of these RNs work in school health (35.3 percent) and academic education (21.4 percent).

RNs care for patients across the age span in all settings (Table 3-16). Among RNs who worked in hospital settings, more than half reported that they spend at least 50 percent of their patient care time with adult patients (56 percent). Geriatric patients are the second most commonly reported patient population in hospital settings, with 11.9 percent of RNs spending the majority of time with these patients. Slightly more than 10 percent of hospital-based RNs provided care for children.

Among RNs employed in nonhospital settings, adult patients consume roughly one-third of RN time for patient care. An additional 25 percent of RN time is devoted to geriatric patients. Pediatric and adolescent patients are the majority of patients for 13.4 percent of nonhospital RNs probably reflecting school based and home care nursing populations.

Table 3-16.Primary patient age group cared for in principal nursing position, for hospital and
nonhospital settings*

| Patient Type | Hospital setting (number) | Hospital setting (percent) | Nonhospital setting (number) | Nonhospital setting (percent) |
|------------------------|------------------------------|-------------------------------|------------------------------------|-------------------------------------|
| Adult | 891,482 | 56.0 | 342,703 | 35.5 |
| Geriatric | 189,969 | 11.9 | 240,652 | 25.0 |
| Prenatal | 34,119 | 2.1 | 13,116 | 1.4 |
| Newborn or neonatal | 88,723 | 5.6 | 6,368 | 0.7 |
| Pediatric/adolescent | 90,185 | 5.7 | 129,546 | 13.4 |
| Multiple age groups | 192,472 | 12.1 | 88,587 | 9.2 |
| No patient care | 104,312 | 6.6 | 143,255 | 14.9 |
| Total number of nurses | 1,591,261 | 100.0 | 964,227 | 100.0 |

*Only registered nurses who provided setting and patient type information are included in the calculations used for this table.

Source: 2008 National Sample Survey of Registered Nurses

Table 3-17 presents the clinical specialty in which RNs spend most of their time in their principal nursing position.

Among RNs who worked in hospital settings, the most common clinical area is general medical or surgical care, in which 29.3 percent of RNs provide care. Critical care and cardiac/cardiovascular care are common clinical specialties for hospital-employed RNs, with 19.2 and 16.7 percent of RNs, respectively.

Among RNs working in nonhospital settings, the most common clinical area is general medical or surgical care (21.1 percent), with primary care being reported by nearly as many RNs (19.7 percent). A relatively high percentage of nonhospital RNs report that their primary clinical area is chronic care

(13.6 percent), indicating that RNs may be providing care to chronically ill patients in homes and extended care facilities.

| Clinical Specialty | Hospital setting (number) | Hospital setting (percent) | Nonhospital setting (number) | Nonhospital setting (percent) |
|---------------------------------|---------------------------------|----------------------------------|------------------------------------|-------------------------------------|
| Critical care | 301,271 | 19.2 | 16,465 | 2.0 |
| Cardiac care | 262,175 | 16.7 | 46,518 | 5.7 |
| Chronic care | 60,237 | 3.8 | 110,942 | 13.6 |
| Emergency/trauma care | 186,451 | 11.9 | 31,888 | 3.9 |
| Gastrointestinal | 85,806 | 5.5 | 34,610 | 4.2 |
| General medical surgical | 459,315 | 29.3 | 172,075 | 21.1 |
| Gynecology (women's health) | 60,739 | 3.9 | 37,878 | 4.6 |
| Hospice | 22,459 | 1.4 | 58,713 | 7.2 |
| Infectious/communicable disease | 50,647 | 3.2 | 44,084 | 5.4 |
| Labor and delivery | 86,271 | 5.5 | 8,180 | 1.0 |
| Neurological | 67,151 | 4.3 | 17,771 | 2.2 |
| Obstetrics | 102,300 | 6.5 | 30,296 | 3.7 |
| Oncology | 89,611 | 5.7 | 29,919 | 3.7 |
| Primary care | 57,861 | 3.7 | 161,041 | 19.7 |
| Psychiatric/mental health | 79,638 | 5.1 | 53,130 | 6.5 |
| Pulmonary/respiratory | 85,602 | 5.5 | 34,774 | 4.3 |
| Renal/dialysis | 57,094 | 3.6 | 31,295 | 3.8 |
| Other specialty** | 95,595 | 6.1 | 140,565 | 17.2 |
| No patient care | 17,526 | 1.1 | 42,628 | 5.2 |
| Total number of nurses | 1,566,645 | | 816,982 | |

| Table 3-17. | Primary clinical specialty in principal nursing position for hospital and nonhospital |
|-------------|---|
| | settings* |

*Only registered nurses who provided setting and clinical specialty information are included in the calculations used for this table. Percents add to more than 100 as registered nurses may report more than one clinical specialty.

**Other clinical specialty includes dermatology, endocrinology, infusion, occupational health, ophthalmology, orthopedic, otolaryngology, plastic/cosmetic surgery, radiology, urology, no specific area, and other specialties.

Source: 2008 National Sample Survey of Registered Nurses

3.9.1 Education and Areas of Clinical Specialty

An RN's highest nursing or nursing-related education is associated with the clinical specialty in which the RN works (Table 3-18).

- General medical-surgical care is the most common specialty of RNs, regardless of education. However, the percentage in this area is lower for RNs with graduate degrees.
- RNs with master's degrees and doctorates are more likely to work in primary care (19.2 percent) than are RNs at other education levels. They also are more likely to work in gynecology/women's health (6.2 percent), obstetrics (6.9 percent), and mental health

(8.9 percent). These are fields in which advanced practice registered nurses with graduate degrees often have specific preparation.

- RNs with bachelor's degrees are more likely to work in critical care (17.4 percent), cardiac care (15.0 percent), labor and delivery (4.6 percent), neurological (4.1 percent), and oncology (6.1 percent) than are RNs with other education preparation.
- RNs with ADNs are more likely to work in general medical surgical care (29.3 percent), chronic care (8.5 percent), emergency/trauma (10.7 percent), hospice (4.1 percent), infectious disease (4.3 percent), pulmonary/respiratory (5.7 percent), and renal/dialysis (4.6 percent) than are RNs with other education preparation.
- RNs with diplomas are somewhat more likely to work in gastrointestinal (6 percent) than are other RNs. More than 10 percent work in primary care.

Table 3-18.Primary clinical specialty in principal nursing position, by highest nursing and
nursing-related education*

| | Diploma | Associate degree | Bachelor's degree | Master's or doctorate |
|---------------------------------|-----------|---------------------|----------------------|--------------------------|
| Clinical Specialty | (percent) | (percent) | (percent) | (percent) |
| Critical care | 10.6 | 12.8 | 17.4 | 9.5 |
| Cardiac care | 11.1 | 14.1 | 15.0 | 9.4 |
| Chronic care | 7.6 | 8.5 | 6.5 | 7.2 |
| Emergency/trauma care | 7.6 | 10.7 | 9.7 | 7.3 |
| Gastrointestinal | 6.0 | 5.7 | 5.2 | 3.3 |
| General medical surgical | 27.6 | 29.3 | 27.2 | 23.9 |
| Gynecology (women's health) | 4.8 | 4.1 | 3.8 | 6.2 |
| Hospice | 3.8 | 4.1 | 3.3 | 2.3 |
| Infectious/communicable disease | 3.7 | 4.3 | 4.2 | 4.1 |
| Labor and delivery | 3.7 | 4.0 | 4.6 | 3.6 |
| Neurological | 2.7 | 3.8 | 4.1 | 3.5 |
| Obstetrics | 6.6 | 5.1 | 5.9 | 6.9 |
| Oncology | 4.7 | 4.5 | 6.1 | 5.2 |
| Primary care | 10.4 | 8.4 | 7.8 | 19.2 |
| Psychiatric/mental health | 5.9 | 6.4 | 4.3 | 8.9 |
| Pulmonary/respiratory | 4.9 | 5.7 | 5.5 | 3.6 |
| Renal/dialysis | 2.9 | 4.6 | 4.0 | 2.2 |
| No patient care | 3.1 | 2.1 | 2.7 | 4.1 |
| Other specialty** | 12.0 | 10.3 | 9.8 | 10.5 |

*Only registered nurses who provided clinical specialty and highest education information are included in the calculations used for this table. Percents add to more than 100 as registered nurses may have reported more than one clinical specialty.

**Other clinical specialty includes dermatology, endocrinology, infusion, occupational health, ophthalmology, orthopedic, otolaryngology, plastic/cosmetic surgery, radiology, urology, no specific area, and other specialties.

3.10 Composition of Nursing Work

RNs spend the majority of their time performing direct patient care and recordkeeping related to patient care called "charting" (Figure 3-15 and Appendix A, Table 27). Nearly half of RNs (46.8 percent) spend at least 75 percent of their day in these activities, and nearly 70 percent spend at least half their day in this area.





Non-nursing tasks, which include housekeeping and searching for supplies do not appear to be significant tasks for RNs; only 3.3 percent of RNs reported that non-nursing activities occupy 25 percent or more of their time.

Consultation with other professionals and agencies occupies at least 25 percent of the time of 7 percent of RNs. Nearly 40 percent of RNs do not engage in consultation at all (38.9 percent).

Source: 2008 National Sample Survey of Registered Nurses

The Employment of Registered Nurses

3

Teaching, precepting, or orienting students or new hires occupies at least some work time of 52.6 percent of RNs. Only 6.9 percent of RNs say they spend at least 25 percent of their time in teaching-related activities.

3.10.1 Composition of Work and Job Title

The activities of RNs vary by job title (Figure 3-16).

- Staff nurses spend an average of 71.8 percent of their time performing patient care and keeping patient care records. Non-nursing tasks occupy an average of 7.6 percent of staff nurse time.
- RNs in first-line management positions spend an average of 45 percent of their time supervising others and 25.3 percent of their time providing patient care and charting.
- Middle and senior managers spend an average of 38.3 percent of their time in supervision, 25 percent performing administrative responsibilities, and 11.5 percent providing patient care and charting. An average of 9.9 percent of their time is spent consulting with other providers or agencies.
- Nurse practitioners spend an average of 79.7 percent of their time providing patient care and charting patient care information.

3.10.2 Composition of Work and Education

The highest nursing or nursing-related education of RNs is associated with the activities performed by RNs (Table 3-19 and Appendix A, Table 28). While RNs with graduate degrees spend less time doing patient care and charting on average than do other RNs, this is probably associated with the fact that RNs who are managers and educators would have less patient care activities than staff nurses. Only nurse practitioners, nurse midwives, and nurse anesthetists would have patient contact similar to staff nurses (Table 3-5).




Figure 3-16. Time spent by registered nurses in various activities, by job title

*Excludes registered nurses for whom job title was not known.

Source: 2008 National Sample Survey of Registered Nurses

| Table 3-19. | Average percent time spent on various activities, by highest nursing or nursing- |
|-------------|--|
| | related education |

| | | Associate | Bachelor's | Master's | |
|-----------------------|---------|-----------|------------|----------|-----------|
| Functional Area | Diploma | degree | degree | degree | Doctorate |
| Patient care/charting | 60.3 | 61.4 | 58.1 | 48.2 | 18.1 |
| Non-nursing tasks | 6.0 | 6.9 | 6.5 | 3.0 | 1.6 |
| Consultation | 7.2 | 6.7 | 7.9 | 8.9 | 8.8 |
| Supervision | 10.3 | 9.9 | 10.2 | 10.7 | 8.7 |
| Administration | 3.3 | 3.2 | 3.9 | 9.7 | 19.6 |
| Research | 1.5 | 1.7 | 1.9 | 2.5 | 9.7 |
| Teaching | 5.6 | 6.1 | 7.0 | 12.4 | 26.7 |
| Other | 6.0 | 4.1 | 4.6 | 4.7 | 6.7 |

3.11 Temporary and Travel Work

An estimated 88,495 RNs, or 3.4 percent of all employed RNs, are employed through temporary employment agencies. Temporary agencies may provide either temporary local placements for RNs, often on a per-shift basis, or short- to mid-term placements with nonlocal employers. The latter type of agency focuses on "travel" placements that typically last from 3 months to 1 year and often occur in a State other than where the travel RN resides.

About 2.5 percent of RNs employed in hospital settings report that they are employed by a temporary service for their principal nursing position, and 1 percent say they are employed by a temporary service for at least one secondary nursing position (but not their principal position) (Table 3-20 and Appendix A, Table 29). Temporary agency employment is somewhat more common among RNs whose principal nursing position is in a hospital.

Table 3-20.Employment by temporary employment services (TES), for primary and secondary
nursing positions, by principal nursing position in a hospital

| | Hospital setting (percent) | Nonhospital setting (percent) |
|---------------------------------------|-------------------------------|----------------------------------|
| TES principal position only | 2.5 | 2.0 |
| TES secondary position only | 1.0 | 1.1 |
| Both principle and secondary position | 0.1 | 0.1 |
| Not employed by TES | 96.4 | 96.8 |
| Total | 100.0 | 100.0 |

Source: 2008 National Sample Survey of Registered Nurses

More than half of RNs who have secondary employment through a nontravel temporary placement agency report that they are placed in a hospital (Table 3-21).

Table 3-21.Employment settings of secondary employment through a nontravel temporary
employment service

| Employment setting | Number | Percent |
|---------------------|--------|---------|
| Hospital | 13,096 | 53.8 |
| Nonhospital setting | 11,241 | 46.2 |
| Total | 24,337 | 100.0 |

3.12 Crossing State Lines and Compact States

As of December 2008, 23 States participated in the Nurse Licensure Compact. A multi-State or mutual recognition licensure model, the Compact allows a nurse to have an RN license in the State where she or he resides and practice with that license in other States that are part of the Compact. Thus, an RN may practice across State lines unless otherwise restricted. RNs who do not reside in a Compact State must have a separate license for each State in which they work. Advanced practice nursing licenses are not part of the Compact model in any state.

An estimated 824,662 RNs live in a Compact State (Table 3-22). Of these, 96 percent live and work in the same State, 1.6 percent work in a different State that is part of the Compact, and 2.4 percent work in a State different from that in which the nurse lives, which is not a Compact State.

Table 3-22. Licensure in a Compact State, by where registered nurses live and work

| | Number | Percent |
|---|---------|---------|
| Live and work in same Compact State | 791,444 | 96.0 |
| Live in one Compact State and work in another Compact State | 13,537 | 1.6 |
| Live in a Compact State and work in non-Compact State | 19,681 | 2.4 |
| Total | 824,662 | 100.0 |

Source: 2008 National Sample Survey of Registered Nurses

3.13 Unionization in Principal Nursing Positions

Table 3-23 presents unionization by employment setting for registered nurses' principal nursing positions. Among RNs who work in hospital settings, 18.6 percent are unionized.

Unionization is most prevalent among RNs whose principal positions are in school health (40.4 percent), public health (26.0 percent), and academic education (19.2 percent). This is consistent with the comparatively high rate of unionization in school systems, among academic faculty, and among public sector and government employees.



| Employment setting | Unionized (percent) |
|----------------------------|------------------------|
| Hospital | 18.6 |
| Nursing home/extended care | 5.8 |
| Academic education | 19.2 |
| Home health | 5.9 |
| Public/community health** | 26.0 |
| School health | 40.4 |
| Ambulatory care | 6.0 |
| Other*** | 7.3 |

*Only registered nurses who provided employment setting information are included in the calculations used for this table.

**Public/community health includes occupational health.

***Other includes insurance/benefits/utilization review.

Table 3-23.

Source: 2008 National Sample Survey of Registered Nurses

3.14 Earnings of Registered Nurses

Average annual earnings for full-time employed RNs were \$66,973 in 2008, rising 15.9 percent since the 2004 average of \$57,785 (Figure 3-17 and Appendix A, Table 32). When annual earnings are adjusted for inflation using the Consumer Price Index (CPI), earnings in 2008 were \$26,826, which is only a 1.7 percent increase from average real (inflation-adjusted to a 1980 base) 2004 earnings of \$26,366. Thus, growth in earnings of full-time RNs between 2004 and 2008 only slightly outpaced inflation.

Total annual earnings from nursing work are closely associated with the total number of hours RNs work per year in principal and secondary positions (Table 3-24). In general, total earnings from all nursing positions increase with the number of hours worked in these positions, up to 3,500 hours per year.




Figure 3-17. Actual and inflation-adjusted "real" average earnings for full-time registered nurses in their principal nursing position*

*Only registered nurses who provided earnings information are included in the calculations used for this figure.

Source: 1980-2008 National Sample Survey of Registered Nurses

Table 3-24. Average annual earnings from all nursing positions, by total number of hours worked per year in all nursing positions*

| Hours per year | Average annual earnings |
|-----------------------|-------------------------|
| 1 to 499 hours | 13,850 |
| 500 to 999 hours | 26,271 |
| 1,000 to 1,499 hours | 42,582 |
| 1,500 to 1,999 hours | 58,140 |
| 2,000 to 2,499 hours | 68,327 |
| 2,500 to 2,999 hours | 80,537 |
| 3,000 to 3,499 hours | 92,254 |
| 3,500 to 3,999 hours | 85,117 |
| More than 4,000 hours | 93,865 |

*Only registered nurses who provided earnings information are included in the calculations used for this table.

3.14.1 Earnings and Job Titles

The average annual earnings of full-time RNs are reported by job title in Table 3-25 and Appendix A, Table 30.

- The highest earnings are reported by nurse anesthetists, who average \$154,221 per year.
- The lowest earnings are reported by patient educators, at \$59,421 per year.
- Staff nurses earn \$61,706 per year, on average.
- Nurse practitioners average \$85,025 per year, while nurse midwives average \$82,111.
- RNs whose positions involved instruction of health professions students average \$65,844 per year.

Table 3-25.Average annual earnings, by job title of principal nursing position, for registered
nurses working full-time*

| Job title | Average annual earnings |
|----------------------------|-------------------------|
| Staff nurse | 61,706 |
| Management/administration | 78,356 |
| First-line management | 72,006 |
| Middle management | 74,799 |
| Senior management | 96,735 |
| Nurse anesthetist | 154,221 |
| Clinical nurse specialist | 72,856 |
| Nurse midwife | 82,111 |
| Nurse practitioner | 85,025 |
| Patient educator | 59,421 |
| Instruction | 65,844 |
| Patient coordinator | 62,978 |
| Informatics nurse | 75,242 |
| Consultant | 76,473 |
| Researcher | 67,491 |
| Surveyor/auditor/regulator | 65,009 |
| Other** | 64,003 |
| Total | 66,973 |

*Only registered nurses who provided earnings and job title information are included in the calculations used for this table.

**Other job title includes registered nurses for whom job title is unknown.

3.14.2 Earnings and Employment Setting

Earnings of RNs vary across the setting of their principal nursing position (Table 3-26 and Appendix A, Table 31).

- RNs who work full-time in hospitals earn an average of \$69,079 per year and staff nurses in hospitals earn an average of \$63,809 per year.
- The setting with the lowest average pay for full-time RNs is school health service, although this may be because school health RNs typically work only an academic year rather than a full calendar year.

Table 3-26.Average annual earnings, by employment setting of principal nursing position, for
registered nurses working full-time*

| Employment setting | Overall average | Staff nurse |
|--|-----------------|-------------|
| Hospital | 69,079 | 63,809 |
| Nursing home/extended care facility | 61,507 | 54,239 |
| Academic education program | 66,701 | 66,542 |
| Home health setting | 63,490 | 59,289 |
| Public or community health setting | 60,580 | 55,708 |
| School health service | 47,851 | 46,416 |
| Occupational health | 65,947 | 62,086 |
| Ambulatory care setting (not hospital) | 65,976 | 55,657 |
| Insurance claims/benefits/utilization review | 67,897 | 61,370 |
| Other** | 72,750 | 67,660 |
| Total | 66,973 | 61,706 |

*Only registered nurses who provided earnings and setting information are included in the calculations used for this table.

**Other employment setting includes registered nurses for whom employment setting is unknown.

Source: 2008 National Sample Survey of Registered Nurses

3.14.3 Earnings From Secondary Jobs

Secondary nursing positions are held by 12.6 percent of RNs employed in nursing. Appendix A, Table 19 presents details about the earnings of RNs in their principal and secondary nursing positions, according to whether they hold only one nursing job or multiple nursing jobs.

 RNs who work full-time in their principal position and have multiple nursing jobs earn more in their principal position (\$68,618) than do RNs who do not have multiple jobs (\$66,742).

• Average earnings in secondary positions are \$15,201 for RNs whose principal position is full-time and \$13,392 for those with a part-time principal position.

These data indicate that RNs who hold multiple positions earn more overall and in their principal positions. The reasons for this are not known based on 2008 NSSRN data findings. RNs with two or more positions earn an average of \$74,894 annually from all nursing jobs. For those whose principal position is full-time, the average for all earnings is \$83,818 per year.

3.14.4 Earnings and Experience

RNs employed full-time receive higher earnings from their principal nursing position as they gain experience up to 30 years (Table 3-27). However, after 30 years of experience, average annual earnings decline, largely due to older RNs working fewer hours. Nonetheless, earning differentials based on years of experience are relatively modest. RNs at the height of their earning potential with 30 years experience earn, on average, only \$19,281 more than RNs with less than 5 years experience.

| Table 3-27. | Average annual earnings for registered nurses working full-time in their princip | |
|-------------|--|--|
| | nursing position, by years since initial nursing education* | |

| Years since initial nursing education | Average annual earnings |
|---------------------------------------|-------------------------|
| Less than 5 years | 56,118 |
| 6 to 10 years | 65,604 |
| 11 to 15 years | 67,212 |
| 16 to 20 years | 69,388 |
| 21 to 25 years | 71,216 |
| 26 to 30 years | 75,399 |
| 31 to 35 years | 73,733 |
| 36 to 40 years | 71,551 |
| More than 40 years | 70,358 |

*Only registered nurses who provided earnings information are included in the calculations used for this table.

Source: 2008 National Sample Survey of Registered Nurses

3.14.5 Earnings and Education

The highest nursing or nursing-related education level of RNs is associated with job titles and subsequent earnings. Some RNs with graduate degrees earn substantially more than do other nurses (Table 3-28 and Appendix A, Table 32). In part, they earn more because they command high earnings when they hold management job titles or are in advanced specialties. The earnings of RNs

who work as faculty are somewhat less than other RNs with advanced degrees. RNs working as faculty are discussed further in Chapter 4.

- RNs with graduate degrees earn an average of at least \$20,000 more than do RNs with other levels of education.
- Staff nurses with graduate degrees average about \$6,000 more than other staff RNs.
- Diploma RNs earn, on average, more than ADN nurses but not more than bachelor's degree RNs for the same job titles.

Table 3-28. Average annual earnings of registered nurses working full-time in their principal nursing position, by highest nursing or nursing-related education and job title*

| | | Associate | Bachelor's | Master's/ | Overall |
|--------------------------|---------|-----------|------------|-----------|---------|
| Job title | Diploma | degree | degree | doctorate | average |
| All nurses | 65,349 | 60,890 | 66,316 | 87,363 | 66,973 |
| Staff nurse | 63,027 | 59,310 | 63,382 | 69,616 | 61,706 |
| First-line management | 68,089 | 66,138 | 75,144 | 85,473 | 72,006 |
| Senior/middle management | 74,090 | 69,871 | 79,878 | 101,730 | 81,391 |
| Patient coordinator | 62,693 | 60,240 | 64,068 | 71,516 | 62,978 |

*Only registered nurses who provided earnings and job title information are included in the calculations used for this table.

Source: 2008 National Sample Survey of Registered Nurses

3.15 Job Satisfaction

The 2004 and 2008 surveys asked RNs to report their satisfaction with their principal job. In 2008, 29.3 percent of RNs reported that they were extremely satisfied with their principal nursing position, compared with 27.5 percent in 2004. In 2004, 50.5 percent where moderately satisfied while in 2008, 51.8 percent were moderately satisfied (Figure 3-18). Only 11.1 percent were dissatisfied in 2008, compared with 13.8 percent in 2004. While the differences from 2004 to 2008 are small, RN job satisfaction may be somewhat improved.

The Employment of Registered Nurses

3




Figure 3-18. Job satisfaction of registered nurses*

* Only registered nurses who provided job satisfaction information are included in the calculations used for this figure.

3.15.1 Job Satisfaction and Employment Setting

Employment setting is associated with overall satisfaction. Appendix A, Table 33 presents satisfaction ratings for RNs according to the setting of their principal nursing position. Figure 3-19 provides data for the most common employment settings.

- Registered nurses who work in academic education programs, ambulatory care, and home health settings, although paid less than RNs in other settings, reported the highest rates of job satisfaction (86.6 percent, 85.5 percent, and 82.8 percent, respectively).
- Working in nursing homes/extended care was associated with the lowest rate of RNs being moderately or extremely satisfied with their work (74.5 percent). These RNS also reported the most dissatisfaction with their work (moderate or extreme dissatisfaction at 16.9 percent; extreme dissatisfaction at 4.8 percent).
- Almost 12 percent of RNs employed in hospitals reported moderate or extreme dissatisfaction (11.8 percent).

3.15.2 Job Satisfaction and Job Titles

Registered nurses' satisfaction with their jobs also varies with their job title. Appendix A, Table 34 provides information for all job titles; Figure 3-20 provides data for the largest groups.

- Staff nurses were less likely to report that they were moderately or extremely satisfied than RNs with other job titles.
- Being "extremely satisfied" was reported more often by RNs in senior or middle management (39.2 percent) or other job titles (40.2 percent). The "other job titles" category includes advanced practice fields and education.
- Moderate or extreme dissatisfaction was reported more often by RNs in first-line management positions (12.3 percent) and staff nurses (12 percent).
- "Extremely dissatisfied" was reported more often by RNs in senior or middle management positions (2.9 percent).

These data suggest that RNs in positions of the most autonomy and agency, middle managers, faculty, and advanced practice registered nurses, were most likely to be extremely satisfied with their jobs. Staff nurses were least likely to be fully satisfied with their positions.




Figure 3-19. Job satisfaction, by employment setting of principal nursing position*

*Only registered nurses who reported employment setting information are included in the calculations used for this figure. Percents may not add to100 due to the effect of rounding.

**Public/community health includes school and occupational health.

***Other includes insurance, benefits, and utilization review, and other settings.




Figure 3-20. Job satisfaction, by job title in principal nursing position*

*Only registered nurses who provided job title information are included in the calculations used for this figure.

3.15.3 Job Satisfaction and Age

RN satisfaction generally increases with age (Figure 3-21 and Appendix A, Table 35). This may be because those who are dissatisfied have changed to positions or employment situations which are more satisfying for them.

- The percentage of employed RNs who are extremely satisfied continues to increase with age through age 74 years.
- The percent of employed RNs who are moderately to extremely satisfied averages around 78 percent for RNs under 35 years old. It averages about 80 percent for RNs who are ages 35 to 49 years. About 82 percent of RNs in their 50s report that they are moderately to extremely satisfied, and after age 60 job satisfaction rises to more than 85 percent.
- Moderate and extreme dissatisfaction is greatest between ages 40 and 50, averaging 12.4 and 12.5 percent, respectively.

This should not necessarily be interpreted to mean that age is a factor in job satisfaction. Rather, perhaps young RNs or recent graduates who are dissatisfied with a position may change jobs until they find a position that is more satisfactory. Registered nurses who cannot find a satisfactory position may leave nursing altogether.

3.15.4 Job Satisfaction and Education

Figure 3-22 and Appendix A, Table 36 present data on RNs' job satisfaction according to their highest nursing or nursing-related education.

- RNs with graduate-level education report the highest rates of being extremely satisfied (39.2 percent).
- A high percentage of diploma-educated RNs (31.4 percent) are extremely satisfied; diploma-educated RNs also tend to be older than other nurses, and age is correlated with satisfaction.
- RNs with ADNs most often report being moderately to extremely dissatisfied (13.1 percent).





Figure 3-21. Job satisfaction of registered nurses employed in nursing, by age

Figure 3-22. Job satisfaction of registered nurses employed in nursing, by highest nursing or nursing-related education*



*Only registered nurses who provided information on education preparation are included in the calculations used for this figure.

3.16 Job Changes of Registered Nurses

The 2008 NSSRN asked RNs to provide information about their employment status in 2007. Nearly 80 percent of RNs were employed in nursing in both 2007 and 2008 (Figure 3-23). Another 12.3 percent were not employed in nursing in either year. Nearly 3 percent were employed in nursing in 2007, but not in 2008. Conversely, 2.4 percent were employed in 2008 but had not been employed in 2007. Of those who were employed in 2007 but not in 2008, 27.3 percent (24,430) reported that they stopped working due to retirement. A group comprising 2.5 percent of RNs graduated in 2007 or 2008 and was employed only in 2008. These data indicate that departures from nursing work are often short-lived.

Table 3-29 provides more information about principal nursing employment in 2008 compared with 2007.

- Among RNs who worked full-time in 2007, 91.5 percent continued to work full-time in 2008. Another 5.8 percent worked part-time in 2008.
- Among RNs who worked part-time in 2007, 82.9 percent continued to work part-time in 2008, while 11.2 percent changed to full-time work.

Employment changes of RNs who were employed in both 2007 and 2008 are described in Table 3-30 and Appendix A, Table 37. Many RNs change positions and employers from year to year.

- Among RNs who worked full-time in 2007, 11.6 percent changed employer by 2008.
 Another 6.5 percent worked with the same employer but held a different position.
- Among RNs who worked part-time in 2007, 9.7 percent changed employers. These data suggest that part-time RNs are slightly less likely to change employers or positions than are full-time RNs.

Appendix A, Table 38 presents the employment settings of RNs in 2008 as compared with 2007. Among RNs employed in hospital settings in 2008, 90.8 percent had been employed in a hospital in 2007. However, other settings draw many RNs from hospitals. For example, while 82.5 percent of RNs employed in home health in 2008 had worked in that setting in 2007, 7.9 percent had worked in a hospital in 2007. Among RNs employed in ambulatory care in 2008, 8.2 percent had worked in a hospital in 2007.




Figure 3-23. Comparison of employment status of registered nurse population in 2007 and 2008

Source: 2008 National Sample Survey of Registered Nurses

Table 3-29.Employment status in 2008, compared with status in 2007

| | Year 2007 employed in nursing full-time (number) | Year 2007 employed in nursing full-time (percent) | Year 2007 employed in nursing part-time (number) | Year 2007 employed in nursing part-time (percent) |
|------------------------------------|--|---|--|---|
| 2008 employed in nursing full-time | 1,753,305 | 91.5 | 69,166 | 11.2 |
| 2008 employed in nursing part-time | 110,737 | 5.8 | 513,551 | 82.9 |
| 2008 not employed in nursing | 52,628 | 2.7 | 36,773 | 5.9 |



| | Year 2007 employed in nursing full- time (number) | Year 2007 employed in nursing full- time (percent) | Year 2007 employed in nursing part- time (number) | Year 2007 employed in nursing part- time (percent) |
|---|---|--|---|--|
| Same position and employer in 2008 | 1,525,839 | 81.9 | 497,398 | 85.4 |
| Different position, same employer in 2008 | 120,474 | 6.5 | 28,764 | 4.9 |
| Different employer in 2008 | 216.327 | 11.6 | 56,430 | 9.7 |

Table 3-30.Employment changes among registered nurses employed in nursing in both 2007
and 2008

Source: 2008 National Sample Survey of Registered Nurses

Workplace issues are significant reasons for job change among RNs. The reasons RNs changed principal employment are presented in detail in Appendix A, Table 39. RNs could indicate multiple reasons for a change of employment.

- More than 73 percent of RNs reported that they changed positions or employers due at least in part to workplace issues such as stressful work environment (29.6 percent), lack of good management (27.8 percent), or inadequate staffing (20.1 percent).
- Personal career reasons, often involving advancement, were a factor in job changes for 37.5 percent of RNs. Such reasons included interest in another position (29.5 percent) and improved pay or benefits (19.2 percent).
- Nearly 30 percent of RNs changed jobs at least in part for personal family reasons, which included relocation (14.9 percent), disability (1.3 percent), and unspecified personal/family issues (15.4 percent).

3.17 Plans for Future Employment

Employed RNs were asked if they planned to leave their principal nursing position within 3 years and if so, whether they intended to continue to work in nursing. Among RNs under 55 years old, the percentage of RNs that intended to leave nursing within 3 years is quite small, at 3 percent or less (Figure 3-24). Among RNs 55 and older who work in nursing, 76,915 (12.5 percent) intend to leave the nursing profession within 3 years; another 54,539 (8.9 percent) intend to leave their current nursing jobs and are unsure if they will remain in nursing afterward. Overall, a high proportion of RNs in all age groups intend to stay in nursing.




Figure 3-24. Intentions to stay in nursing over the next 3 years, by age group*

*The total percent of registered nurses across age groups may not equal the estimated total of RNs due to the effect of rounding.

**Stay in nursing includes registered nurses who do not plan to leave their current nursing position in the next 3 years, those who plan to leave their current position but stay in nursing, and those who are undecided about staying in their current nursing position who were not asked about their intention regarding nursing work in general.

Source: 2008 National Sample Survey of Registered Nurses

3.18 Non-Nursing Jobs

Some RNs are employed in positions outside nursing, including those who also hold a nursing position. RNs who work in non-nursing positions report a variety of fields of employment. The largest percentage (31.4 percent) is employed in non-nursing health-related service fields (Figures 3-25 and 3-26 and Appendix A, Table 42). Other common fields of employment are retail

and food sales and services (14.4 percent), elementary and secondary education (10.4 percent), and pharmaceuticals and medical equipment (9.0 percent).

Among all RNs, an estimated 208,427 held a non-nursing position. Of the RNs employed outside of nursing, 97,365 also had a nursing position (Table 3-31). Among RNs not working in nursing, 23.8 percent held a non-nursing position.

More than half of RNs (54.7 percent) who are not working in nursing and hold a non-nursing job report that their position is health-related (Appendix A, Table 41). The percent of RNs who do not work in nursing and have a non-nursing position generally increases with the number of years since they last worked in nursing (Figure 3-25). Forty percent of RNs who have not worked in nursing for 20 years or more hold non-nursing jobs.

- Most RNs who work in health-related occupations that are non-nursing work full-time (73.7 percent) (Appendix A, Table 41).
- Only half of RNs who work in nonhealth-related occupations work full-time (50.6 percent).

RNs hold many job titles in their non-nursing fields of employment (Figure 3-27 and Appendix A, Table 43).

- More than 75 percent of RNs who work outside nursing and have a management job title say their position is health-related.
- Only 26.8 percent of RNs who work outside nursing and are business owner/proprietors say their work is health-related.




Figure 3-25. Non-nursing positions among registered nurses not employed in nursing, by years since last worked in nursing




Figure 3-26. Employment settings of registered nurses employed outside of nursing*

*Excludes registered nurses who are employed in nursing but have other employment outside of nursing.

**Other includes agriculture/farming/ranching, fine arts/performing arts, emergency response, fitness/sport, religious organizations, and others.

Source: 2008 National Sample Survey of Registered Nurses

Table 3-31. Employment in non-nursing occupations, by nursing employment status*

| | Licensed nurses (number) | Working in non-nursing occupations (number) | Working in non-nursing occupations (percent) |
|-------------------------|-----------------------------|---|--|
| Employed in nursing | 2,596,599 | 97,365 | 3.7 |
| Not employed in nursing | 466,564 | 111,062 | 23.8 |
| Total | 3,063,162 | 208,427 | 6.8 |

*Includes an estimated 5,073 registered nurses who are employed in non-nursing and are actively seeking nursing employment. Excludes an estimated 97,365 registered nurses who are employed in nursing and have other employment in non-nursing.




Figure 3-27. Job titles of registered nurses employed outside of nursing*

*Excludes registered nurses who are employed in nursing but have other employment outside of nursing.

**Other includes such job titles as emergency response, clinical research, construction, clergy, chef, IT, etc.

4.1 Introduction

Registered nurses who are nursing education program faculty fill an important role in the nursing workforce and help to ensure that there are adequate numbers of new nurses entering the profession. RNs can serve as faculty or instructors in both their principal and secondary nursing positions. Most RNs whose principal position is in a full-time faculty position teach theoretical and didactic material in the classroom as well as provide clinical education. Those whose faculty work is part-time usually focus on clinical education by teaching and supervising students in clinical settings, or by teaching only one or two didactic courses a year. Depending on the type of nursing program (ADN, diploma, BSN, or higher degree) faculty may be required to teach or may also have other faculty responsibilities such as research, producing scholarly publications, engaging in community and professional organization service, serving on academic committees, and advising students. Faculty members who teach in nurse practitioner programs must also continue clinical practice to maintain national certification and/or State licensure as Advanced Practice Registered Nurses (APRNs) in their area of clinical specialty, which is usually a requirement for RNs who teach clinical courses in nurse practitioner programs.

This chapter includes not only those RNs who reported that their principal position was instructor/lecturer or professor and that their setting was an academic RN program, but also those RNs who reported a secondary nursing job in an academic education program. Nurses whose principal job title was managerial, administrative, or related to patient care in an academic setting are not included as faculty. Nurses with these titles do not primarily provide instruction to nursing students; rather, they provide patient care within academic medical centers, manage education programs, and have other nonteaching responsibilities. Nurses who reported that at least one of their secondary nursing positions is in an academic setting are assumed to have a secondary position as faculty; job titles were not reported for secondary positions.

4.2 Number of Nurses Working in Faculty Roles

In 2008 an estimated 31,056 RNs worked as faculty in their principal nursing position. This number is smaller than the total who worked in academic settings because nurses can work in academic settings while also providing patient care or working in administrative roles (Table 4-1). The majority of RNs in faculty roles worked in programs that provide education at the BSN and higher level (60 percent). Another 38.3 percent worked in ADN or LPN/LVN education programs.

Table 4-1.Registered nurses employed as faculty in their principal nursing position, by type of
nursing education program*

| | Number | Percent |
|--|--------|---------|
| Associate and LPN/LVN program | 11,901 | 38.3 |
| Bachelor's and higher, Associate and BSN program | 18,626 | 60.0 |

*Registered nurses who work in diploma nursing programs were not included in the table because there were too few respondents to estimate the data accurately.

Source: 2008 National Sample Survey of Registered Nurses

An additional 34,666 RNs worked in academic education settings as a secondary nursing position (Table 4-2). More than 45 percent of RNs with secondary positions in academic settings reported that their job title for their principal nursing position is staff nurse. Nurses whose job title is management or nurse practitioner accounted for another 15 percent each. Just over 12 percent of nurses whose job title was in instruction also held a secondary position in an academic setting.

| Table 4-2. | Registered nurses employed as faculty in their secondary nursing position(s), by job |
|------------|--|
| | title of principal nursing position* |

| | Number | Percent |
|--------------------|--------|---------|
| Management | 5,395 | 15.6 |
| Instruction | 4,230 | 12.2 |
| Staff nurse | 15,743 | 45.4 |
| Nurse practitioner | 5,455 | 15.7 |
| Other** | 3,843 | 11.1 |
| Total | 34,666 | 100.0 |

*Only registered nurses who provided job title information are included in the calculations for this table.

**Other includes consultant, nurse midwife, CNS, NA, researcher, informatics, patient coordinator, and patient educator.

4.3 Faculty and Age

More than 40 percent of RNs working as faculty in their principal nursing position are between the ages of 50 to 59 (Figure 4-1). More than 19 percent of RNs whose principal position is as faculty were 60 years or older in 2008. Therefore, almost 60 percent of nursing faculty are over age 50. RNs whose secondary position is in an academic setting tend to be younger; among RNs under 50 years old, more worked in academic settings as a secondary position than as their primary position. Only 8.7 percent of RNs who have a secondary position in academic settings were 60 years or older. Nonetheless, the lack of younger RNs in faculty positions is apparent.




Figure 4-1. Age distribution of registered nurses who work as faculty

4.4 Faculty and Education

Most RNs who worked as faculty in their principal nursing positions had a master's or doctorate degree (Figure 4-2). Education levels were lower among RNs who worked as faculty as their secondary nursing position.




Figure 4-2. Highest education of registered nurses who work as faculty*

*Only registered nurses who provided education preparation information are included in the calculations used for this figure. Source: 2008 National Sample Survey of Registered Nurses

The education level of faculty is associated with the type of nursing program in which they work. RNs who worked in ADN and diploma programs (and also who prepare licensed practical nurses,) had lower education levels, on average, than did those who worked in programs that prepare nurses at the bachelor's and higher level (Figure 4-3). This is because most RN education programs at the bachelor's degree and higher level require full-time, tenure-track faculty to be prepared at the doctorate level while ADN and diploma programs typically require full-time faculty to have a minimum of a master's degree.




Figure 4-3. Highest education of registered nurses who work as faculty in their principal position, by type of program in which they work*

*Only registered nurses who provided education preparation information are included in the calculations used for this figure. Source: 2008 National Sample Survey of Registered Nurses

4.5 Faculty and Hours of Work

Most RNs who worked as faculty in their principal position were employed for at least 9 months a year (96.1 percent), whereas only 3.9 percent were employed as faculty for 8 or fewer months per year (Table 4-3). Most faculty RNs reported that they work at least 32 hours per week in their principal faculty job. About one-quarter worked fewer than 32 hours per week, indicating the importance of part-time faculty work.

| | Number | Percent |
|------------------|--------|---------|
| Months per year | | |
| 1 to 8 months | 1,223 | 3.9 |
| 9 to 10 months | 18,773 | 60.4 |
| 11 to 12 months | 11,061 | 35.6 |
| Hours per week | | |
| 1-15 hours | 1,812 | 5.8 |
| 16-23 hours | 2,404 | 7.8 |
| 24-31 hours | 3,220 | 10.4 |
| 32-39 hours | 1,975 | 6.4 |
| 40-47 hours | 11,473 | 37.0 |
| 48-55 hours | 7,330 | 23.6 |
| 56 or more hours | 2,842 | 9.2 |

Table 4-3.Months worked per year and hours worked per week, by registered nurses who are
faculty in their principal nursing position

Source: 2008 National Sample Survey of Registered Nurses

4.6 Faculty Earnings and Satisfaction

Nursing faculty received somewhat lower annual earnings than RNs in general. Overall earnings from a principal nursing position by RNs who work full-time averaged \$66,973 in 2008 (Table 3-28), while earnings for faculty RNs were \$63,985 on average (Table 4-4). This may be a reflection of the fact that two-thirds of faculty work fewer than 10 months. Faculty earnings were even lower when compared with RNs who have earned a graduate degree. RNs with master's or doctorate degrees earned an average of \$87,363 per year from their principal full-time nursing position (Table 3-28). Faculty who work in programs that prepare nurses at the bachelor's and higher level earned slightly more on average than those who work in ADN programs. Faculty with a job title of instructor or lecturer earned much less than professors (\$54,944 v. \$69,691).

Registered nurses who work as faculty in their principal position were generally quite satisfied with their work (Figure 4-4). Nearly 90 percent of faculty reported that they are moderately or extremely satisfied with their position, as compared with 81.1 percent of all working RNs (Figure 3-18). The satisfaction of faculty is similar to that of nurses with graduate degrees, of whom 87.3 percent were moderately or extremely satisfied (Figure 3-22).

• Faculty that prepare RNs in bachelor's and higher degree programs were somewhat less satisfied than are those who prepare RNs in ADN programs.

• Faculty with instructor or lecturer job titles were less satisfied than those who were professors.


Table 4-4.Average annual earnings of registered nurses who work full-time as faculty in their
principal nursing position*

| | Annual earnings |
|---|-----------------|
| All faculty | 63,985 |
| Earnings by type of program | |
| Faculty in Diploma/Associate degree program | 62,689 |
| Faculty in Bachelor's degree program | 64,789 |
| Earnings by faculty job title | |
| Instructor/Lecturer | 54,944 |
| Professor | 69,691 |

*Only registered nurses who provided earnings information are included in the calculations used for this table.

Source: 2008 National Sample Survey of Registered Nurses

Figure 4-4. Satisfaction of registered nurses who work as faculty in their principal nursing position





This page deliberately blank.

5.1 Introduction

For more than four decades, Advanced Practice Registered Nurses (APRNs) have been an integral part of the health care system and have served in a variety of roles in acute and ambulatory care settings. APRNs are RNs who have received advanced education to develop knowledge and skills in areas not usual for RNs such as diagnosing and managing common acute and chronic diseases, ordering diagnostic tests, prescribing medications, and performing minor procedures. Commonly recognized APRN designations are nurse practitioner (NP), nurse midwife (NM), nurse anesthetist (NA), and clinical nurse specialist (CNS). While most APRNs are prepared in only one advanced practice area, some may be prepared in more than one of these fields. For this report, RNs were categorized as APRNs if they reported that they were prepared as an APRN in one or more of the recognized specialties.

APRNs are required to complete an approved educational program before taking an exam to obtain certification from a nationally recognized professional organization. Each State establishes criteria for State licensure, certification, or other approval of APRNs, which may include requiring graduate education and/or certification from a professional organization.

In 2008, an estimated 250,527 RNs reported that they were prepared as an APRN in one or more advanced specialties or fields. Of these, 220,494 were employed in nursing (Figure 5-1 and Appendix A, Table 44). The largest group of APRNs is NPs, and their estimated numbers rose between 2004 and 2008. CNSs are the second largest group, but their numbers declined between 2004 and 2008, the only group of APRNs to experience a decline. More information about each type of APRN is provided in dedicated sections in this chapter.

5.2 Nurse Practitioners

NP preparation has its roots in public health nursing and school nursing roles in the early part of the 20th century. In the late 1950s, RNs with clinical experience began to collaborate with physicians in the delivery of primary care, particularly in rural areas and in clinics affiliated with medical centers.




Figure 5-1. Registered nurses prepared for advanced practice, 2004-2008*

*The totals in each bar may not equal the estimated numbers for RNs in each Survey year due to incomplete information provided by respondents and the effect of rounding. Only those who claimed advanced preparation as a nurse midwife, nurse anesthetist, clinical nurse specialist, and/or nurse practitioner are included in the calculations used for this figure.

Source: 2004-2008 National Sample Survey of Registered Nurses

In 1965, the first formal education program for NPs was established at the University of Colorado, with a curriculum focused on health promotion, disease prevention, and children's health. Historically, NP programs could confer a certificate or a degree; now, nearly all States require that new NPs complete a graduate degree in nursing, although most permit previously licensed or certified NPs to continue to practice without a master's degree.¹ Registered nurses with a master's degree in nursing, but who have not completed a master's in an NP program, can enroll in a graduate program to obtain a post-master's certificate allowing them to become NPs.² National certifying Boards require a minimum of a master's degree for new graduates of NP programs. Since

¹ Historical perspective is from Julie A. Fairman. Making Room in the Clinic: Nurse Practitioners and the Evolution of Modern Health Care. New Brunswick: Rutgers University Press, 2008.

² These courses typically include pathophysiology, advanced pharmacology, advanced health assessment and physical exam techniques, and course work with clinical NP patient care practice.

1999, some nurse education programs have offered the Doctor of Nursing Practice (DNP), in addition to or instead of a master's degree, although the number of NPs educated in these programs is small thus far.

Early practice for NPs focused on work particularly with women and children in rural and underserved inner city areas where physicians were scarce. NPs are now prepared and certified to work across populations in any geographic region with focus specialty areas including family care, pediatrics, geriatrics, adult health, women's health, psychiatry, neonatology, and acute hospital care of adults and children (Appendix A, Table 46). To work with the title of an NP, nurses must be certified or recognized by the State in which they practice, which usually requires certification by a national professional organization. The State recognized Scope of Practice for NPs differs depending on State regulations. While regulation of NPs may differ by State, the education, knowledge, and practices of NPs are generally the same nationwide.

To assure that NP data were accurate, care was taken to include only NPs with preparation as nurse practitioners. In this Survey, registered nurses were counted as NPs if they responded as an NP and also provided at least one additional piece of corroborating information indicating either that (1) their primary specialty was a specialty nationally recognized for NPs, (2) they had national or State certification or recognition, (3) they had a master's or higher degree in nursing, or (4) they held a job title of nurse practitioner. This section focuses on all RNs prepared as NPs, and all data refer to this group unless otherwise specified. The analysis in this chapter includes all RNs who have been prepared as NPs, even if they are not currently licensed or certified as such or do not practice in the field. These NPs may be retired or pursuing other occupations within or outside nursing. Understanding the employment patters of RNs, including APRNs, is a central goal of the NSSRN.

5.2.1 Number of Nurse Practitioners

In 2008, an estimated 158,348 nurses had NP preparation (Figure 5-2 and Appendix A, Table 45). Of these, 16,370 were also prepared as CNSs, accounting for 10.3 percent of NPs. Another 2.2 percent of NPs in 2008 also had preparation as NMs, NAs, or in more than two APRN fields.




Figure 5-2. Registered nurses prepared in advanced practice nursing

Source: 2008 National Sample Survey of Registered Nurses

Almost 50 percent of NPs were age 50 or older while 35 percent of NPs were under 45 years old (Figure 5-3). NPs younger than 35 years of age comprise almost 14 percent of the population.




Figure 5-3. Age distribution of registered nurses prepared as nurse practitioners

Source: 2008 National Sample Survey of Registered Nurses

5.2.2 Nurse Practitioners and Education

More than 88 percent of NPs have a graduate degree in nursing or a nursing-related field, with 84.3 percent reporting a master's degree and 3.9 percent holding a doctorate degree (Figure 5-4). The 11 percent of NPs without graduate degrees are mostly those who received their NP education before the majority of States required graduate degrees for NP recognition. The majority (73.4 percent) of NPs reported completing an educational program 13 to 36 months long, while 16.2 percent reported a program length of 8-12 months.

More than 93 percent of NPs employed in nursing report having at least either national certification or State Board recognition. Of those with national certification, 71.6 percent said that national

Advanced Practice Registered Nurses

5

certification is required for their job. Of those with State Board recognition, 82.8 percent said that State Board recognition is required (Table 5-1 and Appendix A, Table 45).




Figure 5-4. Nurse practitioner preparation, by highest nursing or nursing-related degree*

*Only those who provided nursing education preparation information are included in the calculations used for this figure.

Source: 2008 National Sample Survey of Registered Nurses

Table 5-1.Certification and State Board recognition of nurse practitioners employed in nursing,
and whether certification or recognition is required for their job

| | Percent of NPs certified/ recognized | Percent of those certified or recognized for whom it is required for their job |
|---|--|--|
| National organization certification | 83.8 | 71.6 |
| State Board recognition | 88.1 | 82.8 |
| Either national organization certification or State Board recognition | 93.2 | 82.2 |

5.2.3 Nurse Practitioners and Employment

Most NPs are employed in nursing, with 141,286 employed in 2008 (89.2 percent) (Appendix A, Table 45). The vast majority worked at least 10 months per year in their principal position, with only 4.5 percent working 9 or fewer months. More than 63 percent of NPs reported that they work at least 40 hours per week in their principal nursing position; more than 25 percent work 48 hours or more per week (Figure 5-5).






Figure 5-5. Hours worked per week by employed nurse practitioners in their principal nursing position



An estimated 69.6 percent of NPs had a principal position job title of nurse practitioner and the vast majority hold national certification. Of the estimated 97,876 NPs who report nurse practitioner as their job title, 99.6 percent have national certification or State recognition (Appendix A, Table 45).

There were 11.4 percent of nurses prepared as NPs who reported that their job title was staff nurse or equivalent (Figure 5-6). Another 7.4 percent had a title in the area of instruction, and 6.3 percent had a management or administration title.





*Excludes NPs for whom job title was not known.

Source: 2008 National Sample Survey of Registered Nurses

NP education has historically focused on the provision of primary care but NPs may also work in acute care or chronic care settings. About 36 percent of NPs reported primary care as their clinical specialty in their principal position (Table 5-2). General medical-surgical and gynecology/women's health were each the primary clinical areas of 12.1 percent of NPs. Another 8.6 percent specialized in cardiac care, and 6.9 percent focused on chronic care.

| Clinical specialty* | Percent |
|---------------------------------|---------|
| Critical care | 5.8 |
| Cardiac care | 8.6 |
| Chronic care | 6.9 |
| Emergency/trauma care | 3.5 |
| Gastrointestinal | 3.4 |
| General medical surgical | 12.1 |
| Gynecology (women's health) | 12.1 |
| Infectious/communicable disease | 3.7 |
| Neurological | 3.2 |
| Obstetrics/labor & delivery | 8.6 |
| Oncology | 4.4 |
| Primary care | 36.1 |
| Psychiatric/mental health | 6.3 |
| Other specialty** | 16.7 |

Table 5-2.Primary clinical specialties of employed nurse practitioners with job title of nurse
practitioner

*The number of clinical specialties will exceed the total number for all clinical specialties because each NP may report more than one specialty.

**Other clinical specialty includes dermatology, endocrinology, hospice, occupational health, ophthalmology, orthopedic, otolaryngology, plastic/cosmetic surgery, radiology, pulmonary/respiratory, renal/dialysis, urology, no specific area, and other specialties.

Source: 2008 National Sample Survey of Registered Nurses

More than 64 percent of NPs whose job title is nurse practitioner reported that they principally deliver ambulatory care or primary care (Table 5-3). Among the other most commonly reported job titles of NPs, between 14 and 24 percent predominately provide ambulatory or primary care as well.

| Table 5-3. | Employed nurse practitioners whose primary level of care is ambulatory or primary |
|------------|---|
| | care, by job title* |

| | Number | Percent |
|--------------------|--------|---------|
| Staff nurse | 3,029 | 19.0 |
| Management | 1,254 | 14.6 |
| Instruction | 1,857 | 18.3 |
| Nurse practitioner | 61,513 | 64.5 |
| Other** | 1,760 | 24.0 |

*Excludes NPs for whom job title was not known.

**Other job titles include NA, CNS, NM, patient educator, patient coordinator, quality improvement/utilization review, informatics, consultant, legal nurse, researcher, surveyor/auditor/regulator, and no job title.

Source: 2008 National Sample Survey of Registered Nurses

NP preparation historically focused on ambulatory care, and thus NPs often work in primary care settings; however, NPs may also work in specialty clinics, acute care hospitals, schools, public health departments, and extended care facilities. More than 38 percent of NPs were employed in hospital

settings, which can include primary care clinics located in or owned by a hospital (Figure 5-7). Another 35.3 percent worked in ambulatory care settings outside hospitals.

NPs who were employed in ambulatory care settings reported that ambulatory care was the level of care they provide most often (Figure 5-8). However, among NPs who worked in hospital settings, only 33.7 percent listed ambulatory care as their primary level of care. Higher percentages of NPs employed in school health (59.5 percent) and public/community health (47.4 percent) reported that their main level of care was ambulatory or primary care. Only 7.7 percent of NPs work in academic education, and of these NPs, 15.1 percent reported ambulatory or primary care as their main level of care.

5.2.4 Nurse Practitioner Earnings and Job Satisfaction

Registered nurses working both full- and part-time and prepared as NPs earned an average of \$73,776 per year in their principal position. Those who worked full-time earned an average of \$83,192. NPs were, on average, very satisfied with their work (Figure 5-9). More than 87 percent were moderately or extremely satisfied with their principal position, and only 7.6 percent were moderately or extremely dissatisfied.




Figure 5-7. Employment settings in which nurse practitioners work*

 $\ensuremath{^{\star}\text{Excludes}}$ NPs for whom employment setting was not known.

**Public/community health includes occupational health.

***Other includes insurance/benefits/utilization review.




Figure 5-8. Percent of nurse practitioners whose primary level of care is ambulatory or primary care, by employment setting*

*Excludes NPs for whom level of care or employment setting was not known.

**Public/community health includes occupational health.

***Other includes insurance/benefits/utilization review.




Figure 5-9. Nurse practitioner satisfaction with principal position

5.3 Nurse Midwives

Midwifery has a long history. Nurse midwifery arose both in New York City and in Kentucky in the late 19th and early 20th centuries. The earliest U.S. nurse midwifery programs were designed to meet the needs of special populations in urban, rural, and impoverished populations. In 1925, nurse Mary Breckinridge established the Frontier Nursing Service in Kentucky. The educational program of the Frontier Nursing Service drew from public health nursing and British nurse midwifery traditions, focusing primarily on the care of women from preconception through postpartum care, as well as the care of infants. After the American College of Nurse Midwives formed in 1955, practice and education opportunities for nurse midwives expanded. National certification for nurse midwives (NMs) was established in 1970.^{3,4,5}

Source: 2008 National Sample Survey of Registered Nurses

³ Dye, N.S. (1983). Mary Breckinridge, the Frontier Nursing Service and the Introduction of Nurse-Midwifery in the U.S. Bulletin of the History of Medicine, 57(4), pp. 485-507.

⁴ Tom, S. A. C.N.M., M.S. (1982). The evolution of nurse-midwifery: 1900-1960. Journal of Nurse-Midwifery, (27)4, pp. 4-13.

⁵ Dawley, K. C.N.M., PhD. (2003). Origins of nurse-midwifery in the Unites States and its expansion in the 1940s. Journal of Midwifery & Women's Health, (48)2, pp. 86-95.

Nurse midwifery education requirements vary across States. Few nondegree programs exist in the United States and most programs, including the Frontier Nursing Service program, confer a master's or doctorate degree. To practice as a nurse midwife, NMs must be certified or recognized by the State in which they practice or by a national professional organization. Much like NPs, the Scope of Practice of NMs varies by State legislation. Many States require a graduate degree in nursing, but some still permit NMs to practice after completing a bachelor's degree and a post-licensure certification program.

To assure that NM data were accurate, care was taken to include only NMs with preparation as nurse midwives. In this Survey, nurses who responded as having been prepared as an NM were only considered as such if they also provided at least one additional piece of corroborating information indicating either that (1) their primary APRN area of study was family care, maternal-child health, neonatal care, nurse-midwifery, obstetrics/gynecology, or women's health; (2) they reported national or State certification or recognition; (3) they had a master's or higher degree in nursing; or (4) their job title was reported as nurse midwife.

5.3.1 Number of Nurse Midwives

NMs comprised the smallest number of the four APRN groups in 2008, with an estimated 18,492 registered nurses prepared as NMs (Appendix A, Table 45). Of these, an estimated 3,164 also have been prepared in another APRN field.

NMs were older than nurses prepared as NPs. More than 54 percent were 50 years or older (Figure 5-10).




Figure 5-10. Age distribution of registered nurses prepared as nurse midwives

Source: 2008 National Sample Survey of Registered Nurses

5.3.2 Nurse Midwives and Education

Nurse midwives were the least likely of the APRN groups to have been educated at the graduate level, with only 55.5 percent reporting that they had a master's or doctorate degree (Figure 5-11). More than 30 percent of NMs reported a diploma or an ADN as their highest nursing education. Nurse midwives were more likely than other APRNs to have been educated in shorter programs, with 56 percent having completed programs that lasted 13 to 36 months, and 34.5 percent completing 8- to 12-month programs. This is consistent with the fact that earlier NM programs did

not require students to have a bachelor's degree nor did the early programs confer a master's degree.⁶




Figure 5-11. Nurse midwife preparation, by highest nursing or nursing-related degree*

*Only NMs who provided nursing education preparation information are included in the calculations used for this figure. Source: 2008 National Sample Survey of Registered Nurses

More than 91 percent of NMs employed in nursing had national certification or recognition from their State Board of Nursing. National certification is more common than State Board recognition (Table 5-4). Of those NMs who are employed in any nursing position and have national certification, 57.9 percent report that their certification is required for their job. Of those employed with State Board recognition, 71.9 percent said their recognition was required.

⁶ In some states, individuals can be certified or recognized as non-nurse midwives. The requirements for this vary by State and are generally considered separate from the Nurse Midwife credential.



| | Percent of NMs certified/ recognized | Percent of those certified or recognized for whom it is required for their job |
|---|---|--|
| National organization certification | 89.7 | 57.9 |
| State Board recognition | 70.0 | 71.9 |
| Either national organization certification or State Board recognition | 91.5 | 60.4 |


Table 5-4.Certification and State Board recognition of nurse midwives employed in nursing,
and whether certification or recognition is required for their job

Source: 2008 National Sample Survey of Registered Nurses

5.3.3 Nurse Midwives and Employment

In 2008, 15,581 NMs were employed in nursing positions (84.3 percent of all NMs) (Appendix A, Table 45) and only 35.6 percent of NMs worked less than 40 hours per week. More than one quarter of NMs work more than 48 hours per week (Figure 5-12).

Figure 5-12. Hours worked per week by employed nurse midwives in their principal nursing position








Figure 5-13. Job titles of nurse midwives*

*Excludes NMs for whom job title was not known. Source: 2008 National Sample Survey of Registered Nurses

An estimated 41.7 percent of employed NMs reported their principal position's job title as nurse midwife (Figure 5-13). All of these NMs also had State recognition or national certification in this field (Appendix A, Table 45). Many NMs have a job title of staff nurse (37.5 percent), but 20.7 percent have other titles such as patient coordinator, faculty, or instructor.

The majority of NMs (57.9 percent) worked in hospital settings in 2008, and 25.1 percent worked in ambulatory care (Figure 5-14). NMs often work in clinical specialties related to midwifery, even when their job title is not nurse midwife. Among those who worked in a hospital setting, 41.8 percent specialize in labor and delivery, 33.7 percent in obstetrics, and 13.6 percent in gynecology or women's health (Table 5-5).





*Excludes NMs for whom setting was not known.

Source: 2008 National Sample Survey of Registered Nurses

Table 5-5.Clinical specialties of registered nurses prepared as nurse midwives, for selected
job titles in the hospital setting

| Clinical specialty* | Percent with nurse midwife title | Percent of nurse midwives in hospital setting |
|---------------------------|-------------------------------------|--|
| Labor and delivery | 56.0 | 41.8 |
| Obstetrics | 68.9 | 33.7 |
| Gynecology/women's health | 50.2 | 13.6 |

*NMs could report more than one clinical specialty.

Source: 2008 National Sample Survey of Registered Nurses

5.3.4 Nurse Midwife Earnings and Job Satisfaction

Registered nurses prepared as NMs and working full- or part-time earned an average of \$69,222 per year in their principal nursing position, and those who were employed full-time earned an average of

Advanced Practice Registered Nurses

5

\$75,340 and reported high levels of satisfaction with their principal nursing position (86.2 percent being moderately or extremely satisfied) (Figure 5-15).




Figure 5-15. Nurse midwife satisfaction with principal position

Source: 2008 National Sample Survey of Registered Nurses

5.4 Nurse Anesthetists

Registered nurses have been providing anesthesia care in the United States since the Civil War. The first school of nurse anesthesia, which was established in 1909 at Saint Vincent Hospital in Oregon, consisted of a 6-month course. Most nurse anesthesia programs that opened after that program conferred an anesthesia bachelor's degree, diploma, or certificate. In 1982, the American Association of Nurse Anesthetists recommended that NA education be a minimum of a master's degree and by 1998, all NA programs were at the master's degree level. Previously licensed NAs are allowed to

continue their practice. Nurse anesthetists take a national certification exam at the end of their NA education and are licensed in most States through State nursing boards.^{7,8}

5.4.1 Number of Nurse Anesthetists

Nurse anesthetists were the third largest group of APRNs, next to NPs and CNSs, with an estimated 34,821 nurses prepared as anesthetists in 2008 (Appendix A, Table 45). There was a 7.1 percent growth in the number of NAs between 2004 and 2008, compared with a 9 percent increase between 2000 and 2004.

Nurse anesthetists have a larger proportion of younger nurses than other APRN groups. In 2008, nearly 40 percent were under 45 years old (Figure 5-16). More than 40 percent of NAs are men (Figure 7-2) (discussed in more detail in Chapter 7).

⁷ Bankert, M. Watchful Care: A History of America's Nurse Anesthetists. New York: Continuum. 1989.

⁸ Thatcher, V.S. History of Anesthesia with Emphasis on the Nurse Specialist. Philadelphia: JB Lippincott Company. 1953.





Figure 5-16. Age distribution of registered nurses prepared as nurse anesthetists


Source: 2008 National Sample Survey of Registered Nurses

5.4.2 Nurse Anesthetists and Education

Currently, a minimum of a master's degree is required prior to NA national certification. Because this requirement is relatively new and there is a long history of NA education at less than the graduate level, in 2008 only 65.4 percent of NAs held a master's or doctorate degree as their highest nursing or nursing-related degree (Figure 5-17). Nearly all NAs (96.4 percent) were educated in programs 13 to 36 months long.

Virtually all NAs had national certification (90.9 percent), and most had State recognition (82.1 percent) (Table 5-6). Nearly 90 percent of NAs report that national certification is required for their job.





*Only NAs who provided nursing education preparation information are included in the calculations used for this figure.


Source: 2008 National Sample Survey of Registered Nurses


Table 5-6.Certification and State Board recognition of nurse anesthetists employed in nursing,
and whether certification or recognition is required for their job

| | Percent of all NAs certified/ recognized | Percent of those certified or recognized for whom it is required for their job |
|---|--|--|
| National organization certification | 90.9 | 89.6 |
| State Board recognition | 82.1 | 89.5 |
| Either national organization certification or State Board recognition | 99.3 | 91.8 |

5.4.3 Nurse Anesthetists and Employment

In 2008, 31,868 NAs were employed in nursing positions (91.5 percent of all NAs) (Appendix A, Table 45). Nurse anesthetist worked more hours per week, on average, than other APRN groups, with 76.5 percent working 40 or more hours per week (Figure 5-18). Similarly to NMs, more than one quarter of NAs work more than 48 hours per week.






Source: 2008 National Sample Survey of Registered Nurses

An estimated 92.3 percent of NAs reported their job title in their principal nursing position as nurse anesthetist. All of these NAs also reported that they have national certification and/or recognition by their State Board of Nursing, even if it is not required for their current position (Appendix A, Table 45).

The majority of NAs worked in hospital settings (82.2 percent) in 2008, and another 14.8 percent worked in ambulatory care.

5.4.4 Nurse Anesthetist Earnings and Job Satisfaction

Registered nurses prepared as NAs earned an average of \$135,776 per year in their principal nursing position, making them the highest paid group of APRNs, and the highest paid of all RNs. They were the most satisfied of the APRN groups, with 93.5 percent being moderately or extremely satisfied with their principal position (Figure 5-19).





Source: 2008 National Sample Survey of Registered Nurses

5.5 Clinical Nurse Specialists

The CNS role was conceived in 1954 and developed in acute care settings and mental health hospital settings. The first clinical specialist certification exam was launched in that year in psychiatric/mental health nursing. From the 1960s through the 1990s, clinical nurse specialist master's education programs were developed in many schools of nursing. The CNS education and role focuses on improving clinical care, primarily in hospitals and extended care facilities by having an expert nurse (the CNS) who could help coordinate care for individuals, educate nursing personnel who provided direct care, and help identify and improve aspects of the health system organization that affected patients and nursing staff. CNSs have expertise in one or more clinical areas such as oncology, pediatrics, geriatrics, psychiatric/mental health, adult health, obstetrics, acute/critical care, and community health.⁹

CNSs have been required to have a master's degree in nursing since the inception of this advanced practice field, and while national certification is available for many CNS specialties, certification is not required by State Boards of Nursing nor are most CNSs recognized in many States' licensure regulations as APRNs. In some States, the job title of clinical nurse specialist is not legally limited to nurses who have CNS preparation or certification. Thus, some RNs who reported that their job title is clinical nurse specialist do not have a graduate degree and were not counted as a prepared CNS in this Survey.

5.5.1 Number of Clinical Nurse Specialists

In 2008, there were an estimated 59,242 nurses prepared as CNSs, some of whom also were prepared in other advanced specialties (Appendix A, Table 45). CNSs are the second largest group of APRNs. There was a marked decline of 18.3 percent in the number of CNSs between 2004 and 2008. However, over the same period there was an increase of 11.4 percent in the number of registered nurses prepared as both NPs and CNSs.

CNSs were older than other APRN groups in 2008, with 63.6 percent over 50 years old and only 18.2 percent under 45 years (Figure 5-20). Few CNSs, only 10.2 percent, are under the age of 40.

⁹ Montemuro, M.A. MHSc, RN. (1987). The Evolution of the Clinical Nurse Specialist: Response to the Challenge of Professional Nursing Practice. *Clinical Nurse Specialist, (1)*3, pp. 106-110.

5.5.2 Clinical Nurse Specialists and Education

CNS education focuses on how clinical care is affected by the patient's personal and family characteristics and environment, the nursing personnel who provide care, and the health system organization. CNSs have been required to have a master's degree in nursing since the inception of this advanced practice field. The vast majority of CNSs (92.8 percent) reported a master's degree as their highest preparation (Figure 5-21). An additional 7.2 percent completed a doctorate degree for their CNS preparation. Most CNSs reported that their education program lasted 13 to 36 months (79.3 percent).

Many registered nurses prepared as CNSs do not have national certification or recognition from their State Board of Nursing; only 54.6 percent have recognition or certification from one or both. Less than 40 percent have national certification, and 41.6 percent have State Board recognition (Table 5-7). Of those with national certification, 39.5 percent said their certification was required for their job, while 47.2 percent of those with State Board recognition reported that their certification was required for their principal position.




Figure 5-20. Age distribution of registered nurses prepared as clinical nurse specialists




Figure 5-21. Clinical nurse specialist preparation, by highest nursing or nursing-related degree*

*Only CNSs who provided nursing education preparation information are included in the calculations used for this figure.

Source: 2008 National Sample Survey of Registered Nurses

Table 5-7.Certification and State Board recognition of clinical nurse specialists employed in
nursing, and whether certification or recognition is required for their job

| | Percent of CNSs certified/ recognized | Percent of those certified or recognized for whom it is required for their job |
|---|---|--|
| National organization certification | 39.7 | 39.5 |
| State Board recognition | 41.6 | 47.2 |
| Either national organization certification or State Board recognition | 54.6 | 41.4 |

5.5.3 Clinical Nurse Specialists and Employment

CNSs had the lowest rate of employment in nursing of all the advanced specialties, with 84 percent working in nursing in 2008 (Appendix A, Table 45). However, nearly 70 percent of CNSs employed in nursing worked at least 40 hours per week, which is a larger percentage than either NPs or NMs (Figure 5-22). Only NAs had as high a percentage working more than a 40 hour week.




Figure 5-22. Hours worked per week by employed clinical nurse specialists in their principal nursing position

Source: 2008 National Sample Survey of Registered Nurses

An estimated 18.8 percent of CNSs reported their job title as clinical nurse specialist (Figure 5-23). State Board recognition or national certification was held by 85.4 percent of nurses with the CNS job title (Appendix A, Table 45). Other common job titles among nurses prepared as CNSs are instruction (21.1 percent), and management or administration (17.8 percent). An additional 16 percent reported a staff nurse job title, and 10.5 percent had other titles such as public health nurse, school nurse, or patient coordinator (Figure 5-23).





Figure 5-23. Job titles of employed nurses prepared as clinical nurse specialists*


*Excludes CNSs for whom job title was not known.

**Other job titles include consultant, NA, informatics, patient educator, patient coordinator, researcher, school nurse, and public/community health.

Source: 2008 National Sample Survey of Registered Nurses

Some nurses prepared as CNSs also had preparation in other advanced specialties. Dual CNS-NP preparation was reported by 27.6 percent of CNSs, a trend that has increased since 2004. Among RNs prepared as both NPs and CNSs, the most common job title was nurse practitioner (54.8 percent) (Figure 5-24). The remainder had a variety of job titles, including clinical nurse specialist, professor or instructor, management or administration titles, or a staff nurse title.





*Excludes dual-prepared CNSs and NPs for whom job title was not known.

**Other job titles include CNS, consultant, management, instruction, staff nurse, patient coordinator, and researcher.

Source: 2008 National Sample Survey of Registered Nurses

Nearly half of nurses prepared as CNSs worked in hospitals in 2008 (49.4 percent, Figure 5-25). Another 15.7 percent were in academic settings, and 13.3 percent worked in ambulatory care.

5.5.4 Clinical Nurse Specialist Earnings and Job Satisfaction

Registered nurses prepared as CNSs working full- or part-time earned an average of \$74,918 per year in their principal nursing position, and those who were employed full-time earned an average of \$83,787. Those who were prepared in both NP and CNS specialties earned an average of \$76,350 overall and \$86,330 for full-time positions. CNSs were very satisfied with their principal positions (Figure 5-26), with 91 percent being moderately or extremely satisfied.




Figure 5-25. Employment settings in which clinical nurse specialists work*

 $\ast \mbox{Excludes CNSs}$ for whom employment setting was not known.

**Public/community health includes school and occupational health.




Figure 5-26. Clinical nurse specialist satisfaction with principal position

Source: 2008 National Sample Survey of Registered Nurses

Nurses Who Do Not Work in Nursing 6

6.1 Introduction

This chapter examines the characteristics of RNs who do not work in nursing and their plans regarding future nursing work. An estimated 466,564 licensed RNs (just over 15 percent of all licensed RNs) were not employed in nursing in 2008. Nurses choose to not work in nursing for a variety of reasons. Some are retired but may choose to retain a nursing license for a variety of reasons including potential future employment or because they feel personally connected to the profession and want to retain the link of licensure. Others stop working to attend to family needs, because of illness, to receive further education, or for other personal reasons. Still others are not working in nursing, either temporarily or permanently, because they are unhappy with some aspect of the profession.

6.2 Demographics of Nurses Not Working in Nursing

6.2.1 Nurses Not Working in Nursing and Age

Nearly one-third of RNs who were not working in nursing in 2008 were 65 years of age or older; almost half were 60 years of age or older (Figure 6-1). This suggests that the majority of nurses who have retained licenses but are no longer working may have left the workforce for age-related reasons.

A disproportionate percentage of licensed RNs aged 50 to 54 have not been working in nursing for 11 to 15 years (17.6 percent of RNs out of work this long) or for 16 or more years (24.3 percent) (Appendix A, Table 48). These individuals left nursing by 1997 or earlier, when they were in their 20s and 30s. During the mid-1990s, RN employment rates were somewhat lower, and the percentage of nurses employed in hospitals declined. These data suggest that nurses who were under age 40 at that time, left the profession and never returned to nursing work.




Figure 6-1. Age distribution of nurses who are not working in nursing*

*Includes only registered nurses who are not working in nursing.

Source: 2008 National Sample Survey of Registered Nurses

6.2.2 Presence of Children

The decision to work in nursing is more closely tied to age than to whether an RN has children. The percent of RNs not working in nursing in 2008 increases with age group (Figure 6-2 and Appendix A, Table 49). Nurses whose children were all under age 6 were less likely to work in nursing than were nurses whose children were all at least 6 years old. Moreover, older mothers were less likely to work in nursing than younger mothers.




Figure 6-2. Percentage of nurses not working in nursing, by age and presence of children*

*Includes only registered nurses who are not working in nursing.


Source: 2008 National Sample Survey of Registered Nurses

6.3 Length of Time Away From Nursing Work

Many nurses who left employment in nursing did so for a short period of time (Figure 6-3 and Appendix A, Table 51). About half of RNs who were not working in nursing in 2008 had been out of nursing work for 4 or fewer years.

- Among RNs who were under 50 years old, 13.3 percent had been out of nursing for less than 1 year, and another 44.2 percent stopped working 1 to 4 years prior.
- Only 21.2 percent of RNs under 50 years old stopped working in nursing 10 or more years ago, compared with 31.4 percent of nurses 50 years and older.

Figure 6-3. Length of time since nonworking nurses last worked in nursing, by age*



*Includes only registered nurses who are not working in nursing.

Source: 2008 National Sample Survey of Registered Nurses

6.4 Reasons for Not Working in Nursing

Nurses were asked their reasons for not working in the nursing profession. They were able to indicate more than one reason. Significantly, reported reasons for not working in nursing were associated with RNs' age. Among nurses 50 years and older, the main reason for not working in nursing was retirement (54.4 percent), although personal and family reasons (41.5 percent),

workplace reasons (35.5 percent), and personal career factors (32.4 percent) also were important. Among nurses under 50 years old, the dominant reason for not working is associated with personal family factors (68.5 percent), such as the need to care for children (Figure 6-4). Another 39.8 percent were not working for personal career reasons.

Of concern, however, is the fact that 41 percent of RNs under 50 years and 35.5 percent of RNs 50 years of age and older reported that they were not working in nursing due to workplace problems such as burnout, stressful work environment, and poor management.




Figure 6-4. Reasons nurses are not working in nursing, by age group*

*Percents do not add to 100 because registered nurses may have provided more than one reason. Includes only registered nurses who are not working in nursing.

6.5 Plans Regarding Nursing Work

Nurses who were not working in nursing in March 2008 were asked about their plans regarding nursing employment. Their answers were associated with both age and marital status (Table 6-1).

- Nurses under 50 years old who were never married were more likely than their married counterparts to have already returned to work in nursing when they completed the Survey.
- Nurses under 50 years old who were married or in a domestic partnership were more likely than their divorced or never married counterparts to have plans to return to nursing, with only 14.2 percent saying they did not plan to return to nursing work. However, a large percentage of these RNs were undecided about their future plans regarding nursing work (26.4 percent).
- Nurses 50 years and older were less likely than RNs under 50 to plan to return to nursing work. This was particularly true for nurses who were never married (54.7 percent).

| | Under 50 years old (percent) | | 50 years and older (percent) | | | |
|--|---------------------------------|----------------------|---------------------------------|---------|----------------------|------------------|
| Intentions | Married | Divorced/ widowed | Never married | Married | Divorced/ widowed | Never married |
| Have returned to nursing Actively looking for work in | 17.2 | 32.4 | 36.3 | 6.1 | 8.0 | 9.4 |
| nursing | 7.5 | 16.5 | 8.1 | 2.9 | 4.4 | 3.9 |
| Plan to return in the future | 34.7 | 17.8 | 17.4 | 9.2 | 11.0 | 6.3 |
| No future intentions to return | 14.2 | 15.1 | 15.2 | 48.2 | 50.5 | 54.7 |
| Undecided | 26.4 | 18.2 | 23.0 | 33.6 | 26.1 | 25.6 |
| Total | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 |

Table 6-1. Intentions of nurses who are not working in nursing, overall, and by marital status*

*Includes only registered nurses who are not working in nursing.

Source: 2008 National Sample Survey of Registered Nurses


Only 5.5 percent of RNs who were not working in nursing reported that they were seeking a nursing position at the time they completed the Survey (Appendix A, Table 51). Most of the RNs who were seeking a nursing position had been out of nursing work for less than 1 year.


Pursuing nursing employment is closely associated with age. Among RNs who were under 50 years old, 27.7 percent who had not been working in nursing for less than 1 year were seeking nursing employment (Figure 6-5). About 8 percent of nurses under 50 years old who had not been working

for 1 to 4 years were seeking nursing employment. In comparison, among RNs 50 years and older, 7.7 percent who had not been working for less than 1 year were seeking nursing work, while less than 4 percent who had been out of the RN workforce for 1-4 years were searching for a nursing job.

Among RNs not working in nursing who were seeking nursing employment, only 25.8 percent wanted a full-time nursing position (Appendix A, Table 52). Another 39.3 percent preferred part-time work, and 34.9 percent were seeking any level of nursing employment.

Most RNs who were seeking nursing work in 2008 had been doing so for only a short time (Appendix A, Table 52). While 31.6 percent had been seeking a nursing position for 4-9 months, 37.8 percent had been looking for a nursing job for up to 1 month.





*Includes only registered nurses who are not working in nursing.

This page deliberately blank.



Demographics of the Registered Nurse Workforce

7.1 Introduction

This chapter examines the age, gender, race/ethnicity, language competency, and family characteristics of registered nurses, and how these factors are associated with employment. It also presents the geographic distribution of RNs. Nursing employment continues to vary significantly across the country, with a two-fold difference in the employed RN-to-population ratio among the States. Geographic variation occurs in the employment setting, educational level, racial/ethnic minority representation, and median earnings of employed RNs. While still a workforce of primarily White, non-Hispanic women, the population of RNs in the United States continues to become more diverse with an increase of men and racial/ethnic minority groups among new entrants to the workforce.

7.2 Registered Nurses and Age

In 2008, for the first time in the past three decades, the rate of aging of the nursing workforce slowed (Figure 3-3). In 1988, half of the working population of RNs was less than 38 years of age (Table 7-1 and Appendix A, Table 1). By 2004 the median age rose to 46 years. In 2008, however, the median age did not increase but remained steady at 46 years of age.

This plateau in the trend toward aging of the RN workforce was the consequence of an increase in employed RNs under 30 years of age—the first increase seen in this age group since the inaugural NSSRN in 1977—which helped to offset the steady rise in the number of employed RNs age 60 and older (Figure 7-1 and Appendix A, Table 1). The increase in the youngest cohort of RNs is the result of rapidly rising enrollments in schools of nursing, in particular in bachelor's nursing programs, whose student population is younger than that of other initial nursing education programs. This rising trend in BSN enrollment started a few years earlier, and graduates from it are now entering the workforce.




| Survey year | Median age |
|-------------|------------|
| 2008 | 46 |
| 2004 | 46 |
| 2000 | 44 |
| 1996 | 42 |
| 1992 | 40 |
| 1988 | 38 |

Table 7-1. Median age of registered nurses

Source: 1988-2008 National Sample Survey of Registered Nurses

Figure 7-1. Percent of registered nurses working in nursing, by age cohort, 1988-2008



Source: 1988-2008 National Sample Survey of Registered Nurses

7.3 Registered Nurses and Gender

Men comprised a growing percentage of RNs in 2008. To provide a comparison over time within the 2008 Survey, male RNs who were licensed before 2000 are compared to those licensed in 2000

or later. Male RNs made up 6.2 percent of employed RNs who were licensed before 2000 and 9.6 percent of those licensed in 2000 or later (Table 7-2). Men licensed as RNs since 2000 are older than women licensed in the same period, with a median age of 35 compared to 31 for women. In contrast, male and female RNs licensed before 2000 have a similar age profile.

Table 7-2. Gender of employed registered nurses, by year of initial registered nurse license

| | Licensed | Licensed before 2000 | | n 2000 or later |
|----------------------------|----------|----------------------|------|-----------------|
| | Male | Female | Male | Female |
| Percent of employed nurses | 6.2 | 93.8 | 9.6 | 90.4 |
| Median age | 49 | 50 | 35 | 31 |

Source: 2008 National Sample Survey of Registered Nurses

Male and female RNs were equally likely to have pursued a bachelor's or higher degree in nursing or nursing-related fields (Table 7-3). However, men were somewhat more likely than women to have received non-nursing higher degrees.

Table 7-3. Highest education of registered nurses working in nursing, by gender

| | Percent with bachelor's and higher in nursing or nursing-related field | Percent with bachelor's and higher in nursing and any non-nursing field |
|-------|---|--|
| Men | 49.9 | 62.2 |
| Women | 50.3 | 55.1 |

Source: 2008 National Sample Survey of Registered Nurses

The employment profile of men who work in nursing is different from that of women. A greater proportion of male RNs work in hospitals compared to women; nearly 76 percent of men report primary employment in a hospital, compared with 61.2 percent of women (Table 7-4).

Differences in the employment of male and female RNs are even more pronounced in certain job titles. Men made up just over 7 percent of employed RNs overall in 2008, yet they constituted 41.1 percent of nurse anesthetists (Figure 7-2).



| | Men (percent) | Women (percent) |
|--------------------|------------------|--------------------|
| Hospital | 75.6 | 61.2 |
| Ambulatory care | 5.6 | 10.9 |
| Home health | 3.9 | 6.6 |
| Nursing home | 4.4 | 5.3 |
| Academic education | 2.3 | 3.9 |
| Other** | 8.0 | 12.0 |
| Total | 100.0 | 100.0 |


Table 7-4. Employment settings, by gender*

*Excludes registered nurses for whom employment setting was not known.

**Other includes public/community health, school health, occupational health, and insurance/benefits/utilization review.

Source: 2008 National Sample Survey of Registered Nurses

Figure 7-2. Percent of men employed in nursing, by job title*





*Excludes registered nurses for whom job title was not known.

**Other includes consultant, CNS, researcher, informatics, surveyor/auditor/regulator, patient educator, and nurses who reported having no job title.

7.4 Registered Nurses and Race and Ethnicity

Racial/ethnic minority groups remain underrepresented in the RN population when compared with their profile in the general population, but gains have been made over time (Figure 7-3). The population of RNs from minority racial/ethnic groups has increased from 119,512 in 1980 to 333,368 in 2000; by 2008, it increased further to 513,860 (Appendix A, Table 1). Nurses from minority racial and ethnic groups still represent only 16.8 percent of all nurses. This percent may change in the future since RNs from racial and ethnic minority groups represent a larger percentage of recently graduated nurses than of nurses who graduated in earlier graduate cohorts (Figure 7-4). American Indian RNs represent a very small percent of RNs working in nursing, which limits the ability to analyze their representation across variables such as employment setting and job title.











Figure 7-4. Racial/ethnic distribution, by year of graduation from initial nursing education

Source: 2008 National Sample Survey of Registered Nurses

The racial and ethnic profile of the RN population is substantially different from that of the U.S. population. As seen in Figure 7-5, 65.6 percent of the U.S. population is White, non-Hispanic, while 83.2 percent of RNs were White, non-Hispanic in 2008. Hispanics, Blacks, and American Indians/Alaska Natives are underrepresented in the RN population. Asians are slightly overrepresented among RNs, with 5.8 percent of RNs reporting a racial background of Asian, Native Hawaiian, or Pacific Islander, compared with 4.5 percent of the U.S. population. As discussed in Chapter 8, a substantial number of RNs who are educated in countries other than the United States have been recruited and received their initial nursing education in the Philippines or India. This may be a contributing factor to the comparatively high share of Asians among RNs.



Source: 2008 National Sample Survey of Registered Nurses

The age profile of the RN workforce differs by racial/ethnic group. White, non-Hispanic RNs are the oldest racial/ethnic category of RNs working in nursing yet they were the youngest group at the time of their initial licensure (Table 7-5). Nearly all other racial/ethnic groups were slightly older at the point of initial licensure but were slightly younger than White, non-Hispanic nurses in the 2008 workforce.

U.S. population

□ RN population



| | Age at licensure | Current age |
|---|------------------|-------------|
| White, non-Hispanic | 25 | 47 |
| Pacific Islander, non-Hispanic | 25 | 39 |
| Hispanic/Latino, any race | 28 | 40 |
| Two or more races, non-Hispanic | 28 | 45 |
| Asian, non-Hispanic | 29 | 41 |
| Black/African American, non-Hispanic | 29 | 44 |
| American Indian/Alaska Native, non-Hispanic | 31 | 45 |
| Total | 25 | 46 |

Table 7-5.Median current age and age at licensure of registered nurses working in nursing, by
race/ethnicity

Source: 2008 National Sample Survey of Registered Nurses

Educational attainment varied among the racial/ethnic groups captured in the 2008 Survey. The most common type of initial RN education among all racial/ethnic groups except Asian RNs is the ADN (Table 7-6). Among Asian RNs, 69.6 percent entered the profession with a bachelor's or higher degree. This high rate of initial BSN education may be in part due to the large share of Asian RNs who were internationally educated. Among internationally educated RNs, bachelor's education for those who emigrate to the United States is more common than among RNs educated in the United States.

| | Diploma (percent) | Associate degree (percent) | Bachelor's and higher (percent) |
|--------------------------------------|----------------------|-------------------------------|---------------------------------------|
| White, non-Hispanic | 19.3 | 48.2 | 32.5 |
| Black/African American, non-Hispanic | 13.2 | 54.7 | 32.1 |
| Hispanic/Latino, any race | 5.5 | 55.1 | 39.4 |
| Asian, non-Hispanic | 12.6 | 17.8 | 69.6 |
| Total | 17.8 | 47.2 | 34.9 |

Table 7-6. Initial nursing education of registered nurses employed in nursing, by race/ethnicity

Source: 2008 National Sample Survey of Registered Nurses

White, non-Hispanic RNs were the least likely racial/ethnic group in 2008 to have attained a bachelor's degree or higher in nursing, nursing-related, or non-nursing fields (Table 7-7). Deconstructing this educational attainment profile reveals that Hispanic and Asian RNs were more likely to have pursued a bachelor's degree for initial RN education but less likely to have pursued graduate degrees than were White, non-Hispanic RNs. In contrast, Black RNs were more likely than White, non-Hispanic nurses to have achieved both bachelor's and graduate degrees.



| | Bachelor's degree (percent) | Master's or doctorate degree (percent) | Total (percent) |
|--------------------------------------|--------------------------------|--|--------------------|
| Nursing and nursing-related | | | |
| White, non-Hispanic | 35.0 | 13.4 | 48.4 |
| Black/African American, non-Hispanic | 37.9 | 14.6 | 52.5 |
| Hispanic/Latino, any race | 41.0 | 10.5 | 51.5 |
| Asian, non-Hispanic | 67.3 | 8.3 | 75.6 |
| Total | 37.3 | 13.0 | 50.3 |
| Nursing and non-nursing | | | |
| White, non-Hispanic | 39.2 | 14.5 | 53.7 |
| Black/African American, non-Hispanic | 41.5 | 16.7 | 58.1 |
| Hispanic/Latino, any race | 45.6 | 11.0 | 56.6 |
| Asian, non-Hispanic | 70.0 | 10.6 | 80.6 |
| Total | 41.4 | 14.3 | 55.6 |

Table 7-7. Highest education of registered nurses employed in nursing, by race/ethnicity

Source: 2008 National Sample Survey of Registered Nurses

Racial/ethnic diversity also varied among APRNs. White, non-Hispanic RNs comprised 83.3 percent of all nurses prepared in advanced practice fields, only slightly higher than the 82.2 percent of all employed RNs who were White, non-Hispanic (Table 7-8). This higher representation by White, non-Hispanic RNs is found across all APRN disciplines except NMs. Black RNs have a higher degree of representation among APRNs because of a higher relative percentage among NMs and NPs. Asian and Hispanic RNs are significantly underrepresented among APRNs.

| Table 7-8. | Advanced practice preparation of registered nurses, by race/ethnicity |
|------------|---|
|------------|---|

| | White, non-Hispanic (percent) | Black/African American, non-Hispanic (percent) | Hispanic/Latino, any race (percent) | Asian, non-Hispanic (percent) |
|------------------------------|-------------------------------------|---|---|-------------------------------------|
| Employed nurses | 82.2 | 5.6 | 3.9 | 5.8 |
| All advanced-practice nurses | 83.3 | 6.3 | 3.5 | 4.2 |
| Nurse practitioners | 84.1 | 5.8 | 3.9 | 3.5 |
| Clinical nurse specialists | 87.9 | - | - | - |
| Nurse midwives | 57.0 | 21.8 | - | _ |
| Nurse anesthetists | 89.7 | - | - | - |

-Too few cases to report estimated number (fewer than 30 respondents).

Source: 2008 National Sample Survey of Registered Nurses

Among the four largest racial/ethnic groups, hospital employment was the most common place of employment (Table 7-9). A smaller percent of Asian, non-Hispanic RNs worked in home health than any other racial/ethnic group. Hispanic/Latino RNs were least represented in nursing homes.

In the aggregate, American Indian RNs in 2008 were less likely to be employed in hospitals or as staff nurses than were all other racial/ethnic groups. Generally, the racial/ethnic pattern of employment varied across all other employment settings.

| Percent of nurses | White, non-Hispanic (percent) | Black/African American, non-Hispanic (percent) | Hispanic/Latino, any race (percent) | Asian, non-Hispanic (percent) | Total (percent) |
|-------------------|-------------------------------------|---|---|-------------------------------------|--------------------|
| Hospital | 60.9 | 63.4 | 68.8 | 75.4 | 62.2 |
| Ambulatory care | 11.3 | 6.7 | 6.9 | 5.3 | 10.5 |
| Home health | 6.8 | 6.4 | 6.8 | 2.7 | 6.4 |
| Nursing home | 5.2 | 6.0 | 3.5 | 6.0 | 5.3 |
| Total | 84.2 | 82.5 | 86.0 | 89.3 | 84.5 |

Table 7-9. Employment setting of registered nurses, by race/ethnicity*

*Excludes registered nurses for whom employment setting was not known.

Source: 2008 National Sample Survey of Registered Nurses

More than 80 percent of RNs working in nursing were employed in three job titles. Among the four largest racial/ethnic groups, White, non-Hispanic nurses were less likely than the other three groups to be employed as staff nurses and more likely to be in patient care coordinator positions (Table 7-10). Black RNs had the highest percentage of employment among the four groups in management and administration positions at 13.8 percent, followed closely by White, non-Hispanic RNs at 12.9 percent. Management jobs were held by 10.9 percent of Hispanic/Latino RNs and 7.2 percent of Asian RNs.

| Table 7-10. | Job titles of registered nurses, by race/ethnicity* |
|-------------|---|
| | |

| Job title | White, non-Hispanic (percent) | Black/African American, non-Hispanic (percent) | Hispanic/Latino, any race (percent) | Asian, non-Hispanic (percent) | Total (percent) |
|---------------------|-------------------------------------|---|---|-------------------------------------|--------------------|
| Staff nurse | 64.8 | 67.1 | 72.0 | 83.0 | 66.3 |
| Management | 12.9 | 13.8 | 10.9 | 7.2 | 12.5 |
| Patient coordinator | 5.6 | 5.2 | 4.1 | 3.5 | 5.4 |
| Total | 83.3 | 86.1 | 87.0 | 93.7 | 84.3 |

*Excludes registered nurses for whom job title was not known.

7.5 Registered Nurses and Language Competency

In communities where English is not the first language of many patients, the dominant factor in cultural competency for RNs may be how well they can communicate with the population they serve. In 2008, most working RNs spoke only English fluently. Just 5.1 percent of RNs spoke Spanish, and 3.6 percent spoke Tagalog or another Filipino language. About 1.1 percent spoke French, while less than 1 percent or RNs spoke any Chinese dialect, German, American Sign Language, or other language.

7.6 Registered Nurses and Marital Status and Dependents

Nearly 74 percent of RNs were married or in a domestic partnership in 2008 (Appendix A, Table 54). Similar percentages of married and not married nurses reported having dependent adults in their households (15.2 percent v. 18.1 percent) (Table 7-11). In contrast, married nurses were twice as likely as nurses who were not married to have dependent children in the household (51.7 percent v. 25.3 percent).

Table 7-11. Marital status and presence of children or adult dependents at home

| Marital status | Dependent adults in home (percent) | Children in home (percent) | |
|------------------------------|---------------------------------------|-------------------------------|--|
| Married/Domestic partnership | 15.2 | 51.7 | |
| Not married | 18.1 | 25.3 | |

Source: 2008 National Sample Survey of Registered Nurses

Approximately 55 percent of RNs working in nursing reported having household dependents, either adult or children (Table 7-12 and Appendix A, Tables 54 and 55). That number decreases to 42.6 percent among RNs working outside of nursing, and decreases to 38.7 percent among those not working. This decrease was associated with lower numbers of RNs reporting dependent children in the household among those working outside of nursing and those not employed. The percentage of households with dependent adults was virtually the same across all three groups.



| | Dependent adults in home (percent) | Children in home (percent) | Both adults and children (percent) |
|----------------------------|---------------------------------------|-------------------------------|---------------------------------------|
| Working in nursing | 10.1 | 38.8 | 5.9 |
| Working outside of nursing | 10.6 | 27.2 | 4.8 |
| Not working | 10.3 | 25.1 | 3.3 |

Table 7-12. Employment of registered nurses, by presence of children or adult dependants in the home

Source: 2008 National Sample Survey of Registered Nurses

7.7 Registered Nurses and Household Income

Household income for the majority of RNs employed in nursing is equal to or significantly better than the median household income in 2008.¹ Approximately 40 percent of RNs in 2008 lived in households in which total household income was at least \$100,000 (Table 7-13 and Appendix A, Table 56). Among RNs employed full-time as staff nurses, 33.8 percent reported an annual household income of at least \$100,000. RNs in administrative and management positions were more likely to be in high-income households, with 54.9 percent reporting household income of \$100,000 or more. Total household income tended to be lower among RNs who were not employed in nursing.

Table 7-13. Annual household income of registered nurses, by employment status of the nurse

| | Annual household income less than or equal to \$50,000 (percent) | Annual household income \$50,001- \$100,000 (percent) | Annual household income greater than \$100,000 (percent) |
|-------------------------------------|--|---|---|
| All nurses | 12.5 | 47.5 | 40.1 |
| Nurses employed in nursing | 9.2 | 49.7 | 41.2 |
| Nurses not employed in nursing | 31.0 | 35.1 | 34.0 |
| All full-time nurses | 7.0 | 51.7 | 41.2 |
| Full-time staff nurses | 9.3 | 56.8 | 33.8 |
| Full-time management/administration | 2.8 | 42.3 | 54.9 |

¹ Median household income 2008 is \$52,029. Source: U.S. Census Bureau Quick Facts, http://quickfacts.census.gov/qfd/states/00000.html

7.8 Geography, Nurse Demographics, and Employment

There was significant variation in the distribution of the RN workforce across the country in 2008, with a two-fold difference among the States in the rate of employed RNs (Appendix A, Tables 57 and 58). Employment by setting also varied geographically (Figure 7-6 and Appendix A, Table 59). A smaller percentage of RNs in the New England region were employed in hospitals (57.1 percent) and ambulatory care settings (7.9 percent), while the percentage employed in nursing homes in New England (9.2 percent) was higher than other regions. RNs in the Pacific region were more likely to be employed in ambulatory care settings (13.2 percent) than RNs in other regions.




Figure 7-6. Distribution of employment settings, by geographic region*


*Excludes registered nurses for whom employment setting was not known.


Source: 2008 National Sample Survey of Registered Nurses

Earnings varied by geographic region (Figure 7-7 and Appendix A, Table 60). Median earnings for staff RNs in 2008 ranged from a low of \$50,000 in the West North Central and East South Central



regions to \$75,000 in the Pacific region. For nurses with a primary full-time position in administration or management, the range in median earnings among regions was comparably large, from \$65,000 to \$95,000.





*Excludes registered nurses for whom earnings was not known.

Source: 2008 National Sample Survey of Registered Nurses

Education levels vary notably by region. The East-South Central region had the smallest percentage of RNs who had obtained at least a bachelor's degree, with only 44.7 percent at the bachelor's or higher level in nursing or a nursing-related field (Figure 7-8 and Appendix A, Table 61). The highest percentage of RNs with at least a bachelor's degree in nursing or a nursing-related field was in the Pacific and New England regions.

70% 61.6% 60.1% 59.5% 60% 56.7% 55.6% 55.3% 55.1% 54.7% 53.6% 53.9% 53.0% 52.8% 51.9% 50.5% 50.3% 49.4% 49.3% 48.9% 50% 47.6% 44.7% 40% 30% 20% 10% 0% New England Middle Atlantic East North West North South Atlantic East South West South Mountain Pacific Total U.S. Central Central Central Central Nursing or nursing related Nursing and non-nursing

Figure 7-8. Percent of employed registered nurses with a bachelor's degree and higher, by region

Source: 2008 National Sample Survey of Registered Nurses

Median age of employed RNs varied little across regions, ranging from 44 to 48 years (Appendix A, Table 62). There were notable differences among geographic regions in the percentage of employed RNs representing racial/ethnic minority groups (Figure 7-9 and Appendix A, Table 63). The Pacific and West-South Central regions had the highest percentages of RNs who were non-White or Hispanic. In the West-North Central region, only 5.5 percent of RNs were from racial or ethnic minority groups; in the New England region, only 7.1 percent were from minority groups.



Pacific

Mountain

Total U.S.

35% 32.5% 30% 26.4% 25% 20.8% 20% 18.4% 17.8% 15.5% 15% 11.2% 11.1% 10% 7.1% 5.5% 5% 0%

West North South Atlantic

Central

East South

Central

West South

Central

Figure 7-9. Percent of employed registered nurses in racial/ethnic minority groups, by geographic region

Source: 2008 National Sample Survey of Registered Nurses

Central

New England Middle Atlantic East North

Internationally Educated Nurses (IENs)

8.1 Introduction

This chapter reviews the characteristics and employment of internationally educated nurses (IENs), formerly termed foreign-trained or foreign-educated nurses. These nurses have completed primary nursing education, often in British or U.S. modeled nursing education programs, outside of the United States or in the U.S. territories. Since the early 1900s, IENs have worked in the United States RN workforce. Prior to World War II, IENs came predominantly from Canada and western European English-speaking countries. Since that time, IENs have come to the United States from the Caribbean, Asia, South America, and most recently some African nations, in addition to Canada and Europe. Internationally educated nurses include RNs who individually immigrate for personal and/or economic reasons and also may include larger populations of IENs who are recruited to fill U.S. hospital nursing shortages by employment agencies specializing in international nursing recruitment.

In 2008, IENs from the Philippines continued to be the largest group of IENs in the U.S. workforce, followed by RNs from Canada. In recent years, IENs constitute a small but an increasing percentage of the U.S. RN workforce.

8.2 IEN Licensure and National Origin

In 2008 there were an estimated 165,539 IENs living in the United States, accounting for 5.4 percent of all licensed RNs (Table 8-1 and Appendix A, Table 65). Of these, 146,097 were employed in nursing in 2008, comprising 5.6 percent of the RN workforce. Approximately one-quarter of working IENs received their first U.S. nursing license in 2004 or later. The number of IENs is slowly growing: IENs licensed between 2004-2008 represented 8.1 percent of the total of recently licensed employed RNs, compared with 5.1 percent of those licensed before 2004.

IENs from the Philippines comprised just over 50 percent of all IENs and 52 percent of those working in nursing (Figure 8-1 and Appendix A, Table 65). Nearly 12 percent of IENs came from


Canada, and 9.6 percent were educated in India. The United Kingdom supplied 6 percent of IENs, while Korean and Nigerian IENs comprised 2.6 and 2.1 percent, respectively.

| | Total licensed nurses | Total IENs | IEN percent of total | Total employed nurses | Employed IENs | IEN percent of employed |
|----------------------|-----------------------------|------------|----------------------------|-----------------------------|------------------|-------------------------------|
| Licensed before 2004 | 2,618,494 | 128,028 | 4.9 | 2,169,846 | 111,576 | 5.1 |
| Licensed 2004-2008 | 444,668 | 37,511 | 8.4 | 426,753 | 34,521 | 8.1 |
| Total | 3,063,162 | 165,539 | 5.4 | 2,596,599 | 146,097 | 5.6 |

Table 8-1. Internationally educated nurses in the United States

Source: 2008 National Sample Survey of Registered Nurses





Source: 2008 National Sample Survey of Registered Nurses

The National Council of State Boards of Nursing reported an increase in the number of IEN graduates who passed the National Council Licensure Examination (NCLEX) after 2000; fewer than

5,000 IENs passed in 1998 but more than 22,000 IENs passed in 2007.¹ More than half of these nurses were from the Philippines in 2007 (13,342), and 4,053 were from India. The growth in the number of IENs taking and passing the NCLEX is another indicator of the growth in the number of IENs looking to live and work in the United States.

Most IENs did not obtain their first U.S. nursing license until 4 or more years after completing their initial RN education (Table 8-2). In fact, 33.5 percent of IENs did not obtain a U.S. license until 10 or more years after their initial RN degree was received.

| Table 8-2. | Years between initial nursing education and first U.S. licensure |
|------------|--|
|------------|--|

| | U.Seducated | Internationally educated |
|------------------|-------------|--------------------------|
| 0 years | 82.6 | 7.2 |
| 1 year | 15.8 | 6.2 |
| 2-3 years | 1.5 | 17.3 |
| 4-9 years | <0.1 | 35.9 |
| 10 or more years | <0.1 | 33.5 |

Source: 2008 National Sample Survey of Registered Nurses

8.3 IEN Demographic Characteristics

Half of RNs working in nursing in 2008—both IENs and U.S.-educated—were 46 years of age or older (Table 8-3). The median age at licensure of U.S.-educated RNs was younger than that of IENs, 25 years of age compared to 30 for IENs. Half of IENs arrived in the United States to work at least 6 years after completing their primary nursing education in their home country.

Table 8-3.Median current age and age at licensure of registered nurses employed in nursing,
by international education status

| | U.Seducated | Internationally educated |
|------------------|-------------|--------------------------|
| Current age | 47 | 46 |
| Age at licensure | 25 | 30 |

Source: 2008 National Sample Survey of Registered Nurses

Men comprised comparable percentages of recently licensed IENs and U.S.-educated nurses, at about 10 percent of each group (Table 8-4). A slightly higher percentage of IENs than of U.S.-educated nurses licensed before 2004 were male.

¹ Data provided by the National Council of State Boards of Nursing, not collected through the 2008 NSSRN.



| | | U.Seducated working nurses | | ally educated g nurses |
|----------------------|---------|-------------------------------|--------|---------------------------|
| | Number | Percent | Number | Percent |
| Licensed before 2004 | 132,449 | 6.4 | 8,496 | 7.6 |
| Licensed 2004-2008 | 38,296 | 9.8 | 3,500 | 10.1 |

Table 8-4.Men among U.S. and internationally educated employed nurses, by year of licensure

Source: 2008 National Sample Survey of Registered Nurses

More than 68 percent of IENs working in nursing in 2008 had at least a bachelor's degree in nursing or a nursing-related field, compared with 49.2 percent of U.S.-educated nurses (Table 8-5). A variety of external factors have influenced the educational makeup of the IEN workforce. One factor in the last two decades is the requirement from the U.S. Citizenship and Immigrations Services that IENs possess a minimum of a bachelor's degree to obtain a U.S. work visa.² In addition, many States and U.S. graduate nursing programs require that IENS obtain a U.S. RN license to enroll in a U.S. graduate nursing education program. A third factor is that there are predominantly bachelor's education programs for nurses, as opposed to ADN programs, in some of the countries from which many IENs immigrate, such as the Philippines. These factors mean that the IEN workforce educational attainment is significantly different than education of U.S.-educated RNs, the majority of whom receive their primary education in 2-year ADN programs. Therefore, IENs appear to be better educated as a group than U.S. RNs.

Table 8-5.Registered nurses working in nursing with at least a bachelor's degree, by locationof education

| | U.Seducated | Internationally educated |
|-------------------------|-------------|--------------------------|
| Nursing/nursing-related | 49.2 | 68.6 |
| Nursing or other field | 54.7 | 71.3 |

Source: 2008 National Sample Survey of Registered Nurses

The majority of IENs resided in a small number of States (Appendix A, Table 66). Approximately one-quarter of IENs lived in California, with New York, Florida, and Texas each home to 10 to 12 percent of IENs.

² U.S. Citizenship and Immigration Services <u>http://www.uscis.gov/portal/site/uscis</u>

8.4 IEN Employment

In 2008, 88.3 percent of IENs were employed in nursing (Appendix A, Table 65), slightly more than RNs in general (84.8 percent, Chapter 3). IENs were also more likely than their U.S.-educated counterparts to work full-time (Table 8-6).

Table 8-6. Registered nurses working full-time, by location of education and year of licensure

| | U.Seducated | Internationally educated |
|----------------------|-------------|--------------------------|
| Licensed before 2004 | 71.7 | 83.5 |
| Licensed 2004-2008 | 85.3 | 93.7 |

Source: 2008 National Sample Survey of Registered Nurses

Internationally educated nurses worked slightly more hours per year and per week in their principal nursing position than did U.S.-educated RNs (Table 8-7).

Table 8-7.Average hours worked per year and per week in principal nursing position, by full-
time and part-time status and location of education

| | U.Seducated | Internationally educated |
|------------------------|-------------|--------------------------|
| Average hours per year | | |
| Full-time employment | 2,202.7 | 2,232.4 |
| Part-time employment | 1,173.7 | 1,306.2 |
| Average hours per week | | |
| Full-time employment | 42.9 | 43.7 |
| Part-time employment | 24.0 | 27.6 |

Source: 2008 National Sample Survey of Registered Nurses

A greater percent of IENs held secondary nursing positions than did RNs in general. Nationwide,

12.2 percent of RNs who were employed full-time held secondary nursing jobs (Figure 3-11), while 17.2 percent of IENs have at least one additional RN position (Figure 8-2).




Figure 8-2. Secondary nursing positions held by IENs working full-time in their principal nursing position

Source: 2008 National Sample Survey of Registered Nurses

Hospitals were the employment setting for most IENs in 2008, regardless of whether they were licensed in the United States prior to 2004 or since 2004. Approximately three-quarters of recently licensed IENs were employed in hospital settings, with 71.8 percent similarly employed among those licensed before 2004 (Table 8-8 and Appendix A, Table 67). The employment pattern was quite different among U.S.-educated nurses, where 82.9 percent of those recently licensed were hospital-employed, yet only 57.5 percent of those licensed before 2004 were employed in hospitals. This suggests that IENS have more of a tendency to stay in hospital employment than U.S.-educated nurses.

| Table 8-8. | Number and percent of nurses employed in a hospital setting for their principal | | |
|------------|---|--|--|
| | nursing position, by location of initial education and year of licensure | | |

| | U.Seducated | | Internationally educated | |
|----------------------|-------------|---------|--------------------------|---------|
| - | Number | Percent | Number | Percent |
| Licensed before 2004 | 1,170,276 | 57.5 | 79,080 | 71.8 |
| Licensed 2004-2008 | 323,349 | 82.9 | 26,240 | 76.5 |
| Total | 1,493,625 | 61.6 | 105,320 | 72.9 |

U.S.-educated nurses were more than twice as likely to be found in ambulatory care settings compared with IENs (Table 8-9). However, a higher percentage of IENs were employed in academic education settings, compared with U.S.-educated RNs.

Table 8-9.Registered nurses employed in ambulatory care and academic education, by
location of initial education

| | U.Seducated | | Internationally educated | |
|--------------------|-------------|---------|--------------------------|---------|
| | Number | Percent | Number | Percent |
| Ambulatory care | 263,998 | 10.9 | 6,276 | 4.3 |
| Academic education | 88,656 | 3.7 | 9,490 | 6.6 |

Source: 2008 National Sample Survey of Registered Nurses

There were differences in the job titles of IENs compared with U.S.-educated RNs. More than 90 percent of all recently licensed nurses had staff nurse positions—91.2 percent of U.S.-educated RNs and 94.1 percent of IENs (Table 8-10). That percentage declines among those licensed prior to 2004 to 74 percent for IENs and 60.7 percent for U.S.-educated nurses. This may reflect the fact that IENs tend to stay in hospital employment positions.

Table 8-10.Registered nurses employed as staff nurses, by location of initial education and
year of licensure

| | U.Seducated | | Internationally educated | |
|----------------------|-------------|---------|--------------------------|---------|
| | Number | Percent | Number | Percent |
| Licensed before 2004 | 1,241,440 | 60.7 | 81,057 | 74.0 |
| Licensed 2004-2008 | 353,487 | 91.2 | 32,217 | 94.1 |
| Total | 1,594,927 | 65.6 | 113,274 | 78.8 |

Source: 2008 National Sample Survey of Registered Nurses

Nearly one-quarter of IENs intended to leave their current nursing position within the next 3 years, compared to 30.1 percent of U.S.-educated nurses (Table 8-11). However, the majority of these nurses intended to continue working in nursing.

Table 8-11. Employment intentions, by location of initial education

| | U.Seducated | Internationally educated |
|---|-------------|--------------------------|
| Plan to leave current position within 3 years | 30.1 | 24.7 |
| For those who plan to leave their job: | | |
| will continue to work in nursing | 72.1 | 72.3 |

This page deliberately blank.



9.1 Introduction

In this report recent graduates are defined as RNs who completed initial nursing education after 2000 (from 2001 to 2008). Within this group, another cohort was identified as the most recent graduates who completed their initial nursing education after 2004 (2005–2008). The sample of most recent graduates includes few RNs who graduated and obtained their licenses in 2008 because the survey sample was selected in early 2008.

In 2008, RNs who completed initial registered nursing programs after 2000 comprised nearly 20 percent of all RNs in the United States and 22.3 percent of those employed. These nurses are indicative of expansions in RN education programs responding to a national RN shortage. When these recent graduates completed their nursing program, they were nearly 4 years older at graduation than RNs who graduated before 2001. Many more recent graduates completed postsecondary education in another field prior to starting their RN education, and a considerable number worked in a health care position before completing initial RN education. These new graduates are more diverse than previous cohorts, with men and minorities comprising a greater percentage. More than 95 percent of new nurses were employed in nursing positions, and they were most likely to be employed in a hospital and as a staff nurse, compared with other employment setting and job title options. They changed employers at a rate of 17.7 percent between 2007 and 2008, often to advance their careers. As expected, most recent graduates had lower salaries and a narrower range of job titles than did RNs who graduated between 2001 and 2004.

9.2 Demographic Characteristics of Recent Registered Nurse Graduates

The number of RNs who received their first registered nursing degree between 2001 and 2008, and live in the United States, is estimated to be 603,402, comprising nearly 20 percent of the total RN population in 2008 (Table 9-1). The earlier group of these RNs—who graduated between 2001 and 2004—represent nearly 10 percent of all RNs and more than 10 percent of RNs employed in

nursing. Nearly 12 percent of RNs employed in nursing graduated from their first RN education program between 2005 and 2008.

| | Total estimated population | Graduated before 2001 | Graduated 2001-2004 | Graduated 2005-2008 |
|--|----------------------------------|-----------------------|------------------------|------------------------|
| Number residing in the United States | 3,063,162 | 2,459,780 | 289,760 | 313,622 |
| Percent of RN population Number employed residing in the United | 100.0 | 80.3 | 9.5 | 10.2 |
| States | 2,596,599 | 2,019,906 | 273,833 | 302,859 |
| Percent of employed nurses | 100.0 | 77.8 | 10.6 | 11.7 |

Table 9-1. Estimated population of registered nurses, by graduation cohort

Source: 2008 National Sample Survey of Registered Nurses

9.3 Recent Registered Nurse Graduates and Education

Recent graduates in 2008 were more likely to have received initial RN education in a bachelor's or master's level nursing program than were RNs who graduated before 2001 (Table 9-2). About 39 percent of RNs who graduated after 2000 had a BSN as their initial nursing education, and about 1 percent entered with a master's degree. The most notable change across the graduation cohorts is that diploma nursing education was rare among recent graduates (3 percent.)

Table 9-2. Initial education of registered nurses, by graduation cohort

| | Graduated before 2001 | Graduated 2001-2004 | Graduated 2005-2008 |
|-----------------------|-----------------------|---------------------|---------------------|
| Initial education | (percent) | (percent) | (percent) |
| Diploma | 24.6 | 3.5 | 3.1 |
| Associate | 42.7 | 56.4 | 56.7 |
| Bachelor's and higher | 32.7 | 40.1 | 40.3 |

Source: 2008 National Sample Survey of Registered Nurses

Very few of the RNs who graduated between 2001 and 2004 (2.8 percent), and an even smaller percentage of the most recent graduates (less than 1 percent) were internationally educated, compared to 6.3 percent of those who graduated before 2001 (Table 9-3). This is not surprising because nearly 70 percent of internationally educated nurses (IENs) did not receive their first U.S. license until at least 4 years after completing their initial RN education and 33.5 percent did not receive a U.S. license until 10 or more years after initial education (Table 8-2).¹ Thus, few IENs who

¹ Internationally educated nurses (IENs) are registered nurses with an active U.S. nursing license who completed their initial nursing education outside of the United States or in the U.S. territories.

graduated in the most recent cohort (2005–2008) were likely to have immigrated by 2008 into the United States.

| | Graduated before 2001 | Graduated 2001-2004 | Graduated 2005-2008 |
|--------------------------|-----------------------|---------------------|---------------------|
| Initial education | (percent) | (percent) | (percent) |
| U.Seducated | 93.7 | 97.2 | 99.2 |
| Internationally educated | 6.3 | 2.8 | 0.8 |

Table 9-3. Location of initial education of nurses, by graduation cohort

Source: 2008 National Sample Survey of Registered Nurses

Even among the most recent graduates, IENs were more likely to have obtained their initial education in a bachelor's program (Table 9-4). Among IENs who graduated since 2001, and were licensed in the United States, more than 74 percent had received a bachelor's degree.

Table 9-4. Initial nursing education, by graduation cohort and location of education

| | U.Seducated (percent) Graduated Graduated before 2001 2001–2008 | | Internationally educated (percent) | |
|-----------------------|--|------|---------------------------------------|------------------------|
| Initial Education | | | Graduated before 2001 | Graduated 2001–2008 |
| Diploma/Associate | 69.0 | 60.5 | 41.8 | 25.8 |
| Bachelor's and higher | 31.0 | 39.5 | 58.2 | 74.2 |

Source: 2008 National Sample Survey of Registered Nurses

The average age at graduation from initial RN education has been increasing over time (Figure 2-2). RNs who graduated before 2001 completed their nursing programs at an average age of 26.7 years, while those who graduated after 2000 were on average a little over 30 years old at graduation (Table 9-5 and Appendix A, Table 2). More than half of RNs who completed their initial education prior to 2001 did so when they were under 25 years old; only 34.9 percent of 2001–2004 graduates and 32 percent of 2005–2008 graduates were younger than 25 years old at graduation.

Table 9-5. Average age at graduation and age distribution at graduation, by graduation cohort

| Age at graduation | Graduated before 2001 (percent) | Graduated 2001-2004 (percent) | Graduated 2005-2008 (percent) |
|-------------------|------------------------------------|----------------------------------|----------------------------------|
| Average age | 26.7 | 30.1 | 30.8 |
| <25 years | 56.7 | 34.9 | 32.0 |
| 25–29 years | 17.1 | 24.7 | 23.5 |
| 30–39 years | 18.4 | 24.9 | 27.0 |
| 40-49 years | 7.0 | 12.0 | 13.2 |
| 50 or more years | 0.8 | 3.6 | 4.3 |

In addition to being older, recently graduated RNs are more likely to have received a postsecondary education degree prior to entering nursing (Table 9-6).

Table 9-6.Registered nurses with postsecondary education degrees before starting initial
nursing education program

| | Graduated before 2001 (percent) | Graduated 2001–2004 (percent) | Graduated 2005-2008 (percent) |
|-----------------------------|---------------------------------------|-------------------------------------|-------------------------------------|
| No degree before RN degree | 81.5 | 68.3 | 62.9 |
| Had degree before RN degree | 18.5 | 31.7 | 37.1 |

Source: 2008 National Sample Survey of Registered Nurses

The most common fields of study for RNs who completed their initial RN education between 2005 and 2008 and who held a postsecondary degree prior to their RN education were health-related fields (32.5 percent); liberal arts, social sciences, or humanities (28.2 percent); biological or physical science (17.3 percent); and business or management (11.3 percent) (Table 9-7 and Appendix A, Table 4). More than half of recent graduates with a postsecondary degree had received a bachelor's degree in another field. The most recent entrants to the registered nursing profession were more likely than previous entrants to be entering nursing after study in another field or as a "second career."

Table 9-7.Fields of study for registered nurses who had postsecondary degrees before initial
nursing education, by graduation cohort

| | Graduated before 2001 (percent) | Graduated 2001–2004 (percent) | Graduated 2005-2008 (percent) |
|---|---------------------------------------|-------------------------------------|-------------------------------------|
| Health-related field | 44.0 | 38.6 | 32.5 |
| Biology/physical science | 11.9 | 18.2 | 17.3 |
| Business/management | 8.6 | 8.8 | 11.3 |
| Education | 7.5 | 3.9 | 4.0 |
| Liberal arts/social sciences/humanities | 24.0 | 25.4 | 28.2 |
| Other nonhealth field* | 4.0 | 5.2 | 6.7 |

*Other includes law, computer science, and social work.

Source: 2008 National Sample Survey of Registered Nurses

Recent RN graduates are more likely to have worked in a health-related occupation prior to their initial RN education (Table 9-8). Among RNs who graduated prior to 2001, 65.5 percent had worked in a health occupation; this rate is more than 73 percent among recent graduates. Recent graduates are somewhat less likely than those who graduated before 2001 to have worked as nursing

aides/assistants and more likely to have worked as clerks in health care settings or in allied health fields. More than 20 percent of RNs, regardless of graduation cohort, were LPN/LVNs prior to initial education.

| | Graduated before 2001 (percent) | Graduated 2001–2004 (percent) | Graduated 2005–2008 (percent) |
|---|---------------------------------------|-------------------------------------|-------------------------------------|
| Percent with health-related employment before nursing | | | |
| education | 65.5 | 75.9 | 73.1 |
| Jobs of those with prior health employment | | | |
| Nursing aide/assistant | 70.0 | 64.5 | 62.2 |
| Home health aide/assistant | 8.0 | 11.2 | 10.8 |
| LPN/LVN | 21.1 | 21.6 | 21.6 |
| Emergency medical technician/paramedic | 4.4 | 8.9 | 8.2 |
| Medical assistant | 5.2 | 8.4 | 9.0 |
| Dental assistant | 2.3 | 2.8 | 2.0 |
| Allied health/technician | 7.4 | 10.0 | 10.6 |
| Manager in health setting | 2.0 | 3.2 | 4.2 |
| Clerk in health setting | 10.9 | 13.7 | 16.7 |
| Military medical corps | 2.3 | 2.1 | 1.5 |
| Other type of health-related position * * | 2.6 | 5.1 | 4.4 |

Table 9-8.Employment in a health occupation by registered nurses before initial nursing
education, by graduation cohort*

*Entries for prior health employment add to more than 100 because an individual registered nurse may have held more than one type of prior health employment.

**Other includes medical doctor, midwife, student nurse externship, mental health/psychiatric, and health education.

Source: 2008 National Sample Survey of Registered Nurses

Recently graduated RNs relied upon different sources of support for their initial education than did RNs who graduated prior to 2001 (Table 9-9). A greater percentage of recent graduates relied on earnings from health care jobs. This is not surprising since recent graduates also were more likely than those who graduated before 2001 to have worked in a health occupation prior to their initial RN education. Recent graduates were also more likely than earlier graduates to have received nursing education financing from federally assisted loans, other types of loans, State and local government scholarships, nongovernment scholarships, personal savings, and employer tuition reimbursement. They were less likely to have received support from "other family resources," which may indicate that these graduates, who were older at graduation than were previous cohorts, received little nursing education financial support from parents or other relatives.

| | Graduated before 2001 (percent) | Graduated 2001–2004 (percent) | Graduated 2005-2008 (percent) |
|--|---------------------------------------|-------------------------------------|-------------------------------------|
| Earnings from health care job | 29.8 | 34.8 | 32.1 |
| Earnings from other job | 22.5 | 23.3 | 24.9 |
| Earnings from other household members | 23.9 | 24.4 | 27.4 |
| Personal savings | 18.3 | 20.0 | 25.5 |
| Other family resources | 45.9 | 29.3 | 29.7 |
| Employer tuition reimbursement | 8.4 | 16.8 | 17.5 |
| Federal traineeship/scholarship | 14.2 | 22.1 | 23.6 |
| Federally assisted loan | 24.6 | 45.7 | 50.5 |
| Other type of loan | 9.1 | 14.4 | 21.0 |
| State/local government scholarship/grant | 14.5 | 20.8 | 25.5 |
| Nongovernment scholarship | 13.1 | 17.4 | 17.6 |
| Other resources | 4.8 | 4.5 | 5.9 |

 Table 9-9.
 Sources of financing for initial nursing education, by graduation cohort*

*Entries for sources of financing add to more than 100 because more than one source of financing may have been used by an individual registered nurse.

Source: 2008 National Sample Survey of Registered Nurses

More than 10 percent of RNs who completed their initial nursing education between 2001 and 2004 later completed additional degrees. Bachelor's degrees were completed by 9.6 percent of graduates from diploma or ADN programs; master's degrees or doctorates were completed by more than 10 percent of BSN graduates and almost 2 percent of diploma or ADN graduates (Table 9-10).

Table 9-10.Highest level of nursing or nursing-related education for recent registered nurse
graduates, by initial nursing education (2001–2004)

| Highest education | Diploma/associate degree (percent) | Bachelor's degree (percent) | Total (percent) |
|--|--|-----------------------------------|--------------------|
| Diploma/Associate | 88.7 | (percent) | 53.2 |
| Bachelor's | 9.6 | 89.5 | 40.6 |
| Master's or doctorate | 1.7 | 10.5 | 6.4 |
| Percent who have received additional degrees | 12.4 | 11.1 | |

Source: 2008 National Sample Survey of Registered Nurses

The most recent RN graduates work in the State in which they were educated.

- Among RNs who graduated in the past 5 years, 80.8 percent worked in the State where they were educated.
- Among RNs who graduated 6 to 10 years ago, 75.8 percent worked in the State where they were educated.

• For RNs who graduated 26 or more years ago, only 58.9 percent still worked in the State where they were educated.

9.4 Recent Registered Nurse Graduates and Diversity

Recently graduated RNs are substantially more ethnically and racially diverse and have more male graduates than nurses who graduated prior to 2001. Nearly 10 percent of graduates between 2005 and 2008 were men, compared with only 5.9 percent of graduates prior to 2001 (Table 9-11).

| Table 9-11. | Gender of registered nurses, by graduation cohort |
|-------------|---|
|-------------|---|

| | Graduated before 2001 (percent) | Graduated 2001-2004 (percent) | Graduated 2005-2008 (percent) |
|--------|------------------------------------|----------------------------------|----------------------------------|
| Male | 5.9 | 9.1 | 9.8 |
| Female | 94.1 | 90.9 | 90.2 |

Source: 2008 National Sample Survey of Registered Nurses

The RN population is becoming more ethnically diverse (Table 9-12). Among recent RN graduates in 2008, the percentage of RNs from racial/ethnic minority groups was higher, compared with previous cohorts (22 percent v. 15.5 percent). The largest ethnic groups after White, non-Hispanic were Blacks (4.9 percent prior to 2001; 7.3 percent from 2001 to 2008) and Hispanics (2.8 percent prior to 2001; 6.7 percent from 2001 to 2008). The percentage of graduates who were non-Hispanic Asian has dropped slightly with more recent cohorts accounting for 5 percent of recent graduates (2001–2008) and 5.7 percent prior to 2001.

| Table 9-12. | Racial/ethnic composition of registered nurse population, by graduation cohort |
|-------------|--|
|-------------|--|

| Race | Graduated before 2001 (percent) | Graduated 2001-2008 (percent) |
|---|------------------------------------|----------------------------------|
| White, non-Hispanic | 84.5 | 78.0 |
| Black/African American, non-Hispanic | 4.9 | 7.3 |
| Asian, non-Hispanic | 5.7 | 5.0 |
| Hispanic/Latino, any race | 2.8 | 6.7 |
| Other or two or more races, non-Hispanic* | 2.1 | 3.0 |

*Other races include American Indian and Pacific Islander.

9.5 Family Situation of Recent Nurse Graduates

In 2008, 71.8 percent of 2001–2004 graduates and 62.1 percent of 2005–2008 graduates were married or in a domestic partnership (Table 9-13).

More than half of recent RN graduates had children living at home, with 61.4 percent of 2001–2004 graduates and 48.1 percent of 2005–2008 graduates reporting the presence of children. Both groups of recent graduates were more likely than pre-2001 graduates to have children under 6 years old at home. More than 50 percent of graduates prior to 2001 and the most recent graduates had no children at home.

| Table 9-13. | Marital status and presence of children, by graduation cohort |
|-------------|---|
| | |

| | Graduated before 2001 (percent) | Graduated 2001–2004 (percent) | Graduated 2005–2008 (percent) |
|--------------------------------|---------------------------------------|-------------------------------------|-------------------------------------|
| Married/Domestic partnership | 75.5 | 71.8 | 62.1 |
| Widowed/Divorced/Separated | 18.3 | 12.3 | 9.6 |
| Never married | 6.2 | 15.9 | 28.3 |
| No children | 60.6 | 38.6 | 51.9 |
| Children under 6 years old | 6.4 | 27.0 | 16.6 |
| Children 6 years old and older | 27.5 | 23.1 | 23.2 |
| Children in both age groups | 5.6 | 11.3 | 8.4 |

Source: 2008 National Sample Survey of Registered Nurses

9.6 Recent Registered Nurse Graduates and Employment

The vast majority of recent RN graduates were employed in nursing positions in 2008 (Table 9-14). Among the most recent group, 83.8 percent worked full-time. In contrast, nurses who graduated from 2001 to 2004 were somewhat more likely to work part-time in their principal nursing positions. They were also more likely to have children living at home. Recent graduates who have children living at home were less likely to work full-time (Table 9-15).

| | Graduated before 2001 (percent) | Graduated 2001–2004 (percent) | Graduated 2005–2008 (percent) |
|---------------------------------|---------------------------------------|-------------------------------------|-------------------------------------|
| Employed full-time in nursing | 59.2 | 75.4 | 83.8 |
| Employed part-time in nursing | 22.9 | 19.1 | 12.8 |
| Not employed in nursing | 17.9 | 5.5 | 3.4 |
| Total employed in nursing, 2008 | 82.1 | 94.5 | 96.6 |

Table 9-14. Employment of registered nurses, by graduation cohort

Source: 2008 National Sample Survey of Registered Nurses

Table 9-15. Employment of new registered nurse graduates, by presence of children

| Graduates 2001-2008 | No children at home (percent) | Children at home (percent) |
|-------------------------------|----------------------------------|-------------------------------|
| Employed full-time in nursing | 85.7 | 74.8 |
| Employed part-time in nursing | 10.0 | 20.7 |
| Not employed in nursing | 4.3 | 4.5 |

Source: 2008 National Sample Survey of Registered Nurses

The majority of recently graduated RNs reported their principal employment setting as a hospital. Hospitals employed 83.3 percent of RNs graduating from 2005 to 2008, and more than 75 percent of RNs graduating from 2001 to 2004 (Table 9-16). Only 56.7 percent of RNs who graduated prior to 2001 worked in hospitals in 2008. In contrast, the percentage of recently graduated RNs working in hospitals in the 2004 NSSRN was somewhat lower—78.8 percent of RNs who had graduated in the 4 years prior to 2004 worked in hospitals, and 69.2 percent of those who graduated 5 to 8 years prior to 2004 worked in hospitals at the time of that Survey.

Table 9-16.Employment setting of principal nursing position, by graduation cohort, 2008 and
2004

| | 2008 NSSRN | | | 2004 NSSRN | |
|---|-----------------------------|----------------------------|----------------------------|----------------------------|----------------------------|
| | Graduated before 2001 | Graduated 2001- 2004 | Graduated 2005- 2008 | Graduated 1997- 2000 | Graduated 2001- 2004 |
| Hospital settings, without hospital-based | | | | | |
| extended care | 56.7 | 75.3 | 83.3 | 69.2 | 78.8 |
| Nonhospital extended care | 5.6 | 4.0 | 4.1 | 6.4 | 5.7 |
| Academic education | 4.4 | 2.4 | 1.1 | 1.7 | 1.4 |
| Home health/public/community health | 11.4 | 7.1 | 5.4 | 8.7 | 4.8 |
| Ambulatory care | 11.8 | 8.1 | 4.5 | 8.4 | 6.1 |
| Other* | 10.1 | 3.1 | 1.6 | 5.6 | 3.2 |

*Other includes school health, occupational health, hospital-based extended care, insurance, benefits, and utilization review.

The most common job title held by all working RNs in 2008 was staff nurse or an equivalent title. Recent RN graduates were even more likely to have this title than were RNs who graduated prior to 2001 (Table 9-17). Over time, RNs are more likely to have job titles other than staff nurse. For example, only 1.4 percent of 2005–2008 graduates have a middle or senior management job title in 2008, while 3.8 percent of 2001–2004 graduates have this title.

| Job Title | Graduated before 2001 (percent) | Graduated 2001–2004 (percent) | Graduated 2005–2008 (percent) |
|-----------------------------|---------------------------------------|-------------------------------------|-------------------------------------|
| First-line management | 4.6 | 3.5 | 1.3 |
| Middle or senior management | 10.1 | 3.8 | 1.4 |
| Instruction | 4.4 | 1.7 | 0.5 |
| Staff nurse | 60.5 | 80.7 | 92.2 |
| Nurse practitioner | 4.5 | 2.7 | 0.4 |
| Patient coordinator | 6.2 | 3.4 | 1.9 |
| Other job titles* | 9.7 | 4.3 | 2.2 |

Table 9-17. Job titles of registered nurses, by graduation cohort

*Other includes registered nurses who reported no job title.

Source: 2008 National Sample Survey of Registered Nurses

The number of hours worked weekly and annually by full-time RNs in 2008 was similar across the graduation cohorts. However, among RNs who work part-time, the most recent cohorts worked more hours than did RNs who graduated before 2001 (Table 9-18). RNs graduating in 2005–2008 who worked part-time averaged 27.1 hours per week and 1,318 hours per year in their principal nursing positions. Part-time RNs who graduated from 2001 to 2004 averaged 24.9 hours per week and 1,232 hours per year, while part-time RNs who graduated before 2001 worked an average of 23.8 hours per week and 1,162 hours per year.

Table 9-18.Average hours worked per year and per week by registered nurses, by full-time/part-
time status and graduation cohort

| | Graduated before 2001 | Graduated 2001-2004 | Graduated 2005-2008 |
|--------------------|--------------------------|------------------------|------------------------|
| Employed full-time | | | |
| Hours per year | 2,220 | 2,178 | 2,141 |
| Hours per week | 43.3 | 42.2 | 41.4 |
| Employed part-time | | | |
| Hours per year | 1,162 | 1,232 | 1,318 |
| Hours per week | 23.8 | 24.9 | 27.1 |

While all groups of graduates reported high levels of satisfaction with their nursing position, recent RN graduates were less satisfied with their principal nursing positions than were RNs who graduated before 2001. Among RNs who graduated from 2005 to 2008, 78.2 percent were moderately or extremely satisfied, compared with 82.1 percent of RNs who graduated before 2001 (Table 9-19). The satisfaction of RNs who graduated from 2001 to 2004 was similar to that of the 2005 to 2008 graduates.

| Satisfaction of employed nurses | Graduated before 2001 (percent) | Graduated 2001–2004 (percent) | Graduated 2005–2008 (percent) |
|-----------------------------------|---------------------------------------|-------------------------------------|-------------------------------------|
| Extremely satisfied | 31.7 | 22.0 | 20.3 |
| Moderately satisfied | 50.5 | 54.7 | 57.8 |
| Neither satisfied or dissatisfied | 7.2 | 10.5 | 9.7 |
| Moderately dissatisfied | 8.2 | 10.6 | 9.7 |
| Extremely dissatisfied | 2.5 | 2.3 | 2.4 |

Table 9-19. Satisfaction with principal nursing position, by graduation cohort

Source: 2008 National Sample Survey of Registered Nurses

9.7 Recent Registered Nurse Graduates and Job Changes

Many factors can lead to job changes, such as job dissatisfaction, relocation, career advancement, and other reasons. In 2008, nearly 40 percent of recent graduates planned to leave their current jobs within 3 years, compared with 27.1 percent of pre-2001 graduates (Table 9-20). Nonetheless, the vast majority of recently graduated RNs intended to remain in the nursing profession, with only 3.3 percent of those who intended to leave their current job stating that they did not plan to stay in nursing.

Table 9-20. Intentions regarding nursing employment, by graduation cohort

| | Graduated before 2001 (percent) | Graduated 2001-2008 (percent) |
|--|------------------------------------|----------------------------------|
| Plans regarding current position | | |
| No plans to leave job | 57.8 | 42.8 |
| Undecided about plans | 15.1 | 17.8 |
| Have left job or plan to leave in 12 months | 14.5 | 23.2 |
| Plan to leave in 1 to 3 years | 12.6 | 16.2 |
| Total that plans to leave job within 3 years | 27.1 | 39.3 |
| For those who plan to leave their job | | |
| Plan to remain in nursing work | 77.9 | 96.7 |
| Plan to leave nursing | 22.1 | 3.3 |

Recent graduates change jobs somewhat more frequently than RNs who graduated prior to 2001. Of all RNs employed in March 2008, 14.9 percent of graduates from 2001 to 2004 had changed employers for their principal nursing position in the previous year and 7.5 percent had changed positions but were with the same employer (Table 9-21). Among the 2005 to 2008 graduates, 14.8 percent had changed employers, and 7.1 percent had changed positions with the same employer.

Table 9-21.Job changes between March 2007 and March 2008, for registered nurses employed
in 2008, by graduation cohort

| | Graduated before 2001 (percent) | Graduated 2001–2004 (percent) | Graduated 2005–2008 (percent) |
|-----------------------------------|---------------------------------------|-------------------------------------|-------------------------------------|
| Not employed in 2007 | 2.9 | 2.5 | 28.0 |
| Same position, same employer | 82.5 | 75.0 | 50.0 |
| Different position, same employer | 5.3 | 7.5 | 7.1 |
| Different employer | 9.4 | 14.9 | 14.8 |

Source: 2008 National Sample Survey of Registered Nurses

In 2008, reasons given for job changes varied, but among all RNs the most common reasons were related to working conditions such as the adequacy of staffing, physical demands of the job, stress, or the work environment (Table 9-22). Recent graduates were more likely than were pre-2001 graduates to report that they changed jobs for career reasons, such as for career advancement, or because they were interested in another job. RNs who graduated between 2001 and 2008 also were more likely than the older graduates to report that personal or family reasons drove their job change.

| Reasons for change | Graduated before 2001 (percent) | Graduated 2001–2004 (percent) | Graduated 2005-2008 (percent) | Overall graduated 2001–2008 (percent) |
|--------------------|---------------------------------------|-------------------------------------|-------------------------------------|---|
| Career | 32.9 | 41.0 | 46.8 | 43.9 |
| Personal/family | 30.0 | 38.4 | 28.7 | 33.5 |
| Workplace | 71.4 | 71.1 | 71.2 | 71.2 |

Table 9-22. Reasons for job changes, by graduation cohort*

*Percents may not add to 100 because registered nurses may have provided more than one reason for job change.

Source: 2008 National Sample Survey of Registered Nurses

Among RNs who changed employers, 45.4 percent of recent graduates changed employment settings, as compared with 56.4 percent of job changers who graduated prior to 2001. Recent graduates were more likely to change their 2007 employment setting if they were employed in nursing homes or extended care (25.8 percent), or public or community health (23.2 percent) (Table 9-23). Only 11.1 percent of recent graduates employed in hospitals in 2007 changed to a nonhospital employment setting if they changed employers by 2008.

| Setting in 2007 | Percent who left setting between 2007 and 2008 (percent) | | | |
|----------------------------|--|--|--|--|
| Hospital | 11.1 | | | |
| Nursing home/extended care | 25.8 | | | |
| Home health | 21.2 | | | |
| Public/community health* | 23.2 | | | |
| Ambulatory care | 20.8 | | | |
| Other settings** | 18.9 | | | |

Table 9-23.Employment setting changes for registered nurses who graduated 2001–2008

*Public/community health includes school health and occupational health.

****Other settings includes academic education and insurance/benefits/utilization review.**

Source: 2008 National Sample Survey of Registered Nurses

9.8 Recent Registered Nurse Graduates and Secondary Nursing Jobs

Secondary jobs in nursing were held by 13.3 percent of nurses who graduated from 2001 to 2004 and 11.1 percent of nurses who graduated from 2005 to 2008 (Table 9-24). The most common settings for second jobs among 2005–2008 graduates were hospitals (54.8 percent), nursing homes and extended care (13.8 percent), and home health (13.7 percent). Graduates in the 2001–2004 cohort reported hospitals (48.3 percent), ambulatory care (14.1 percent), and academic education settings (11.5 percent) as the most common locations for their secondary jobs.

| | Graduated before 2001 (percent) | Graduated 2001-2004 (percent) | Graduated 2005-2008 (percent) |
|------------------------------|---------------------------------------|-------------------------------------|-------------------------------------|
| Has a secondary nursing job? | | | |
| Yes | 12.8 | 13.3 | 11.1 |
| Νο | 87.2 | 86.7 | 88.9 |
| Setting of secondary job(s)* | | | |
| Hospital | 41.1 | 48.3 | 54.8 |
| Nursing home/extended care | 9.0 | 8.6 | 13.8 |
| Academic education program | 11.5 | 11.5 | - |
| Home health setting | 11.4 | 9.1 | 13.7 |
| Public/community health | 7.0 | 7.0 | 8.5 |
| Ambulatory care | 14.0 | 14.1 | 9.3 |
| Other employment settings | 10.3 | 10.9 | 4.2 |

*Secondary employment settings add to more than 100 because registered nurses could report more than one secondary setting.

-Too few cases to report estimated number (fewer than 30 respondents).

9.9 Recent Registered Nurse Graduates and Earnings

Average earnings from all nursing positions for the most recent cohort of nurses were \$52,994; the average earnings from just those with full-time principal nursing positions were \$53,924 (Table 9-25). The total nursing earnings of 2005–2008 graduates were lower than that of both RNs who graduated from 2001 to 2004 (\$57,489) and RNs who graduated before 2001 (\$62,903). Registered nurses from all three cohorts had similar earnings from part-time principal positions. However, recent graduates in principal positions that were part-time worked more hours per week on average than those who graduated before 2001; thus, their effective hourly rate was lower than that of the older nurses whose principal position was part-time.

Table 9-25.Average earnings from principal nursing position, and from all nursing positions, by
graduation cohort

| | Graduated | Graduated | Graduated |
|--|-------------|-----------|-----------|
| | before 2001 | 2001-2004 | 2005-2008 |
| Earnings from all nursing positions | 62,903 | 57,489 | 52,994 |
| Earnings from principal nursing position | | | |
| Full-time positions | 70,299 | 60,491 | 53,924 |
| Part-time positions | 36,975 | 36,686 | 35,236 |

Source: 2008 National Sample Survey of Registered Nurses

9

Registered Nurses Nearing Retirement 10

10.1 Introduction

This chapter describes the demographic, education, and employment characteristics of RNs 50 years of age and older. Historically, NSSRN data have shown a decline in work participation around this age. However, retirement literature fails to provide guidance on when RNs actually retire from the nursing workforce and when they allow their nursing license to lapse. In this chapter, "older RNs" are defined as the RN population that is 50 years of age or older, while "younger RNs" refers to RNs younger than 50 years of age in 2008.¹

10.2 Demographics Characteristics of Registered Nurses Nearing Retirement

Older RNs comprised 44.7 percent of the total RN population in 2008, compared with 41.1 percent in 2004 and 33.4 percent in 2000. The percentage of RNs who were 60 years and older increased from 13.6 percent in 2004 to 15.5 percent in 2008 (Figure 3-3 and Appendix A, Table 1).

Older RNs in 2008 were more likely to be female (94.7 percent) and White, non-Hispanic (87.2 percent) than younger nurses (Table 10-1).

Older nurses were also more likely than younger nurses to have received their initial nursing education through a diploma program and less likely to have received their initial education through a bachelor's program (Table 10-1). A greater percentage of older RNs have attained master's or doctorate degrees (16.8 percent) than have younger nurses (10.3 percent).

¹ The NSSRN data reflect only those RNs with active licenses and do not reflect RNs whose licenses have lapsed and have not been renewed.



| | Under 50 years old (percent) | 50 years or older (percent) | Total (percent) |
|--|------------------------------------|-----------------------------------|--------------------|
| Estimated total population | 1,694,088 | 1,369,074 | 3,063,162 |
| Race/Ethnicity | | | |
| White, non-Hispanic | 80.0 | 87.2 | 83.2 |
| Non-White or Hispanic | 20.0 | 12.8 | 16.8 |
| Gender | | | |
| Male | 7.7 | 5.3 | 6.6 |
| Female | 92.3 | 94.7 | 93.4 |
| Initial nursing education | | | |
| Diploma | 9.0 | 34.5 | 20.4 |
| Associate | 48.5 | 41.6 | 45.4 |
| Bachelor's and higher | 42.5 | 23.9 | 34.2 |
| Highest nursing or nursing-related education | | | |
| Diploma | 6.6 | 23.0 | 13.9 |
| Associate | 40.0 | 31.2 | 36.1 |
| Bachelor's | 43.1 | 28.9 | 36.8 |
| Master's or doctorate | 10.3 | 16.8 | 13.2 |

 Table 10-1.
 Demographic and educational characteristics of registered nurses, by age

Source: 2008 National Sample Survey of Registered Nurses

10.3 Employment Characteristics of Older Registered Nurses

A somewhat smaller share of older RNs worked in nursing in 2008, compared with younger RNs: 76.6 percent v. 91.4 percent. Moreover, the percentage of older RNs working in nursing declines steadily as they age; from 87.4 percent for those 50-54 years of age to 33.3 percent for those aged 75 and older (Figure 3-4). Registered nurses are more likely to work in nursing part-time rather than full-time as they age, particularly after age 65, yet more than 30 percent of RNs age 70 or older still work in nursing (Figure 10-1). This trend toward part-time work is particularly notable among nurses as they pass 65 years of age.

Registered Nurses Nearing Retirement




Figure 10-1. Percent of employed registered nurses working full-time and part-time, by age group

Source: 2008 National Sample Survey of Registered Nurses

10.3.1 Employment Positions and Settings

Older RNs in 2008 were less likely to work in hospital settings as they age (Table 10-2). The decline continues steadily with increasing age from nearly 55.9 percent for RNs aged 50-54 to 35.5 percent for RNs aged 65 and older. While a smaller percentage of older nurses than younger nurses worked in hospital positions, higher percentages were employed in nursing homes/extended care, academic education programs, and home health settings.

Registered Nurses Nearing Retirement

| | Age of nurses | | | | |
|---------------------------------------|---|---------------------------------|---------------------------------|---------------------------------|---|
| | Less than 50 years old (percent) | 50-54 years old (percent) | 55-59 years old (percent) | 60-64 years old (percent) | 65 or more years old (percent) |
| Hospital | 69.5 | 55.9 | 53.0 | 46.5 | 35.5 |
| Nursing home/extended care | 4.2 | 5.9 | 6.5 | 7.6 | 10.7 |
| Academic education program | 2.6 | 4.5 | 5.1 | 6.6 | 10.5 |
| Home health setting | 5.5 | 7.2 | 7.7 | 7.8 | 10.5 |
| Public/community health | 2.8 | 4.4 | 4.7 | 7.3 | 6.2 |
| School health service | 2.4 | 4.7 | 4.3 | 4.9 | 5.4 |
| Occupational health | 0.5 | 0.8 | 0.9 | 1.4 | 3.1 |
| Ambulatory care | 9.6 | 11.5 | 12.6 | 12.1 | 11.3 |
| Insurance/benefits/utilization review | 1.6 | 2.6 | 2.2 | 2.4 | 2.6 |
| Other | 1.4 | 2.5 | 3.0 | 3.4 | 4.2 |

Table 10-2.Employment settings of registered nurses in their principal nursing position, by age
group

Source: 2008 National Sample Survey of Registered Nurses

As with all other age groups, staff nurse is the job title for the majority of older RNs; however, a much lower proportion of older RNs were staff nurses compared with younger nurses.

Age of nurses Less than 50-54 years 60-64 years 55-59 years 65 or more 50 years old old old old years old (percent) (percent) (percent) (percent) (percent) Staff nurse 72.1 60.0 57.9 58.0 56.4 **First-line management** 3.8 5.2 4.4 3.8 3.9 Middle or senior management 6.9 10.4 11.3 11.0 9.6 Nurse practitioner 3.6 4.4 4.5 3.3 3.2 Instruction 3.1 3.9 4.1 5.6 7.3

6.8

9.2

7.5

10.3

6.4

11.9

7.5

12.2

Table 10-3. Job titles of registered nurses in their principal nursing position, by age group

*Other includes consultant, midwife, CNS, NA, research, informatics, surveyor/auditor/regulator, and patient education.

4.4

6.0

Source: 2008 National Sample Survey of Registered Nurses

Patient coordinator

Other*

The majority of all RNs in 2008 were moderately or extremely satisfied with their jobs, regardless of the setting in which they worked. Satisfaction among older RNs was slightly higher than among younger RNs across employment settings. Among older RNs, the highest proportion of satisfied nurses was employed in ambulatory settings; for younger RNs, highest satisfaction was among nurses in academic education settings. The lowest proportion of moderately or extremely satisfied RNs across both age groups worked in nursing homes, followed by hospitals (Table 10-4).



| | Hospital | Nursing home/ extended care | Academic education | Home health | Public/ community health* | Ambulatory | Other** |
|---|----------|--------------------------------------|--------------------|----------------|---------------------------------|------------|---------|
| Percent moderately or extremely satisfied | | | | | | | |
| Under age 50 | 78.9 | 72.5 | 86.4 | 82.2 | 82.7 | 82.9 | 81.4 |
| Age 50 or older | 81.6 | 76.4 | 86.7 | 83.5 | 86.2 | 88.7 | 81.6 |

Table 10-4. Job satisfaction, by principal employment setting

*Public/community health includes school health and occupational health.

**Other includes insurance, benefits, and utilization review.

Source: 2008 National Sample Survey of Registered Nurses

10.3.2 Hours Worked by Older Registered Nurses

In 2008, older RNs contributed significantly to the hours worked in nursing. Those aged 50–54 years and employed full-time worked an average of 43.7 hours per week (Table 10-5), slightly more hours per week than full-time nurses under age 50, who averaged 42.6 hours of work per week. Part-time nurses aged 50 to 54 years also worked more hours per week than did part-time nurses under 50 years old. Beginning at age 60, there is a steady decline in hours worked among part-time nurses.

Table 10-5.Average hours worked per week for registered nurses in their principal nursing
position, by age

| | | | Age of | nurses | | |
|--|------------------------------|--------------------|--------------------|--------------------|--------------------|----------------------------|
| | Less than 50 years old | 50-54 years old | 55-59 years old | 60-64 years old | 65-69 years old | 70 or more years old |
| Average hours worked, full- time nurses | 42.6 | 43.7 | 43.6 | 42.0 | 42.7 | 40.9 |
| Average hours worked, part- time nurses | 24.6 | 25.7 | 25.2 | 22.3 | 19.1 | 18.4 |

Source: 2008 National Sample Survey of Registered Nurses

10.3.3 Secondary Nursing Employment

Older RNs were less likely to have secondary positions in nursing than younger nurses, and the share with more than one nursing job generally declines with age (Table 10-6). Under age 65, nurses whose principal position was part-time were more likely to have secondary nursing employment than were those whose principal position was full-time.




Table 10-6. Percent of employed registered nurses with secondary nursing positions, by age

Source: 2008 National Sample Survey of Registered Nurses

10.3.4 Earnings of Older Registered Nurses

Older RNs in 2008 earned more on average than younger RNs from nursing work. This may be due to a greater percentage of older RNs who were employed in management and other higher-paying positions. As nurses age beyond 50 years, however, their average annual earnings decline (Table 10-7). Among part-time RNs, this is likely associated with the drop in the number of hours they work per week. The number of hours worked per week by older RNs who were employed full-time does not decline notably with age; thus, other factors such as changes in employment settings, and the shift of older RNs away from higher paying hospital positions to lower paying home care or ambulatory care settings, may be associated with the decline in annual earnings.

Table 10-7. Average earnings from principal nursing employment, by age

| | Full-time p | rincipal position | Part-time principal position | | | |
|--------------------|---------------|--------------------|------------------------------|--------------------|--|--|
| | All positions | Principal position | All positions | Principal position | | |
| Under 50 years old | \$66,918 | \$64,689 | \$41,614 | \$37,946 | | |
| 50-54 years old | 73,514 | 71,061 | 45,904 | 40,996 | | |
| 55-59 years old | 72,895 | 70,775 | 43,394 | 38,992 | | |
| 60-64 years old | 71,473 | 69,560 | 35,618 | 33,356 | | |
| 65-69 years old | 69,189 | 67,559 | 26,011 | 25,052 | | |
| 70 years and older | 64,981 | 64,522 | 23,956 | 22,539 | | |

Source: 2008 National Sample Survey of Registered Nurses

10.4 Job Changes and Intentions Regarding Nursing Work

While the majority of all RNs in 2008 remained employed with the same employer, older RNs demonstrated less overall job mobility when compared with younger RNs (Table 10-8). For registered nurses 50 to 59 years old, the percent remaining in the same job between 2007 and 2008

was higher than for younger RNs; for RNs between 60 and 69 years, an increasing percentage change to different employers. As RNs grow older, and especially after age 60, retirement and workplace issues become the dominant reasons for employment changes (Table 10-9).

Table 10-8.Registered nurses employed in nursing in 2007 who changed position or employer,
by age

| | | | Age of | nurses | | |
|--|------------------------------|--------------------|--------------------|--------------------|--------------------|----------------------------|
| | Less than 50 years old | 50-54 years old | 55-59 years old | 60-64 years old | 65-69 years old | 70 or more years old |
| Not employed in 2008 | 1.1 | 1.2 | 1.7 | 3.1 | 6.3 | 6.7 |
| Same position, same employer Different position, same | 77.5 | 84.1 | 83.3 | 82.4 | 76.9 | 80.8 |
| employer | 7.1 | 4.9 | 5.2 | 4.3 | 6.1 | - |
| Different employer | 14.4 | 9.7 | 9.8 | 10.2 | 10.7 | - |

- Too few cases to report estimated percent (fewer than 30 respondents).

Source: 2008 National Sample Survey of Registered Nurses

Table 10-9. Reasons for job changes, by age group

| | | | Age of | nurses | | |
|-----------------|---|---------------------------------|---------------------------------|---------------------------------|---------------------------------|--------------------------------------|
| | Less than 50 years old (percent) | 50-54 years old (percent) | 55-59 years old (percent) | 60-64 years old (percent) | 65-69 years old (percent) | 70 or more years old (percent) |
| Retirement | - | - | 8.4 | 31.6 | 49.6 | 45.8 |
| Career | 40.1 | 31.4 | 33.2 | 20.7 | 15.5 | - |
| Personal/family | 32.6 | 29.3 | 28.5 | 27.9 | 25.2 | - |
| Workplace | 73.3 | 73.9 | 71.3 | 63.0 | 49.4 | 48.1 |

- Too few cases to report estimated percent (fewer than 30 respondents).

Source: 2008 National Sample Survey of Registered Nurses

As described in Chapter 3, in 2008 more than 50 percent of all employed RNs, including older RNs, did not plan to leave their current nursing jobs. In fact, nurses 50 to 59 years old were less likely to report an intention to leave their current nursing position within 3 years than were nurses under 50 years old (Table 10-10). Above age 60, the percentage of RNs who intended to leave their nursing position increased with age, as did the percentage who intended to stop working in nursing after leaving that position.



| | | | Age of | nurses | | |
|--|---|---------------------------------|---------------------------------|---------------------------------|---------------------------------|---|
| | Less than 50 years old (percent) | 50-54 years old (percent) | 55-59 years old (percent) | 60-64 years old (percent) | 65-69 years old (percent) | 70 or more years old (percent) |
| Plans regarding current job | | | | | | |
| No plans to leave job | 53.3 | 64.0 | 59.6 | 42.5 | 33.8 | 34.5 |
| Undecided about plans | 16.8 | 13.9 | 13.4 | 14.2 | 17.6 | 19.0 |
| Have left job or plan to leave in | | | | | | |
| 12 months | 17.6 | 12.8 | 13.3 | 18.1 | 20.6 | 21.4 |
| Plan to leave in 1 to 3 years | 12.2 | 9.3 | 13.7 | 25.2 | 28.0 | 25.1 |
| Total that plans to leave within 3 | | | | | | |
| years | 29.8 | 22.1 | 27.0 | 43.3 | 48.7 | 46.5 |
| For those who plan to leave their job | | | | | | |
| Plan to remain in nursing work | 95.3 | 87.9 | 70.1 | 44.4 | 31.0 | 22.4 |
| Plan to leave nursing | 4.7 | 12.1 | 29.9 | 55.6 | 69.0 | 77.6 |

 Table 10-10.
 Intentions to change employers or leave the nursing profession, by age group

Source: 2008 National Sample Survey of Registered Nurses

10.5 Older Registered Nurses Not Working in Nursing

Among RNs under 70 years old in 2008, of those who were not working in nursing, a large percentage last worked in nursing 1 to 2 years previously (Table 10-11 and Appendix A, Table 48). However, these data may underestimate the rate at which older RNs stop working in nursing because these data do not include RNs whose licenses have lapsed.

| Table 10-11. | Length of time since last employed in nursing, for registered nurses not employed in |
|--------------|--|
| | nursing in 2008, by age group |

| | | | Age of | nurses | | |
|------------------|---|---------------------------------|---------------------------------|---------------------------------|---------------------------------|--------------------------------------|
| | Less than 50 years old (percent) | 50-54 years old (percent) | 55-59 years old (percent) | 60-64 years old (percent) | 65-69 years old (percent) | 70 or more years old (percent) |
| Less than 1 year | 13.3 | 12.0 | 12.5 | 11.8 | 9.6 | 6.5 |
| 1 to 2 years | 29.3 | 13.2 | 27.8 | 30.6 | 26.2 | 20.6 |
| 3 to 4 years | 14.9 | 13.6 | 14.9 | 15.1 | 14.9 | 11.5 |
| 5 to 9 years | 21.3 | 19.4 | 16.6 | 16.3 | 25.3 | 21.6 |
| 10 to 19 years | 18.9 | 27.8 | 16.2 | 17.9 | 16.8 | 29.5 |
| 20 or more years | - | 14.0 | 12.0 | 8.3 | 7.1 | 10.4 |

- Too few cases to report estimated percent (fewer than 30 respondents).

Source: 2008 National Sample Survey of Registered Nurses

With increasing age, retirement became an increasingly important reason for RNs in 2008 to not be working in nursing, particularly above age 60 (Table 10-12). Personal and family reasons, workplace issues, and career considerations were important factors in deciding to not work in nursing for a sizable proportion of RNs between 50 and 69 years old.

| | | | Age of | nurses | | |
|-----------------|---|---------------------------------|---------------------------------|---------------------------------|---------------------------------|--------------------------------------|
| | Less than 50 years old (percent) | 50-54 years old (percent) | 55-59 years old (percent) | 60-64 years old (percent) | 65-69 years old (percent) | 70 or more years old (percent) |
| Retirement | - | 9.1 | 25.2 | 63.2 | 78.7 | 86.9 |
| Career | 39.8 | 48.3 | 41.4 | 30.6 | 25.9 | 18.1 |
| Personal/family | 68.5 | 58.3 | 49.5 | 43.7 | 31.8 | 26.3 |
| Workplace | 41.0 | 51.8 | 45.0 | 36.3 | 31.2 | 14.1 |

Table 10-12. Reasons for not working in nursing, by age group

- Too few cases to report estimated percent (fewer than 30 respondents).

Source: 2008 National Sample Survey of Registered Nurses

Among older RNs not employed in nursing, only 3.3 percent were looking for work in nursing in 2008. Older RNs not working in nursing were less likely than their younger peers to report employment outside nursing (21.3 percent v. 29.4 percent, respectively). Older RNs working outside of nursing represented a small percentage of the nursing population, only 2.2 percent of the total nursing population. Of these, 52 percent were working in a health-related field (Table 10-13).

Table 10-13. Percent of registered nurses not employed in nursing who work in non-nursing positions, by age group

| | Under 50 years old | 50 years or older | Total |
|-------------------------------------|-----------------------|----------------------|-------|
| Percent working outside nursing | 29.4 | 21.3 | 23.8 |
| Of those employed outside nursing, | | | |
| percent whose job is health-related | 58.9 | 52.0 | 54.7 |

Source: 2008 National Sample Survey of Registered Nurses

Registered Nurses Nearing Retirement

This page deliberately blank.



Appendix A

Data Tables

| | | | | Number in | Estimated | Estimated | Number in | Estimated | Estimated |
|---|--------------|-----------|-----------|-------------|-------------|-------------|-------------|-------------|-------------|
| | | Total | Total | sample | number | percent | sample not | number not | percent not |
| Gender, racial/ethnic background | Total number | estimated | estimated | employed in |
| and age group | in sample | number | percent | nursing | nursing | nursing | nursing | nursing | nursing |
| Total | 33,179 | 3,063,162 | 100.0 | 28,554 | 2,596,599 | 100.0 | 4,625 | 466,564 | 100.0 |
| Gender | | | | | | | | | |
| Male | 2,332 | 202,491 | 6.6 | 2,123 | 183,719 | 7.1 | 209 | 18,772 | 4.0 |
| Female | 30,847 | 2,860,671 | 93.4 | 26,431 | 2,412,879 | 92.9 | 4,416 | 447,792 | 96.0 |
| Racial/ethnic background | | | | | | | | | |
| White (non-Hispanic) | 28,353 | 2,549,302 | 83.2 | 24,235 | 2,135,592 | 82.2 | 4,118 | 413,710 | 88.7 |
| Black/African American (non-Hispanic) | 1,688 | 165,352 | 5.4 | 1,510 | 146,459 | 5.6 | 178 | 18,894 | 4.0 |
| Asian (non-Hispanic) | 1,433 | 169,454 | 5.5 | 1,269 | 150,938 | 5.8 | 164 | 18,516 | 4.0 |
| Native Hawaiian/Pacific Islander (non-Hispanic) | 92 | 9,528 | 0.3 | 84 | 8,792 | 0.3 | 8 | 736 | 0.2 |
| American Indian/Alaskan Native (non-Hispanic) | 132 | 8,571 | 0.3 | 122 | 7,838 | 0.3 | 10 | 733 | 0.2 |
| Hispanic/Latino (any race) | 910 | 109,387 | 3.6 | 832 | 101,045 | 3.9 | 78 | 8,342 | 1.8 |
| Two or more races (non-Hispanic) | 571 | 51,568 | 1.7 | 502 | 45,935 | 1.8 | 69 | 5,633 | 1.2 |
| Age group | | | | | | | | | |
| Less than 25 | 730 | 79,641 | 2.6 | 699 | 77,041 | 3.0 | 31 | 2,600 | 0.6 |
| 25-29 | 1,966 | 208,543 | 6.8 | 1,868 | 198,734 | 7.7 | 98 | 9,809 | 2.1 |
| 30-34 | 2,677 | 281,999 | 9.2 | 2,467 | 257,472 | 9.9 | 210 | 24,528 | 5.3 |
| 35-39 | 3,149 | 331,972 | 10.8 | 2,876 | 302,520 | 11.7 | 273 | 29,452 | 6.3 |
| 40-44 | 3,623 | 350,101 | 11.4 | 3,293 | 314,368 | 12.1 | 330 | 35,734 | 7.7 |
| 45-49 | 4,763 | 441,831 | 14.4 | 4,333 | 397,552 | 15.3 | 430 | 44,279 | 9.5 |
| 50-54 | 6,035 | 495,985 | 16.2 | 5,393 | 433,848 | 16.7 | 642 | 62,137 | 13.3 |
| 55-59 | 4,858 | 396,967 | 13.0 | 4,209 | 337,918 | 13.0 | 649 | 59,049 | 12.7 |
| 60-64 | 2,995 | 249,533 | 8.1 | 2,234 | 176,932 | 6.8 | 761 | 72,601 | 15.6 |
| 65-69 | 1,509 | 140,263 | 4.6 | 809 | 69,595 | 2.7 | 700 | 70,667 | 15.1 |
| 70-74 | 562 | 53,670 | 1.8 | 251 | 19,752 | 0.8 | 311 | 33,918 | 7.3 |
| 75 and over | 312 | 32,657 | 1.1 | 122 | 10,867 | 0.4 | 190 | 21,790 | 4.7 |
| Average age | | 47.0 | | | 45.5 | | | 55.3 | |
| Median age | | 48 | | | 46 | | | 57 | |

| Table 2. Year of | graduation fro | om initial nursi | ng educatior | n and the average | ge age at gra | duation, by type | <u>e of initial</u> nu | rse education: | 2008 |
|---------------------------|----------------|------------------|--------------|-------------------|---------------|------------------|------------------------|----------------|----------------|
| | | | | | | Estimated | Estimated | Estimated | Estimated |
| | Total | Total | Total | Estimated | Estimated | number of | percent of | number of | percent of |
| Year of graduation from | number in | estimated | estimated | number of | percent of | associate | associate | bachelor's and | bachelor's and |
| initial nursing education | sample | number* | percent | diploma | diploma | degree | degree | higher | higher |
| Total | 33,179 | 3,063,162 | 100.0 | 623,548 | 100.0 | 1,388,884 | 100.0 | 1,044,306 | 100.0 |
| Year of graduation from | | | | | | | | | |
| initial nursing education | | | | | | | | | |
| 2005 or later | 2,971 | 313,622 | 10.2 | 9,570 | 1.5 | 177,401 | 12.8 | 126,105 | 12.1 |
| 2000-2004 | 3,695 | 363,607 | 11.9 | 11,451 | 1.8 | 207,504 | 14.9 | 143,822 | 13.8 |
| 1995-1999 | 4,325 | 419,229 | 13.7 | 26,065 | 4.2 | 230,015 | 16.6 | 162,545 | 15.6 |
| 1990-1994 | 4,212 | 383,674 | 12.5 | 34,058 | 5.5 | 211,647 | 15.2 | 137,947 | 13.2 |
| 1985-1989 | 3,545 | 329,612 | 10.8 | 43,658 | 7.0 | 169,912 | 12.2 | 114,945 | 11.0 |
| 1984 or earlier | 14,431 | 1,253,419 | 40.9 | 498,746 | 80.0 | 392,404 | 28.3 | 358,941 | 34.4 |
| | | | | | | Average age | | Average age | |
| Average age at graduation | | Average age | | Average age | | associate | | bachelor's and | |
| by year of graduation | | all graduates | | diploma | | degree | | higher | |
| 2005 or later | | 30.8 | | 32.9 | | 33.1 | | 27.5 | |
| 2000-2004 | | 29.9 | | 31.0 | | 31.7 | | 27.0 | |
| 1995-1999 | | 30.3 | | 29.3 | | 32.4 | | 27.6 | |
| 1990-1994 | | 30.2 | | 29.3 | | 32.5 | | 26.9 | |
| 1985-1989 | | 27.7 | | 25.8 | | 30.0 | | 25.1 | |
| 1984 or earlier | | 23.9 | | 22.2 | | 26.4 | | 23.6 | |

*Includes an estimated 13,325 RNs whose initial nursing education was in a master's degree program, an estimated 954 in a doctorate degree program, and an estimated 6,425 reported as "other." "Other" includes foreign credentials and other types of education which did not have a national equivalent category. NOTE: Estimated numbers may not equal total, and percents may not add to 100, because of rounding.

| Table 3. Financial resources used for tuition and fees for initial nursing education, by type of initial nursing education: 200 | Table 3. Financial resources used for tuition and fees for initial nursing | a education, by type of initial nursing education: 2008 |
|---|--|---|
|---|--|---|

| | Total number | Total estimated | Total estimated | Estimated number of | Estimated percent of | Estimated number of associate | Estimated percent of associate | Estimated number of | Estimated percent of | Estimated number of master's/ | Estimated percent of master's/ |
|--|--------------|--------------------|--------------------|------------------------|----------------------|-------------------------------------|--------------------------------------|------------------------|----------------------|-------------------------------------|--------------------------------------|
| Financial resources used | in sample | number* | percent | diploma | diploma | degree | degree | bachelor's | bachelor's | doctorate | doctorate |
| Total** | 33,179 | 3,063,162 | 100.0 | 623,548 | 100.0 | 1,388,884 | 100.0 | 1,030,026 | 100.0 | 14,280 | 100.0 |
| Earnings from healthcare-related employment | 10,447 | 923,996 | 30.5 | 124,633 | 20.0 | 527,801 | 38.0 | 277,945 | 27.0 | | |
| Earnings from nonhealthcare-related employment | 7,842 | 698,292 | 22.8 | 107,624 | 17.3 | 312,293 | 22.5 | 274,847 | 26.7 | | |
| Earnings from other household members | 8,034 | 743,647 | 24.3 | 120,029 | 19.2 | 387,095 | 27.9 | 233,037 | 22.6 | | |
| Personal household savings | 6,525 | 586,853 | 19.2 | 83,135 | 13.3 | 317,914 | 22.9 | 181,724 | 17.6 | | |
| Other family resources (parents or relatives) | 13,826 | 1,308,042 | 42.7 | 358,151 | 57.4 | 361,661 | 26.0 | 581,660 | 56.5 | 5,008 | 35.1 |
| Employer tuition, reimbursement plan | 3,363 | 310,977 | 10.2 | 31,031 | 5.0 | 205,121 | 14.8 | 74,526 | 7.2 | | |
| Federal traineeship, scholarship or grant | 5,531 | 487,972 | 15.9 | 53,340 | 8.6 | 238,828 | 17.2 | 193,252 | 18.8 | | |
| Federally assisted loan | 10,171 | 896,920 | 29.3 | 94,762 | 15.2 | 391,021 | 28.2 | 401,940 | 39.0 | 7,846 | 54.9 |
| Other type of loan | 3,768 | 331,478 | 10.8 | 54,955 | 8.8 | 117,506 | 8.5 | 154,799 | 15.0 | | |
| State/local government scholarship or grant | 5,368 | 498,171 | 16.3 | 61,382 | 9.8 | 232,454 | 16.7 | 202,101 | 19.6 | | |
| Nongovernment scholarship or grant | 4,753 | 427,598 | 14.0 | 88,511 | 14.2 | 146,772 | 10.6 | 189,436 | 18.4 | | |
| Other resources | 1.702 | 149.958 | 4.9 | 31.392 | 5.0 | 68.408 | 4.9 | 48.503 | 4.7 | | |

*Includes an estimated 2,591 RNs whose initial nursing education was reported as "other." "Other" includes foreign credentials and other types of education which did not have a national equivalent category. **Entries for financial resources used may add to more than the total because more than one resource may have been named by an individual registered nurse.

--Too few cases to report estimated number and estimated percent (fewer than 30 cases unweighted). NOTE: Estimated numbers may not equal total, and percents may not add to 100, because of rounding.

| | Total number in | Total estimated | Total estimated | Estimated number of | Estimated percent of | Estimated number of associate | Estimated percent of associate | Estimated number of bachelor's and | Estimated percent of bachelor's and |
|---|--------------------|-----------------|--------------------|---------------------|----------------------|-------------------------------------|--------------------------------------|--|---|
| Characteristics of nurses with prior degrees | sample | number* | percent | diploma | diploma | degree | degree | higher | higher |
| Total | 6,996 | 665,795 | 100.0 | 60,975 | 100.0 | 341,025 | 100.0 | 261,204 | 100.0 |
| Degree obtained before initial nursing education** | | | | | | | | | |
| Associate degree | 3,273 | 318,190 | 47.8 | 24,539 | 40.2 | 206,985 | 60.7 | 86,631 | 33.2 |
| Bachelor's | 3,775 | 351,524 | 52.8 | 36,392 | 59.7 | 142,789 | 41.9 | 169,964 | 65.1 |
| Master's | 471 | 43,124 | 6.5 | 7,075 | 11.6 | 15,462 | 4.5 | 20,255 | 7.8 |
| Doctorate | 50 | 4,335 | 0.7 | | | | | | |
| Other | 34 | 3,283 | 0.5 | | | | | | |
| Major field of study before initial nursing education | | | | | | | | | |
| Health-related field | 2,872 | 274,517 | 41.2 | 26,335 | 43.2 | 141,503 | 41.5 | 105,718 | 40.5 |
| Biological or physical science | 948 | 91,354 | 13.7 | 6,361 | 10.4 | 34,446 | 10.1 | 49,723 | 19.0 |
| Business or management | 639 | 60,656 | 9.1 | 6,242 | 10.2 | 34,087 | 10.0 | 20,185 | 7.7 |
| Education | 464 | 42,515 | 6.4 | 4,434 | 7.3 | 27,272 | 8.0 | 10,809 | 4.1 |
| Liberal arts, social science, or humanities | 1,758 | 165,936 | 24.9 | 15,177 | 24.9 | 85,707 | 25.1 | 64,446 | 24.7 |
| Law | 54 | 5,171 | 0.8 | | | | | | |
| Computer science | 45 | 5,157 | 0.8 | | | 3,661 | 1.1 | | |
| Social work | 106 | 10,541 | 1.6 | | | 6,110 | 1.8 | 3,586 | 1.4 |
| Other nonhealth-related field | 110 | 9,948 | 1.5 | | | 5,245 | 1.5 | 3,644 | 1.4 |
| Year of graduation from initial nursing education | | , | | | | , | | , | |
| 2005 or later | 1,064 | 116,449 | 17.5 | 4,509 | 7.4 | 63,253 | 18.5 | 48,141 | 18.4 |
| 2000-2004 | 1,180 | 115,381 | 17.3 | 3,885 | 6.4 | 64,044 | 18.8 | 46,622 | 17.8 |
| 1995-1999 | 1,279 | 121,005 | 18.2 | 7,722 | 12.7 | 64,741 | 19.0 | 48,162 | 18.4 |
| 1990-1994 | 1,037 | 94,449 | 14.2 | 6,070 | 10.0 | 50,468 | 14.8 | 37,911 | 14.5 |
| 1985-1989 | 683 | 61,954 | 9.3 | 7,232 | 11.9 | 31,797 | 9.3 | 22,549 | 8.6 |
| 1984 or earlier | 1,753 | 156,557 | 23.5 | 31,557 | 51.8 | 66,722 | 19.6 | 57,819 | 22.1 |
| | | , | | , | | Average age | | Average age | |
| | | Average age | | Average age | | associate | | bachelor's and | |
| Average age at graduation by year of graduation | | all graduates | | diploma | | degree | | higher | |
| 2005 or later | | 33.3 | | 32.3 | | 34.9 | | 31.3 | |
| 2000-2004 | | 32.4 | | 33.3 | | 33.7 | | 30.7 | |
| 1995-1999 | | 33.1 | | 32.6 | | 34.4 | | 31.3 | |
| 1990-1994 | | 32.3 | | 31.3 | | 34.1 | | 30.1 | |
| 1985-1989 | | 30.2 | | 29.1 | | 31.1 | | 29.1 | |
| 1984 or earlier | | 26.9 | | 25.1 | | 28.5 | | 25.9 | |

*Includes an estimated 2,591 RNs whose initial nursing education was reported as "other." "Other" includes foreign credentials and other types of education which did not have a national equivalent category.

**The number of degrees may add to more than the total because more than one degree may have been named by an individual registered nurse.

--Too few cases to report estimated number and estimated percent (fewer than 30 cases unweighted).

| | • | | 0 | | | Estimated | Estimated | Estimated | Estimated |
|--|-----------|-----------------|-----------|-----------|------------|-----------|------------|----------------|----------------|
| | Total | | Total | Estimated | Estimated | number of | percent of | number of | percent of |
| | number in | Total estimated | estimated | number of | percent of | associate | associate | bachelor's and | bachelor's and |
| Health occupation prior to initial nursing education** | sample | number* | percent | diploma | diploma | degree | degree | higher | higher |
| Total | 22,675 | 2,059,776 | 100.0 | 306,086 | 14.9 | 1,059,402 | 51.5 | 691,251 | 33.6 |
| Nursing aide/assistant | 15,729 | 1,412,214 | 100.0 | 235,855 | 16.7 | 648,961 | 46.0 | 525,713 | 37.3 |
| Home health aide/assistant | 1,947 | 177,573 | 100.0 | 15,246 | 8.6 | 100,273 | 56.6 | 61,681 | 34.8 |
| Licensed practical/vocational nurse | 5,017 | 436,754 | 100.0 | 32,548 | 7.5 | 349,080 | 80.0 | 54,651 | 12.5 |
| Emergency medical technician (EMT) or paramedic | 1,279 | 108,546 | 100.0 | 7,733 | 7.1 | 74,419 | 68.6 | 26,394 | 24.3 |
| Medical assistant | 1,277 | 123,180 | 100.0 | 13,117 | 10.7 | 76,524 | 62.4 | 33,009 | 26.9 |
| Dental assistant | 516 | 48,323 | 100.0 | 5,803 | 12.0 | 28,612 | 59.2 | 13,908 | 28.8 |
| Allied health technician/technologist | 1,806 | 166,202 | 100.0 | 13,659 | 8.2 | 98,014 | 59.1 | 54,242 | 32.7 |
| Manager in health care setting | 541 | 48,608 | 100.0 | 3,885 | 8.1 | 30,671 | 63.6 | 13,702 | 28.4 |
| Clerk in health care setting | 2,625 | 244,073 | 100.0 | 29,768 | 12.2 | 134,106 | 55.1 | 79,716 | 32.7 |
| Military medical corps | 554 | 44,772 | 100.0 | 4,126 | 9.2 | 27,489 | 61.6 | 13,041 | 29.2 |
| Medical doctor | 56 | 6,373 | 100.0 | | | | | | |
| Nurse midwife | 45 | 3,468 | 100.0 | | | | | | |
| Student nurse extern | 229 | 23,642 | 100.0 | 2,910 | 12.4 | 6,801 | 28.9 | 13,809 | 58.7 |
| Mental health/psychiatric | 229 | 23,642 | 100.0 | | | 3,833 | 55.8 | | |
| Health educator | 73 | 7,043 | 100.0 | | | | | | |
| Other health-related position | 204 | 19,419 | 100.0 | 3,226 | 16.6 | 9,916 | 51.1 | 6,277 | 32.3 |

Table 5. Health occupations prior to initial nursing education, by type of initial nursing education: 2008

*Includes an estimated 3,038 RNs for whom type of initial nursing education was not reported or reported as "other." "Other" includes foreign credentials and other types of education which did not have a national equivalent category. An estimated 1,003,386 RNs reported no health occupations prior to initial nursing education.

**Health occupation entries add to more than the total because more than one occupation may have been named by an individual registered nurse.

--Too few cases to report estimated number and estimated percent (fewer than 30 cases unweighted).

| | | | | | | Estimated | Estimated | Estimated | Estimated |
|---|-----------|---------------|-----------|-------------|------------|-------------|------------|----------------|----------------|
| | Total | Total* | Total | Estimated | Estimated | number of | percent of | number of | percent of |
| Year of graduation from initial nursing | number in | estimated | estimated | number of | percent of | associate | associate | bachelor's and | bachelor's and |
| education | sample | number | percent | diploma | diploma | degree | degree | higher | higher |
| Total | 6,216 | 548,397 | 100.0 | 44,182 | 100.0 | 416,872 | 100.0 | 86,608 | 100.0 |
| Year of graduation from | | | | | | | | | |
| initial nursing education | | | | | | | | | |
| 2005 or later | 640 | 57,537 | 10.5 | | | 51,727 | 12.4 | 5,105 | 5.9 |
| 2000-2004 | 824 | 71,881 | 13.1 | | | 61,682 | 14.8 | 8,427 | 9.7 |
| 1995-1999 | 955 | 85,436 | 15.6 | 3,409 | 7.7 | 69,920 | 16.8 | 12,108 | 14.0 |
| 1990-1994 | 1,060 | 88,783 | 16.2 | 5,149 | 11.7 | 69,763 | 16.7 | 13,872 | 16.0 |
| 1985-1989 | 973 | 86,083 | 15.7 | 5,977 | 13.5 | 68,547 | 16.4 | 11,492 | 13.3 |
| 1984 or earlier | 1,764 | 158,676 | 28.9 | 27,172 | 61.5 | 95,234 | 22.8 | 35,604 | 41.1 |
| | | | | | | Average age | | Average age | |
| Average age at graduation | | Average age | | Average age | | associate | | bachelor's and | |
| by year of graduation | | all graduates | | diploma | | degree | | higher | |
| 2005 or later | | 34.8 | | | | 35.1 | | 31.3 | |
| 2000-2004 | | 34.5 | | | | 34.7 | | 32.6 | |
| 1995-1999 | | 33.8 | | 33.7 | | 34.6 | | 29.6 | |
| 1990-1994 | | 33.9 | | 33.1 | | 34.8 | | 29.3 | |
| 1985-1989 | | 31.8 | | 29.9 | | 32.6 | | 27.9 | |
| 1984 or earlier | | 27.4 | | 25.1 | | 29.2 | | 24.1 | |

Table 6. Registered nurses who were ever licensed as practical/vocational nurse, by type of initial nursing education: 2008

*Includes an estimated 735 RNs for whom type of initial nursing education was reported as "other." "Other" includes foreign credentials and other types of education which did not have a national equivalent category.

--Too few cases to report estimated number and estimated percent (fewer than 30 cases unweighted).

| | | | | | Ту | pe of initial nu | ursing educat | ion | |
|----------------------------------|-----------|-----------|-----------|-----------|------------|------------------|---------------|------------|------------|
| | | | | | | Estimated | Estimated | Estimated | Estimated |
| Number of years since graduation | Total | Total | Total | Estimated | Estimated | number of | percent of | number of | percent of |
| from initial nursing education | number in | estimated | estimated | number of | percent of | associate | associate | bachelor's | bachelor's |
| program and State of location** | sample | number* | percent | diploma | diploma | degree | degree | and higher | and highe |
| Total | 31,679 | 2,892,927 | 100.0 | 571,079 | 100.0 | 1,373,510 | 100.0 | 945,770 | 100.0 |
| Located in same State | 19,622 | 2,016,215 | 69.7 | 360,291 | 63.1 | 1,037,450 | 75.5 | 616,944 | 65.2 |
| Located in different State | 12,057 | 876,712 | 30.3 | 210,788 | 36.9 | 336,060 | 24.5 | 328,825 | 34.8 |
| 5 years or less | 5,147 | 523,878 | 100.0 | 15,724 | 100.0 | 297,665 | 100.0 | 209,521 | 100.0 |
| Located in same State | 3,805 | 423,186 | 80.8 | 14,122 | 89.8 | 251,385 | 84.5 | 157,078 | 75.0 |
| Located in different State | 1,342 | 100,692 | 19.2 | | | 46,280 | 15.5 | 52,443 | 25.0 |
| 6-10 years | 3,721 | 365,118 | 100.0 | 12,983 | 100.0 | 216,437 | 100.0 | 135,442 | 100.0 |
| Located in same State | 2,543 | 276,720 | 75.8 | 10,214 | 78.7 | 174,105 | 80.4 | 92,225 | 68.1 |
| Located in different State | 1,178 | 88,398 | 24.2 | 2,769 | 21.3 | 42,332 | 19.6 | 43,217 | 31.9 |
| 11-15 years | 4,451 | 406,345 | 100.0 | 29,999 | 100.0 | 237,380 | 100.0 | 138,586 | 100.0 |
| Located in same State | 3,029 | 305,053 | 75.1 | 23,587 | 78.6 | 184,026 | 77.5 | 97,205 | 70.1 |
| Located in different State | 1,422 | 101,292 | 24.9 | 6,412 | 21.4 | 53,354 | 22.5 | 41,381 | 29.9 |
| 16-25 years | 7,002 | 634,587 | 100.0 | 82,203 | 100.0 | 341,889 | 100.0 | 210,429 | 100.0 |
| Located in same State | 4,396 | 444,303 | 70.0 | 61,056 | 74.3 | 246,658 | 72.1 | 136,570 | 64.9 |
| Located in different State | 2,606 | 190,284 | 30.0 | 21,147 | 25.7 | 95,231 | 27.9 | 73,859 | 35.1 |
| 26 years or more | 11,358 | 962,999 | 100.0 | 430,169 | 100.0 | 280,139 | 100.0 | 251,792 | 100.0 |
| Located in same State | 5,849 | 566,953 | 58.9 | 251,312 | 58.4 | 181,276 | 64.7 | 133,866 | 53.2 |
| Located in different State | 5,509 | 396,045 | 41.1 | 178,857 | 41.6 | 98,863 | 35.3 | 117,926 | 46.8 |

Table 7. State of location of registered nurses as of March 2008 and State of graduation, by type of initial nursing education and number of years since graduation: 2008*

*Excludes an estimated 165,539 RNs who graduated in a foreign country and an estimated 4,696 RNs who graduated in a U.S. territory (Guam, U.S. Virgin Islands or Puerto Rico).

**State of location is the State in which employed if employed in nursing, or State of residence if not employed in nursing.

--Too few cases to report estimated number and estimated percent (fewer than 30 cases unweighted).

| | | | •g••• | u e i g e i i u | <u></u> | | Number in | Estimated | Estimated | | | |
|-------------------------------|-----------|-----------|-----------|-----------------|------------|------------|------------|------------|------------|----------------|----------------|-----------------|
| | | | | Number in | Estimated | Estimated | sample | number | percent | Number in | Estimated | Estimated |
| | | | | sample | number | percent | initial | initial | initial | sample initial | number initial | percent initial |
| | Total | Total | Total | initial | initial | initial | education: | education: | education: | education: | education: | education: |
| Highest nursing or nursing- | number in | estimated | estimated | education: | education: | education: | associate | associate | associate | bachelor's and | bachelor's and | bachelor's and |
| related education | sample | number* | percent | diploma | diploma | diploma | degree | degree | degree | higher | higher | higher |
| Total | 33,179 | 3,063,162 | 100.0 | 6,901 | 623,548 | 100.0 | 14,849 | 1,388,884 | 100.0 | 11,362 | 1,044,306 | 100.0 |
| Diploma | 4,640 | 426,279 | 13.9 | 4,640 | 426,279 | 68.4 | 0 | | | 0 | | |
| Associate degree | 11,657 | 1,103,471 | 36.0 | 71 | 6,771 | 1.1 | 11,585 | 1,096,601 | 79.0 | 0 | | |
| Bachelor's in related field | 645 | 56,915 | 1.9 | 345 | 28,619 | 4.6 | 300 | 28,296 | 2.0 | 0 | | |
| Bachelor's in nursing | 11,487 | 1,068,250 | 34.9 | 907 | 81,689 | 13.1 | 1,848 | 167,887 | 12.1 | 8,727 | 818,343 | 78.4 |
| Master's in related field | 1,028 | 85,709 | 2.8 | 329 | 28,628 | 4.6 | 259 | 21,136 | 1.5 | 439 | 35,902 | 3.4 |
| Master's in nursing | 3,341 | 290,084 | 9.5 | 521 | 44,351 | 7.1 | 794 | 69,511 | 5.0 | 2,009 | 174,354 | 16.7 |
| Doctorate in related field | 184 | 15,229 | 0.5 | 55 | 4,669 | 0.7 | 37 | 2,827 | 0.2 | 92 | 7,732 | 0.7 |
| Doctorate in nursing | 154 | 13,140 | 0.4 | 33 | 2,541 | 0.4 | 26 | | | 95 | 7,975 | 0.8 |
| Total employed in nursing | 28,554 | 2,596,599 | 100.0 | 5,320 | 461,771 | 100.0 | 13,233 | 1,224,337 | 100.0 | 9,949 | 905,545 | 100.0 |
| Diploma | 3,554 | 315,071 | 12.1 | 3,554 | 315,071 | 68.2 | 0 | | | 0 | | |
| Associate degree | 10,398 | 973,815 | 37.5 | 47 | 3,917 | 0.8 | 10,350 | 969,798 | 79.2 | 0 | | |
| Bachelor's in related field | 478 | 39,757 | 1.5 | 249 | 19,175 | 4.2 | 229 | 20,582 | 1.7 | 0 | | |
| Bachelor's in nursing | 10,060 | 927,217 | 35.7 | 738 | 64,028 | 13.9 | 1,659 | 149,702 | 12.2 | 7,658 | 713,156 | 78.8 |
| Master's in related field | 789 | 62,797 | 2.4 | 231 | 18,446 | 4.0 | 217 | 17,435 | 1.4 | 340 | 26,874 | 3.0 |
| Master's in nursing | 2,971 | 253,341 | 9.8 | 432 | 35,553 | 7.7 | 732 | 63,380 | 5.2 | 1,796 | 153,232 | 16.9 |
| Doctorate in related field | 134 | 10,192 | 0.4 | 39 | 3,319 | 0.7 | 23 | | | 72 | 5,566 | 0.6 |
| Doctorate in nursing | 136 | 11,112 | 0.4 | 30 | 2,262 | 0.5 | 23 | | | 83 | 6,717 | 0.7 |
| Total not employed in nursing | 4,625 | 466,564 | 100.0 | 1,581 | 161,777 | 100.0 | 1,616 | 164,546 | 100.0 | 1,413 | 138,761 | 100.0 |
| Diploma | 1,086 | 111,208 | 23.8 | 1,086 | 111,208 | 68.7 | 0 | | | 0 | | |
| Associate degree | 1,259 | 129,656 | 27.8 | 24 | | | 1,235 | 126,803 | 77.1 | 0 | | |
| Bachelor's in related field | 167 | 17,158 | 3.7 | 96 | 9,444 | 5.8 | 71 | 7,714 | 4.7 | 0 | | |
| Bachelor's in nursing | 1,427 | 141,033 | 30.2 | 169 | 17,661 | 10.9 | 189 | 18,185 | 11.1 | 1,069 | 105,187 | 75.8 |
| Master's in related field | 239 | 22,912 | 4.9 | 98 | 10,182 | 6.3 | 42 | 3,702 | 2.2 | 99 | 9,028 | 6.5 |
| Master's in nursing | 370 | 36,744 | 7.9 | 89 | 8,798 | 5.4 | 62 | 6,132 | 3.7 | 213 | 21,122 | 15.2 |
| Doctorate in related field | 50 | 5,037 | 1.1 | 16 | | | 14 | | | 20 | | |
| Doctorate in nursing | 18 | | | 3 | | | 3 | | | 12 | | |

Table 8. Highest nursing or nursing-related education, by initial nursing education: 2008

*Includes an estimated 13,325 RNs whose initial nursing education was in a master's degree program, an estimated 954 in a doctorate degree program, and an estimated 6,425 in an other education program. Also includes an estimated 224 initial master's upgraded to doctorate in nursing, of which an estimated 64 are employed in nursing and an estimated 160 are not employed in nursing. Includes an estimated 4,084 RNs whose highest nursing or nursing-related education was not known. Of these, an estimated 3,297 were employed in nursing and an estimated 787 were not employed --Too few cases to report estimated number and estimated percent (fewer than 30 cases unweighted).

| | M | aster's degre | e** | Doctorate degree** | | | |
|--|------------------|---------------------|----------------------|---------------------|---------------------|----------------------|--|
| Primary focus of degree | Number in sample | Estimated number | Estimated percent | Number in sample | Estimated number | Estimated percent | |
| Total | 4,544 | 387,199 | 100.0 | 330 | 27,725 | 100.0 | |
| Clinical practice*** | 2,192 | 190,347 | 49.2 | 42 | 3,133 | 11.3 | |
| Administration/business/management | 890 | 76,918 | 19.9 | 33 | 2,885 | 10.4 | |
| Education | 565 | 48,743 | 12.6 | 86 | 6,498 | 23.4 | |
| Public health/community health | 306 | 22,782 | 5.9 | 12 | | | |
| Humanities, liberal arts, or social sciences | 33 | 2,852 | 0.7 | 13 | | | |
| Research | 10 | | | 67 | 6,129 | 22.1 | |
| Law | 5 | | | 29 | | | |
| Biological or physical sciences | 28 | | | 5 | | | |
| Informatics | 21 | | | 0 | | | |
| Social work | 14 | | | 0 | | | |
| Computer science | 1 | | | 0 | | | |
| Other health field | 434 | 35,145 | 9.1 | 39 | 3,518 | 12.7 | |
| Other nonhealth field | 45 | 4,276 | 1.1 | 4 | | | |

Table 9. Primary focus of post-nursing master's and doctorate degrees: 2008*

*Includes degrees in nursing or nursing-related areas.

**RNs may have reported multiple master's or doctorate degrees. In such cases, the master's/doctorate in nursing takes

precedence and is shown here. Only one master's or doctorate degree for each nurse is shown. ***Excludes RNs whose initial nursing education was at the graduate level: an estimated 13,325 in a master's degree program

and an estimated 954 in a doctorate degree program, all of which are in clinical practice.

--Too few cases to report estimated number and estimated percent (fewer than 30 cases unweighted).

| | | Current enrollment | |
|-------------------------------------|------------------|--------------------|----------------------|
| Employment and student status | Number in sample | Estimated number | Estimated percent |
| Total | 2,048 | 201,390 | 100.0 |
| Total employed in nursing full-time | 1,514 | 149,441 | 74.2 |
| Full-time student | 530 | 54,751 | 36.6 |
| Part-time student | 984 | 94,690 | 63.4 |
| Total employed in nursing part-time | 383 | 37,665 | 18.7 |
| Full-time student | 162 | 16,360 | 43.4 |
| Part-time student | 221 | 21,305 | 56.6 |
| Total not employed in nursing | 151 | 14,283 | 7.1 |
| Full-time student | 89 | 8,374 | 58.6 |
| Part-time student | 62 | 5,910 | 41.4 |
| Distance education | Number in sample | Estimated number | Estimated percent |
| Total | 2,048 | 201,390 | 100.0 |
| Total employed in nursing full-time | 1,514 | 149,441 | 74.2 |
| No distance-based | 344 | 38,279 | 25.6 |
| Up to 75 percent distance-based | 386 | 39,152 | 26.2 |
| More than 75 percent distance-based | 784 | 72,010 | 48.2 |
| Total employed in nursing part-time | 383 | 37,665 | 18.7 |
| No distance-based | 105 | 10,417 | 27.7 |
| Up to 75 percent distance-based | 140 | 14,640 | 38.9 |

Table 10. Current enrollment in nursing or nursing-relatededucation programs, by employment status and student status: 2008*

*Of the estimated 251,872 RNs pursuing formal educational programs, this table excludes the estimated 2,065 currently enrolled in post-master's certificate programs, an estimated 25,973 enrolled in other certificate programs, the estimated 22,445 whose certificate or academic programs were in a field not related to nursing, and those whose failed to indicate the type of formal program being pursued.

138

151

54

47

50

12,609

14,283

5,344

4,634

4,305

33.5

7.1

37.4

32.4

30.1

NOTE: Estimated numbers may not equal totals, and percents may not add to 100, because of rounding.

More than 75 percent distance-based

More than 75 percent distance-based

Up to 75 percent distance-based

Total not employed in nursing No distance-based

| | - | Estimated | |
|--|------------------|------------------|-----------------|
| ype of certification | Number in sample | number* | Estimated perce |
| Total certification | 12,209 | 1,094,838 | 35.7 |
| Administrator | 110 | 8,317 | 0.3 |
| Ambulatory | 73 | 5,103 | 0.2 |
| Anesthesiology | 70 | 6,230 | 0.2 |
| Cardiac rehabilitation | 50 | 4,569 | 0.1 |
| Case management | 434 | 31,106 | 1.0 |
| Community health | 46 | 5,024 | 0.2 |
| Critical care | 609 | 58,320 | 1.9 |
| Diabetes educator | 108 | 7,924 | 0.3 |
| Family practice | 61 | 5,277 | 0.2 |
| Gastroenterology | 51 | 5,411 | 0.2 |
| General practice | 43 | 3,686 | 0.1 |
| General surgery | 346 | 28,599 | 0.9 |
| Gerontology | 134 | 9,136 | 0.3 |
| Hospice and palliative care/home health | 183 | 15,058 | 0.5 |
| Infection control | 36 | 3,725 | 0.1 |
| Infusion therapy | 82 | 6,856 | 0.2 |
| Lactation consult | 109 | 9,164 | 0.3 |
| Legal nurse | 70 | 4,822 | 0.2 |
| Life support/resuscitation (BLS, ALS, BCLS, CPR, | - | , - | - |
| NRP, and others) | 10,545 | 956,472 | 31.2 |
| Maternal/neonate | 450 | 43,123 | 1.4 |
| Medical/surgical | 302 | 26,594 | 0.9 |
| Neonatal intensive care | 54 | 4,880 | 0.2 |
| Nephrology | 57 | 5,181 | 0.2 |
| Neuroscience | 68 | 6,764 | 0.2 |
| Occupational health | 74 | 4,868 | 0.2 |
| Oncology | 380 | 36,175 | 1.2 |
| Orthopedic | 61 | 6,110 | 0.2 |
| Pediatrics | 152 | 13,654 | 0.4 |
| Psychiatric/mental health/counselor | 250 | 19,126 | 0.6 |
| Quality care | 46 | 4,182 | 0.1 |
| Registered nurse first assistant | 30 | 2,920 | 0.1 |
| Rehabilitation | 150 | 13,054 | 0.4 |
| Research | 39 | 4,131 | 0.1 |
| School nurse/college health | | 6,121 | 0.2 |
| Trauma nursing/emergency medicine (TNCC, ATCN, | 15 | 0,121 | 0.2 |
| ATN, EMT, ENPC, and others) | 1 510 | 125 0.09 | 4.1 |
| Women's health | 1,512 31 | 125,008 3,080 | 4. 0.1 |
| Wound care | 101 | 3,080 7,590 | 0.2 |
| Other** | 420 | 36,061 | 1.2 |

Table 11. National nursing certifications for registered nurses: 2008

*An estimated 2,356 registered nurses who reported an other skill-based certification (e.g. CPR or basic life support) are not included here.

**Other includes urology, pain management, dialysis/hemodialysis, flight nurse, holistic, infectious disease, informatics, nurse educator, telemetry, addictions, radiology, transplant, and continuing education.

| Employment setting | Number in sample | Estimated number* | Estimated percent |
|---|------------------|----------------------|----------------------|
| Total | 28,554 | 2,596,599 | 100.0 |
| i otai | 20,004 | 2,000,000 | 100.0 |
| Hospital | 16,957 | 1,601,831 | 61.7 |
| Community hospital | 13,780 | 1,315,933 | 50.7 |
| Inpatient unit—community hospital | 10,645 | 1,027,190 | 39.6 |
| Nursing home unit—community hospital | 71 | 5,887 | 0.2 |
| Outpatient clinic community hospital owned | 1,481 | 137,910 | 5.3 |
| Outpatient clinic located at community hospital, not hospital owned | 169 | 16,213 | 0.6 |
| Other administrative or functional area—community hospital | 1,407 | 128,312 | 4.9 |
| Specialty hospital | 1,243 | 123,666 | 4.8 |
| Inpatient unit—specialty hospital | 942 | 94,565 | 3.6 |
| Outpatient clinic specialty hospital owned | 171 | 16,967 | 0.7 |
| Outpatient clinic located at specialty hospital, not hospital owned | 27 | | |
| Other administrative or functional area—specialty hospital | 103 | 9,853 | 0.4 |
| Long-term hospital | 275 | 24,678 | 1.0 |
| Inpatient unit—long-term hospital | 217 | 19,886 | 0.8 |
| Nursing home unit—long-term hospital | 24 | | |
| Other administrative or functional area—long term hospital | 34 | 2,518 | 0.1 |
| Psychiatric hospital | 370 | 31,707 | 1.2 |
| Inpatient unit-psychiatric hospital | 306 | 26,030 | 1.0 |
| Nursing home unit—psychiatric hospital | 5 | | |
| Outpatient clinic psychiatric hospital owned | 24 | | |
| Outpatient clinic located at psychiatric hospital, not hospital owned | 5 | | |
| Other administrative or functional area—psychiatric hospital | 30 | 2,892 | 0.1 |
| Federal Government hospital | 726 | 51,782 | 2.0 |
| Inpatient unit—Federal Government hospital | 373 | 27,371 | 1.1 |
| Nursing home unit—Federal Government hospital | 19 | | |
| Outpatient clinic at Federal Government hospital | 211 | 13,895 | 0.5 |
| Other administrative or functional area—Federal Government hospital | 123 | 8,694 | 0.3 |
| Institution, infirmary, or correctional facility hospital—all units | 71 | 7,723 | 0.3 |
| Other type of hospital | 479 | 45,288 | 1.7 |
| Inpatient unit—other hospital | 290 | 27,801 | 1.1 |
| Nursing home unit—other hospital | 6 | | |
| Outpatient clinic other hospital owned | 55 | 4,284 | 0.2 |
| Outpatient clinic located at other hospital, not hospital owned | 22 | | |
| Other administrative or functional area—other hospital | 104 | 10,323 | 0.4 |
| Nursing home/extended care facility | 1,636 | 135,514 | 5.2 |
| Nursing home/extended care (not in a hospital) | 1,320 | 110,519 | 4.3 |
| Facility for mentally retarded/developmentally disabled | 78 | 6,338 | 0.2 |
| Residential care/assisted living | 217 | 16,978 | 0.7 |
| Other type of extended care | 21 | | |
| Academic education program | 1,137 | 98,268 | 3.8 |
| Nursing aide and/or home health aide program | 66 | 5,972 | 0.2 |
| LPN/LVN program | 91 | 7,332 | 0.2 |
| Diploma nursing program | 124 | 10,737 | 0.0 |
| Associate degree nursing program | 317 | 29,192 | 1.1 |
| Bachelor's and higher degree nursing program | 400 | 32,513 | 1.3 |
| Associate degree nursing program and LPN/LVN program | 44 | 3,226 | 0.1 |
| Associate degree nursing program and BSN program | 35 | 2,756 | 0.1 |
| Other education program, not patient education | 59 | 6,498 | 0.3 |
| Footnotes at end of table. | 29 | 0,490 | 0.0 |

Footnotes at end of table.

| Employment setting | Number in sample | Estimated number* | Estimated percent |
|---|---------------------|----------------------|----------------------|
| | campie | | porcon |
| Home health setting | 1,828 | 165,697 | 6.4 |
| Visiting nurse service (VNS/VNA) | 329 | 28,822 | 1.1 |
| Home health service unit (hospital-based) | 247 | 19,752 | 0.8 |
| Home health agency (nonhospital-based) | 590 | 58,248 | 2.2 |
| Private duty in a home setting | 110 | 10,640 | 0.4 |
| Hospice | 421 | 37,528 | 1.4 |
| Other home health setting | 130 | 10,695 | 0.4 |
| Community/public health setting | 1,114 | 97,210 | 3.7 |
| State health or mental health agency | 228 | 16,467 | 0.6 |
| City or county health department | 353 | 34,637 | 1.3 |
| Correctional facility (nonhospital) | 138 | 14,214 | 0.5 |
| Community mental-health organization | 104 | 8,584 | 0.3 |
| Substance abuse center | 30 | 2,109 | 0.1 |
| Other community setting** | 261 | 21,199 | 0.8 |
| School health service | 913 | 84,418 | 3.3 |
| School system (K-12) | 790 | 73,697 | 2.8 |
| College or university | 79 | 6,792 | 0.3 |
| Other school health setting | 44 | 3,928 | 0.2 |
| Occupational health | 246 | 18,840 | 0.7 |
| Private industry | 129 | 10,456 | 0.4 |
| Government | 55 | 3,214 | 0.1 |
| Other occupational health setting | 60 | 4,764 | 0.2 |
| Ambulatory care setting (not hospital) | 2,904 | 270,556 | 10.4 |
| Medical/physician practice | 1,450 | 134,881 | 5.2 |
| Nurse practice | 55 | 4,968 | 0.2 |
| In-store or retail clinic | 26 | | |
| Community health center | 113 | 10,186 | 0.4 |
| Federal clinic (Military, VA, NIH, or IHS-supported) | 87 | 4,869 | 0.2 |
| Federally supported clinic (not in a community health center) | 31 | 2,706 | 0.1 |
| Hospital owned off-site clinics or surgery center | 253 | 24,824 | 1.0 |
| Ambulatory surgical center (not hospital-owned) | 388 | 39,463 | 1.5 |
| Urgent care | 78 | 6,245 | 0.2 |
| Dialysis center/clinic (not in a hospital) | 233 | 22,591 | 0.9 |
| Other ambulatory setting | 189 | 16,615 | 0.6 |
| Insurance claims/benefits/utilization review | 631 | 49,441 | 1.9 |
| Government insurer/benefits department: Federal, State or local | 83 | 7,472 | 0.3 |
| Insurance company or other private claims/benefits/utilization review | 547 | 41,936 | 1.6 |
| Other | 921 | 51,947 | 2.0 |
| Policy, planning, regulatory, or licensing agency | 85 | 6,197 | 0.2 |
| Consulting organization | 151 | 12,297 | 0.5 |
| Home-based self-employment | 55 | 4,727 | 0.2 |
| Telehealth, telenursing, or call center | 435 | 12,516 | 0.5 |
| Pharmaceutical/medical device/medical software | 92 | 8,062 | 0.3 |
| Adult/senior day care | 29 | | |
| Flight nurse/medical air transport | 47 | 3,766 | 0.1 |
| Other | 27 | | |
| Not known | 267 | 22,875 | 0.9 |

Table 12. Employment setting in principal nursing position: 2008 (continued)

*Includes an estimated 1,659 RNs for whom setting information was known, but subsetting information was not known. Such nurses are represented in the setting totals but not subsetting groups. As a result, subsetting counts and percents may not add to setting totals.

**Other community/public health setting includes blood banks and blood-related services.

--Too few cases to report estimated number and estimated percent (fewer than 30 cases unweighted).

Table 13. Employment setting in principal nursing position, by age group: 2008

| | | | | Age group | | | | | | | | | | |
|--|---------------------|--------------------|----------|-----------|---------|---------|---------|---------|---------|---------|---------|--------|--------|----------------|
| Employment setting | Number in sample | Estimated total | Under 25 | 25-29 | 30-34 | 35-39 | 40-44 | 45-49 | 50-54 | 55-59 | 60-64 | 65-69 | 70-74 | 75 and over |
| Total employed | 28,554 | 2,596,599 | 77,041 | 198,734 | 257,472 | 302,520 | 314,368 | 397,552 | 433,848 | 337,918 | 176,932 | 69,595 | 19,752 | 10,867 |
| Hospital | 16,957 | 1,601,831 | 68,635 | 163,651 | 187,954 | 204,803 | 201,511 | 241,578 | 239,917 | 177,617 | 81,114 | 25,885 | 6,565 | 2,602 |
| Nursing home/extended care facility | 1,636 | 135,514 | | 4,460 | 8,744 | 14,179 | 13,886 | 22,117 | 25,324 | 21,684 | 13,216 | 7,006 | 2,518 | |
| Academic education program | 1,137 | 98,268 | | 3,835 | 4,605 | 7,188 | 8,291 | 15,167 | 19,454 | 17,055 | 11,548 | 7,247 | | |
| Home health setting | 1,828 | 165,697 | | 4,512 | 10,896 | 19,817 | 20,810 | 26,453 | 30,880 | 25,869 | 13,545 | 7,960 | | |
| Public or community health setting | 1,114 | 97,210 | | 3,370 | 5,824 | 6,078 | 11,902 | 15,065 | 19,064 | 15,848 | 12,803 | 4,434 | | |
| School health service | 913 | 84,418 | | | 4,518 | 4,403 | 9,782 | 15,312 | 19,996 | 14,254 | 8,636 | 3,098 | | |
| Occupational health | 246 | 18,840 | | | | | | | 3,231 | 3,147 | 2,410 | | | |
| Ambulatory care setting (not hospital) | 2,904 | 270,556 | | 13,763 | 27,054 | 29,210 | 32,695 | 42,138 | 49,127 | 42,235 | 21,132 | 7,464 | 2,722 | |
| Insurance claims/benefits/utilization review | 631 | 49,441 | | | 3,339 | 6,665 | 5,949 | 7,733 | 11,116 | 7,263 | 4,242 | | | |
| Other | 921 | 51,947 | | | 2,148 | 5,767 | 5,289 | 6,206 | 10,712 | 10,220 | 5,911 | 2,570 | | |
| Not known | 267 | 22,875 | | | | | | 3,460 | 5,027 | 2,725 | 2,376 | | | |

--Too few cases to report estimated number (fewer than 30 cases unweighted). NOTE: Estimated numbers may not equal total because of rounding.

| | | _ | Highest | nursing or nursing-rela | ated education | al preparation |
|--|------------------------|----------------------------|-----------------------------------|--|--------------------------------------|--|
| Employment setting | Total number in sample | Total estimated number* | Estimated number of diploma | Estimated number of associate degree | Estimated number of bachelor's | Estimated number of master's/doctorate** |
| Total | 28,554 | 2,596,599 | 315,071 | 973,815 | 966,974 | 337,441 |
| Hospital | 16,957 | 1,601,831 | 167,942 | 626,119 | 646,695 | 159,090 |
| Nursing home/extended care facility | 1,636 | 135,514 | 22,417 | 71,426 | 33,576 | 7,840 |
| Academic education program | 1,137 | 98,268 | 10,897 | 23,397 | 24,221 | 39,753 |
| Home health setting | 1,828 | 165,697 | 24,043 | 70,783 | 57,734 | 12,766 |
| Public or community health setting | 1,114 | 97,210 | 13,085 | 32,781 | 35,374 | 15,739 |
| School health service | 913 | 84,418 | 11,739 | 17,321 | 38,748 | 16,610 |
| Occupational health | 246 | 18,840 | 3,358 | 6,517 | 5,590 | 3,376 |
| Ambulatory care setting (not hospital) | 2,904 | 270,556 | 43,943 | 87,704 | 76,610 | 61,979 |
| Insurance claims/benefits/utilization review | 631 | 49,441 | 7,052 | 14,646 | 21,301 | 6,361 |
| Other | 921 | 51,947 | 8,302 | 14,882 | 19,063 | 9,647 |
| Not known | 267 | 22,875 | 2,293 | 8,239 | 8,062 | 4,281 |

Table 14. Employment setting in principal nursing position, by highest nursing or nursing-related education: 2008

*Includes an estimated 3,297 RNs for whom highest educational preparation was not known.

**Across all employment settings, an estimated 316,137 registered nurses reported their highest education preparation as a master's degree, and an estimated

21,304 reported their highest education preparation as a doctorate degree.

NOTE: Estimated numbers may not equal totals because of rounding.

| General and specific job titles | Number in sample | Estimated number | Estimated percent |
|---|------------------|---------------------|-------------------|
| | | | |
| Total | 28,554 | 2,596,599 | 100.0 |
| Staff nurse | 18,176 | 1,711,271 | 65.9 |
| Staff nurse or direct care nurse | 14,421 | 1,369,847 | 52.8 |
| Charge nurse or team leader | 2,440 | 222,052 | 8.6 |
| School nurse | 737 | 68,588 | 2.6 |
| Public health nurse | 297 | 26,172 | 1.0 |
| Community health nurse | 281 | 24,612 | 0.9 |
| Management/administration | 3,642 | 322,790 | 12.4 |
| First-line management | 1,139 | 105,829 | 4.1 |
| Middle management/administration | 1,782 | 154,809 | 6.0 |
| Senior management/administration | 721 | 62,152 | 2.4 |
| Certified registered nurse anesthetist | 443 | 29,538 | 1.1 |
| Clinical nurse specialist | 258 | 22,070 | 0.8 |
| Nurse midwife | 77 | 6,455 | 0.2 |
| Nurse practitioner | 1,094 | 98,487 | 3.8 |
| Patient educator | 285 | 18,405 | 0.7 |
| Instruction | 1,084 | 94,946 | 3.7 |
| Staff educator or instructor in clinical setting | 428 | 40,772 | 1.6 |
| Staff development director | 78 | 7,356 | 0.3 |
| Instructor/lecturer | 320 | 26,624 | 1.0 |
| Professor | 258 | 20,194 | 0.8 |
| Patient coordinator | 1,729 | 140,060 | 5.4 |
| Patient care coordinator, case manager, discharge planner | 1,374 | 117,420 | 4.5 |
| Advice/triage nurse | 355 | 22,641 | 0.9 |
| Informatics nurse | 128 | 8,952 | 0.3 |
| Consultant | 290 | 23,115 | 0.9 |
| Researcher | 191 | 17,136 | 0.7 |
| Surveyor/auditor/regulator | 124 | 10,652 | 0.4 |
| Other | 861 | 76,323 | 2.9 |
| Quality improvement nurse, utilization review nurse | 503 | 44,523 | 1.7 |
| Infection control | 110 | 10,166 | 0.4 |
| Legal nurse | 52 | 4,848 | 0.2 |
| Other | 31 | 2,178 | 0.1 |
| No position title specified | 165 | 14,608 | 0.6 |
| Not known | 172 | 16,397 | 0.6 |

Table 15. Job title in principal nursing position: 2008

NOTE: Estimated numbers may not equal totals, and percents may not add to 100, because of rounding. Aggregated job titles as shown in this table are subsequently used to describe job title in a grouped manor.

| Job title | Total estimated number | Estimated number hospital setting | Estimated number nonhospital setting |
|--|------------------------------|---|---|
| Total | 2,596,599 | 1,601,831 | 994,768 |
| Staff nurse | 1,711,271 | 1,232,586 | 478,685 |
| Management/administration | 322,790 | 145,574 | 177,216 |
| Certified registered nurse anesthetist | 29,538 | 23,856 | 5,682 |
| Clinical nurse specialist | 22,070 | 13,943 | 8,127 |
| Nurse midwife | 6,455 | 2,682 | 3,773 |
| Nurse practitioner | 98,487 | 36,533 | 61,954 |
| Patient educator | 18,405 | 9,053 | 9,352 |
| Instruction | 94,946 | 28,857 | 66,089 |
| Patient coordinator | 140,060 | 48,605 | 91,456 |
| Informatics nurse | 8,952 | 6,105 | 2,847 |
| Consultant | 23,115 | 3,788 | 19,327 |
| Researcher | 17,136 | 8,625 | 8,510 |
| Surveyor/auditor/regulator | 10,652 | | 8,686 |
| Other* | 92,720 | 39,658 | 53,062 |

Table 16. Job title in principal nursing position, by hospital and nonhospital setting: 2008

*Other job title includes an estimated 16,397 RNs for whom job title was not known.

--Too few cases to report estimated number (fewer than 30 cases unweighted). NOTE: Estimated numbers may not equal totals because of rounding.

| Table 17. Employment setting in principal nursing position, |
|---|
| by selected job titles: 2008 |

| Employment setting | Total estimated number | Staff nurse | Management/ administration | Patient coordinator | Other job titles |
|--|------------------------------|-------------|-------------------------------|---------------------|------------------|
| Total | 2,596,599 | 1,711,271 | 322,790 | 140,060 | 422,477 |
| Hospital | 1,601,831 | 1,232,586 | 145,574 | 48,605 | 175,066 |
| Nursing home/extended care facility | 135,514 | 54,632 | 54,746 | 7,640 | 18,496 |
| Academic education program | 98,268 | 33,865 | 8,986 | | 54,112 |
| Home health setting | 165,697 | 82,935 | 34,513 | 32,317 | 15,933 |
| Public or community health setting | 97,210 | 58,820 | 15,671 | 4,094 | 18,625 |
| School health service | 84,418 | 74,593 | 3,544 | | 5,517 |
| Occupational health | 18,840 | 10,648 | 2,435 | | 4,311 |
| Ambulatory care setting (not hospital) | 270,556 | 146,842 | 36,574 | 12,801 | 74,339 |
| Insurance claims/benefits/utilization review | 49,441 | | 9,551 | 19,692 | 17,952 |
| Other* | 74,822 | 14,104 | 11,197 | 11,394 | 38,127 |

*Other employment setting includes an estimated 22,875 RNs for whom employment setting was not known. --Too few cases to report estimated number (fewer than 30 cases unweighted). NOTE: Estimated numbers may not equal totals because of rounding.

| | | | | | Hiç | ghest nursin | g or nursing | -related edu | cation prepa | ration | |
|--|-----------|-----------|-----------|-----------|------------|--------------|--------------|--------------|--------------|-------------|------------|
| | | | | | | Estimated | Estimated | | | Estimated | Estimated |
| Job title | | Total | Total | Estimated | Estimated | number of | percent of | Estimated | Estimated | number of | percent of |
| | Number in | estimated | estimated | number of | percent of | associate | associate | number of | percent of | master's/ | master's/ |
| | sample | number* | percent | diploma | diploma | degree | degree | bachelor's | bachelor's | doctorate** | doctorate |
| Total | 28,554 | 2,596,599 | 100.0 | 315,071 | 12.1 | 973,815 | 37.5 | 966,974 | 37.2 | 337,441 | 13.0 |
| Staff nurse | 18,176 | 1,711,271 | 100.0 | 218,399 | 12.8 | 739,456 | 43.2 | 685,446 | 40.1 | 65,198 | 3.8 |
| Management/administration | 3,642 | 322,790 | 100.0 | 38,302 | 11.9 | 108,445 | 33.6 | 112,286 | 34.8 | 63,347 | 19.6 |
| Certified registered nurse anesthetist | 443 | 29,538 | 100.0 | 3,078 | 10.4 | 1,221 | 4.1 | 4,652 | 15.7 | 20,588 | 69.7 |
| Clinical nurse specialist | 258 | 22,070 | 100.0 | | | 4,807 | 21.8 | 4,975 | 22.5 | 10,763 | 48.8 |
| Nurse midwife | 77 | 6,455 | 100.0 | | | | | | | 5,128 | 79.4 |
| Nurse practitioner | 1,094 | 98,487 | 100.0 | | | | | 4,241 | 4.3 | 88,280 | 89.6 |
| Patient educator | 285 | 18,405 | 100.0 | 2,339 | 12.7 | 5,607 | 30.5 | 7,858 | 42.7 | 2,601 | 14.1 |
| Instruction | 1,084 | 94,946 | 100.0 | 5,432 | 5.7 | 14,260 | 15.0 | 29,513 | 31.1 | 45,725 | 48.2 |
| Patient coordinator | 1,729 | 140,060 | 100.0 | 22,488 | 16.1 | 50,177 | 35.8 | 57,007 | 40.7 | 10,308 | 7.4 |
| Informatics nurse | 128 | 8,952 | 100.0 | | | 2,832 | 31.6 | 3,420 | 38.2 | | |
| Consultant | 290 | 23,115 | 100.0 | 3,603 | 15.6 | 4,620 | 20.0 | 8,783 | 38.0 | 6,110 | 26.4 |
| Researcher | 191 | 17,136 | 100.0 | | | 4,469 | 26.1 | 8,193 | 47.8 | 2,749 | 16.0 |
| Surveyor/auditor/regulator | 124 | 10,652 | 100.0 | | | 3,942 | 37.0 | 3,813 | 35.8 | | |
| Other | 861 | 76,323 | 100.0 | 10,928 | 14.3 | 24,075 | 31.5 | 29,924 | 39.2 | 11,376 | 14.9 |
| Not known | 172 | 16,397 | 100.0 | | | 6,310 | 38.5 | 6,337 | 38.6 | | |

Table 18. Job title in principal nursing position, by highest nursing or nursing-related education: 2008

*Includes an estimated 3,297 RNs for whom education preparation was not known.

**Across all job titles, an estimated 316,137 registered nurses reported their highest education preparation as a master's degree, and an estimated 21,304 reported their highest education preparation as a doctorate degree.

--Too few cases to report estimated number and estimated percent (fewer than 30 cases unweighted).

| Employment status | Number in sample | Total estimated number* | Total estimated percent | Average earnings: All nursing jobs (dollars) | Average earnings: Principal nursing job only (dollars) | Average earnings: Nonprincipal nursing jobs (dollars)** |
|---|---------------------|-------------------------------|-------------------------------|---|--|---|
| Total | 28,542 | 2,595,591 | 100.0 | 61,176 | 59,324 | 14,693 |
| Principal position: Full-time | 21,300 | 1,936,698 | 74.6 | 68,818 | 66,970 | 15,201 |
| Principal position: Part-time | 7,242 | 658,893 | 25.4 | 38,715 | 36,849 | 13,392 |
| Nurses with principal nursing position only | 24,822 | 2,268,332 | 100.0 | 59,197 | 59,197 | |
| Principal position: Full-time | 18,609 | 1,701,259 | 75.0 | 66,742 | 66,742 | |
| Principal position: Part-time | 6,213 | 567,073 | 25.0 | 36,562 | 36,562 | |
| Nurses with two or more nursing positions | 3,720 | 327,259 | 100.0 | 74,894 | 60,201 | 14,693 |
| Principal position: Full-time | 2,691 | 235,439 | 71.9 | 83,818 | 68,618 | 15,201 |
| Principal position: Part-time | 1,029 | 91,820 | 28.1 | 52,011 | 38,620 | 13,392 |

Table 19. Employment status and average total earnings for registered nurses withone and more than one nursing position: 2008

*Excludes an estimated 1,008 RNs for whom annual earnings amount was not known.

**Excludes RNs who do not hold a nonprincipal nursing job.

| | | | | Employment status | | | | | | | |
|-------------------------------------|-----------|-----------|---------------|-------------------|-----------|----------------|-----------|-----------|----------------|--|--|
| | Total | Total | Total average | Estimated | Estimated | Full-time | Estimated | Estimated | Part-time | | |
| Employment setting | estimated | estimated | annual hours | number full- | percent | average annual | number | percent | average annual | | |
| | number | percent | worked* | time | full-time | hours worked | part-time | part-time | hours worked | | |
| Total | 2,596,599 | 100.0 | 1,944 | 1,937,405 | 74.6 | 2,205 | 659,194 | 25.4 | 1,178 | | |
| Hospital | 1,601,831 | 100.0 | 1,969 | 1,206,920 | 75.3 | 2,202 | 394,911 | 24.7 | 1,257 | | |
| Nursing home/extended care facility | 135,514 | 100.0 | 2,107 | 107,676 | 79.5 | 2,373 | 27,838 | 20.5 | 1,080 | | |
| Academic education program | 98,268 | 100.0 | 1,852 | 72,654 | 73.9 | 2,146 | 25,615 | 26.1 | 1,020 | | |
| Home health setting | 165,697 | 100.0 | 2,008 | 117,404 | 70.9 | 2,344 | 48,293 | 29.1 | 1,190 | | |
| Community/public health setting | 97,210 | 100.0 | 1,958 | 76,779 | 79.0 | 2,209 | 20,432 | 21.0 | 1,017 | | |
| School health service | 84,418 | 100.0 | 1,443 | 61,500 | 72.9 | 1,701 | 22,918 | 27.1 | 750 | | |
| Occupational health | 18,840 | 100.0 | 1,808 | 13,111 | 69.6 | 2,191 | 5,729 | 30.4 | 933 | | |
| Ambulatory care setting | 270,556 | 100.0 | 1,860 | 186,570 | 69.0 | 2,199 | 83,986 | 31.0 | 1,106 | | |
| Insurance claims/benefits | 49,441 | 100.0 | 2,059 | 42,907 | 86.8 | 2,230 | 6,534 | 13.2 | 940 | | |
| Other | 51,947 | 100.0 | 1,950 | 36,292 | 69.9 | 2,335 | 15,655 | 30.1 | 1,058 | | |
| Not known | 22,875 | 100.0 | 1,782 | 15,593 | 68.2 | 2,141 | 7,282 | 31.8 | 1,015 | | |

Table 20. Average annual hours worked and employment setting in principal nursing position: 2008

*Hours worked includes regular hours of work, paid overtime, unpaid overtime, and on-call hours worked. Annual hours are estimated based on hours per typical work week multiplied by estimated months per year anticipated to be working per year in this principal nursing position.

| | | | | | Estimated | |
|----------------------------|-----------------|---------------|---------------|--------------|--------------|--------------|
| Employment setting | | | Estimated | Average paid | percent with | Average |
| | Total estimated | Average hours | percent with | overtime | mandatory | mandatory |
| | number | worked/week | paid overtime | hours/week | overtime | hours/week |
| Total | 1,937,405 | 42.9 | 27.5 | 7.5 | 7.0 | 6.6 |
| Hospital | 1,206,920 | 42.6 | 33.8 | 7.5 | 8.2 | 6.4 |
| Nursing home/extended care | 107,676 | 45.8 | 25.6 | 8.0 | 6.5 | 6.4 |
| Academic education | 72,654 | 44.0 | 18.7 | 8.4 | 4.8 | 8.9 |
| Home health | 117,404 | 45.3 | 21.4 | 7.8 | 7.9 | 8.1 |
| Public/community health** | 151,390 | 41.2 | 9.2 | 8.9 | 3.1 | 7.1 |
| Ambulatory care | 186,570 | 42.5 | 19.9 | 5.3 | 5.5 | 5.2 |
| Other*** | 79,199 | 44.0 | 7.1 | 9.9 | | |
| | | Average | Estimated | | Estimated | |
| Employment setting | Estimated | on-call | percent with | Average | percent with | Average paid |
| | percent with | hours/week | any standby | standby | paid standby | standby |
| | on-call hours* | worked* | hours | hours/week | hours | hours/week |
| Total | 14.5 | 13.3 | 13.0 | 25.1 | 9.3 | 20.1 |
| Hospital | 15.7 | 12.2 | 14.5 | 21.5 | 11.7 | 18.2 |
| Nursing home/extended care | 15.9 | 16.4 | 14.6 | 35.4 | 3.1 | 16.8 |
| Academic education | 10.1 | 13.7 | 7.4 | 21.5 | 5.1 | 16.9 |
| Home health | 30.1 | 13.2 | 25.5 | 32.2 | 17.9 | 27.9 |
| Public/community health** | 6.1 | 19.5 | 6.0 | 32.8 | 2.7 | 24.3 |
| Ambulatory care | 8.6 | 17.8 | 6.9 | 40.9 | 2.9 | 36.8 |
| Other*** | 6.1 | 20.6 | 3.6 | 28.9 | | |

Table 21. Average hours per week and employment setting for registered nurses working full-time in theirprincipal nursing position: 2008

*Reported on-call hours exclude hours that were standby only.

**Public/community health includes school and occupational health.

***Other includes insurance, benefits, and utilization review.

--Too few cases to report estimated percent and average (fewer than 30 cases unweighted).

| Employment setting | Total estimated number | Average hours worked/week | Estimated percent with paid overtime | Average paid overtime hours/week | Estimated percent with mandatory overtime | Average mandatory hours/week |
|---|---|---|---|--|--|---------------------------------------|
| Total | 659,194 | 24.1 | 15.1 | 5.4 | 3.4 | 4.8 |
| Hospital Nursing home/extended care | 394,911 27,838 | 25.2 22.4 | 19.9 13.5 | 5.5 6.4 | 3.7 | 4.5 |
| Academic education Home health | 25,615 48,293 | 22. 4 22.8 24.4 | 7.3 | 7.3 | | |
| Public/community health** Ambulatory care | 48,293 49,079 83,986 | 24.4 20.5 22.3 | 7.3 8.5 | 7.3 3.5 | | |
| Other*** | 22,189 | 20.5 | | | | |
| Employment setting | Estimated percent with on-call hours* | Average on-call hours/week worked* | Estimated percent with any standby hours | Average standby hours/week | Estimated percent with paid standby hours | Average paid standby hours/week |
| Total | 12.2 | 10.0 | 10.5 | 15.4 | 8.6 | 14.0 |
| Hospital Nursing home/extended care Academic education Home health Public/community health** Ambulatory care Other*** | 13.8 11.1 20.4 8.9 5.2 5.7 | 8.6 10.3 13.3 14.8 12.0 14.8 | 12.9 16.0 3.8 | 13.0 25.9 18.8 | 11.3 13.4 | 12.5 24.8 |

Table 22. Average hours per week and employment setting for registered nurses working part-time in theirprincipal nursing position: 2008

* Reported on-call hours exclude hours that were standby only.

**Public/community health includes school and occupational health.

***Other includes insurance, benefits, and utilization review.

--Too few cases to report estimated percent and average (fewer than 30 cases unweighted).

| | | | | | Clinical | specialty (estimated | number)* | | | |
|--|---------------------------|---------------|--------------|--------------|---------------------------|----------------------|--------------------------------|-----------------------------------|---------|--|
| Level of care/type of work | All clinical specialties* | Critical care | Cardiac care | Chronic care | Emergency/ trauma care | Gastrointestinal | General medical surgical | Gynecology (women's health) | Hospice | Infectious/ communicable disease |
| All levels of care/types of work* | 2,596,599 | 318,779 | 310,544 | 172,342 | 218,794 | 120,737 | 635,007 | 98,668 | 81,607 | 95,177 |
| General or specialty inpatient | 699,106 | 45,811 | 72,480 | 34,435 | 32,677 | 39,279 | 270,596 | 35,045 | 21,185 | 32,857 |
| Critical/intensive care | 345,396 | 262,073 | 93,113 | 12,358 | 39,105 | 15,819 | 41,509 | 5,039 | 4,666 | 13,167 |
| Step down/transitional/progressive/telemetry | 197,457 | 39,277 | 120,430 | 17,890 | 15,022 | 17,550 | 75,236 | 5,543 | 5,634 | 14,924 |
| Subacute care | 70,073 | 8,836 | 14,947 | 13,112 | 7,561 | 7,489 | 31,095 | 3,252 | 5,508 | 7,567 |
| Emergency | 199,333 | 30,833 | 22,847 | 9,778 | 160,539 | 10,767 | 29,673 | 6,268 | | 10,648 |
| Urgent care | 49,087 | 7,468 | 7,118 | 7,617 | 19,760 | 6,403 | 13,395 | 4,562 | | 8,163 |
| Rehabilitation | 86,348 | 4,539 | 12,219 | 18,436 | 3,351 | 4,449 | 25,668 | | 8,931 | 6,178 |
| Long-term care/nursing home | 152,523 | 2,781 | 7,337 | 56,306 | 2,333 | 3,415 | 34,291 | | 21,010 | 7,689 |
| Surgery | 377,112 | 34,817 | 45,260 | 11,709 | 24,484 | 41,726 | 228,224 | 31,796 | 4,773 | 11,354 |
| Ambulatory care | 358,936 | 7,074 | 29,968 | 29,951 | 14,107 | 32,113 | 45,833 | 33,020 | 3,491 | 16,696 |
| Home health | 155,677 | | 13,070 | 27,521 | | 4,947 | 42,282 | | 25,315 | 6,142 |
| Public/community health | 125,859 | | 2,352 | 8,986 | 7,527 | | 12,949 | 7,974 | | 22,465 |
| Education | 181,076 | 8,286 | 12,811 | 22,569 | 16,225 | 7,953 | 33,327 | 7,370 | 5,233 | 15,652 |
| Business/administration/review | 231,166 | 6,672 | 9,604 | 12,702 | 8,231 | 4,474 | 20,493 | 3,930 | 6,553 | 7,546 |
| Other*** | 78,180 | 7,738 | 12,759 | 4,861 | 7,205 | 4,085 | 7,604 | | | 5,000 |

Table 23. Level of care by type of clinical specialty in principal nursing position: 2008

| Level of care/type of work | Clinical specialty (estimated number)* | | | | | | | | | | |
|--|--|--------------|------------|----------|--------------|-------------------------------|---------------------------|--------------------|-----------------|----------------------|--|
| | Labor and delivery | Neurological | Obstetrics | Oncology | Primary care | Psychiatric/ mental health | Pulmonary/ respiratory | Renal/ dialysis | No patient care | Other specialty** | |
| All levels of care/types of work* | 94,929 | 85,238 | 132,911 | 119,894 | 220,678 | 133,791 | 120,880 | 89,372 | 314,916 | 237,380 | |
| General or specialty inpatient | 78,894 | 34,521 | 89,393 | 58,822 | 53,865 | 63,311 | 40,351 | 34,782 | 28,414 | 52,039 | |
| Critical/intensive care | 8,293 | 23,514 | 7,358 | 8,604 | 10,266 | 7,941 | 31,121 | 17,752 | 12,210 | 8,752 | |
| Step down/transitional/progressive/telemetry | | 19,314 | 3,416 | 11,083 | 10,358 | 7,205 | 32,712 | 17,508 | 9,162 | 7,352 | |
| Subacute care | | 4,651 | | 6,996 | 10,042 | 5,009 | 11,770 | 5,968 | 5,532 | 9,103 | |
| Emergency | 4,253 | 7,489 | 4,467 | 4,523 | 9,604 | 10,818 | 11,858 | 4,705 | 9,133 | 12,786 | |
| Urgent care | | 3,877 | | | 14,381 | 6,110 | 6,017 | | | 11,274 | |
| Rehabilitation | | 7,963 | | 3,197 | 12,667 | 7,029 | 9,795 | 2,777 | 10,024 | 11,077 | |
| Long-term care/nursing home | | 3,673 | | 2,522 | 20,597 | 12,340 | 8,716 | 2,419 | 23,610 | 18,764 | |
| Surgery | 12,586 | 19,469 | 14,121 | 13,681 | 11,743 | 4,377 | 18,219 | 10,202 | 12,457 | 54,145 | |
| Ambulatory care | 5,624 | 9,891 | 24,751 | 34,898 | 90,251 | 18,439 | 16,915 | 23,601 | 16,433 | 52,342 | |
| Home health | | 4,673 | | 5,981 | 20,589 | 5,364 | 12,887 | 3,132 | 23,590 | 21,954 | |
| Public/community health | | | 5,257 | | 28,719 | 14,033 | 3,674 | | 22,702 | 20,861 | |
| Education | 4,546 | 6,762 | 9,376 | 6,665 | 22,592 | 12,589 | 10,632 | 5,261 | 65,249 | 19,628 | |
| Business/administration/review | | 4,709 | 5,388 | 7,247 | 10,074 | 8,579 | 5,512 | 4,567 | 139,968 | 20,837 | |
| Other*** | | 4,780 | | 7,531 | 4,962 | 3,572 | 4,820 | | 26,566 | 13,506 | |

*The number of reported clinical specialties and levels of care will exceed the total number of employed RNs because each RN may report more than one clinical specialty and level of care. **Other clinical specialty includes dermatology, endocrinology, infusion, occupational health, ophthalmology, orthopedic, otolaryngology, plastic/cosmetic surgery, radiology, urology, no specific area, and other specialties.

***Other level of care/type of work includes ancillary care, research, informatics, flight nurse, and other.

| | | Patient population | | | | | | | | |
|---------------------------------|-----------------|--------------------|-----------|-----------|-----------------|----------------|--|--|--|--|
| | | | | | Pediatric/ | | | | | |
| Clinical specialty | Total estimated | | Geriatric | Adult and | maternal-child/ | Multiple ages/ | | | | |
| | number** | Adult only | only | geriatric | newborn*** | all ages | | | | |
| All clinical specialties* | 2,281,683 | 1,223,259 | 407,377 | 61,418 | 397,358 | 170,164 | | | | |
| Critical care | 318,779 | 182,167 | 31,640 | 12,476 | 75,745 | 16,509 | | | | |
| Cardiac care | 310,544 | 197,969 | 68,210 | 16,720 | 13,573 | 13,602 | | | | |
| Chronic care | 172,342 | 56,262 | 83,079 | 6,244 | 16,506 | 9,606 | | | | |
| Emergency/trauma care | 218,794 | 122,145 | 15,785 | 5,998 | 21,038 | 53,430 | | | | |
| Gastrointestinal | 120,737 | 76,833 | 16,242 | 5,652 | 10,988 | 10,978 | | | | |
| General medical surgical | 635,007 | 367,222 | 145,459 | 22,485 | 50,655 | 47,352 | | | | |
| Gynecology (women's health) | 98,668 | 57,842 | 2,471 | | 21,769 | 14,303 | | | | |
| Hospice | 81,607 | 25,347 | 45,650 | 3,817 | | 5,378 | | | | |
| Infectious/communicable disease | 95,177 | 43,240 | 18,254 | 4,142 | 18,047 | 10,814 | | | | |
| Labor and delivery | 94,929 | 35,445 | | | 52,843 | 5,588 | | | | |
| Neurological | 85,238 | 50,956 | 15,484 | 3,816 | 8,033 | 6,779 | | | | |
| Obstetrics | 132,911 | 47,486 | | | 74,158 | 10,302 | | | | |
| Oncology | 119,894 | 81,542 | 16,545 | 4,607 | 10,651 | 6,448 | | | | |
| Primary care | 220,678 | 78,865 | 42,866 | 4,700 | 69,600 | 22,836 | | | | |
| Psychiatric/mental health | 133,791 | 80,764 | 20,099 | 3,070 | 18,524 | 10,756 | | | | |
| Pulmonary/respiratory | 120,880 | 57,997 | 32,769 | 5,917 | 13,961 | 9,920 | | | | |
| Renal/dialysis | 89,372 | 56,941 | 17,488 | 5,118 | 3,973 | 5,042 | | | | |
| Other specialty**** | 237,380 | 127,917 | 48,036 | 6,015 | 30,522 | 23,316 | | | | |

Table 24. Clinical specialty by patient population in principal nursing position for registered nurses who provide patient care: 2008

*The number of reported clinical specialties will exceed the total number of employed RNs because each RN may report more than one specialty.

**Includes an estimated 22,106 RNs for whom patient population for principal nursing position was not known.

***An estimated 313,759 RNs provide patient care to a pediatric population, including newborn, neonatal, pediatric, and adolescent.

****Other clinical specialty includes dermatology, endocrinology, infusion, occupational health, ophthalmology, orthopedic, otolaryngology, plastic/cosmetic surgery, radiology, urology, no specific area, and other specialties.

| | | | Clinical specialty (estimated number)* | | | | | | | | | |
|--|---------------------------|---------------|--|--------------|---------------------------|--------------------|--------------------------------|-----------------------------------|---------|--|--|--|
| Level of care/type of work | All clinical specialties* | Critical care | Cardiac care | Chronic care | Emergency/ trauma care | Gastrointestinal | General medical surgical | Gynecology (women's health) | Hospice | Infectious/ communicable disease | | |
| All levels of care/types of work* | 1,479,993 | 301,271 | 262,175 | 60,237 | 186,451 | 85,806 | 459,315 | 60,739 | 22,459 | 50,647 | | |
| General or specialty inpatient | 572,291 | 42,636 | 66,210 | 25,378 | 25,178 | 35,634 | 244,796 | 32,590 | 15,141 | 25,679 | | |
| Critical/intensive care | 318,265 | 251,241 | 89,159 | 11,497 | 35,711 | 14,829 | 38,487 | 4,879 | 4,377 | 12,591 | | |
| Step down/transitional/progressive/telemetry | 179,632 | 35,935 | 115,564 | 16,631 | 14,375 | 16,375 | 71,457 | 5,315 | 5,194 | 14,097 | | |
| Subacute care | 41,713 | 7,478 | 12,782 | 6,956 | 5,517 | 6,578 | 23,401 | 3,047 | 2,479 | 5,960 | | |
| Emergency | 166,588 | 26,434 | 19,439 | 7,325 | 145,941 | 8,810 | 23,769 | 5,554 | | 6,791 | | |
| Urgent care | 21,966 | 6,652 | 5,867 | 3,789 | 11,771 | 4,263 | 7,947 | | | 4,151 | | |
| Rehabilitation | 41,650 | 3,423 | 9,356 | 6,078 | | 3,309 | 14,524 | | 3,218 | 3,897 | | |
| Surgery | 294,142 | 32,671 | 42,241 | 9,705 | 23,683 | 30,671 | 192,535 | 27,244 | 4,475 | 10,740 | | |
| Ambulatory care | 129,801 | 5,807 | 17,501 | 13,320 | 8,133 | 20,726 | 24,047 | 10,093 | | 7,686 | | |
| Education | 39,204 | 5,799 | 8,312 | 7,543 | 7,531 | 4,485 | 12,807 | 3,847 | | 5,004 | | |
| Other*** | 84,753 | 10,696 | 16,378 | 11,728 | 10,043 | 5,992 | 20,916 | | 4,237 | 7,918 | | |
| | | | | | Clinical specialt | y (estimated numbe | r)* | | | | | |

Table 25. Level of care by clinical specialty in principal nursing position for registered nurses who provide patient care in hospital settings: 2008

| Level of care/type of work | Labor and delivery | Neurological | Obstetrics | Oncology | Primary care | Psychiatric/ mental health | Pulmonary/ respiratory | Radiology | Renal/ dialysis | Other specialty** |
|--|--------------------|--------------|------------|----------|--------------|-------------------------------|---------------------------|-----------|-----------------|----------------------|
| All levels of care/types of work* | 86,271 | 67,151 | 102,300 | 89,611 | 57,861 | 79,638 | 85,602 | 19,215 | 57,094 | 78,318 |
| General or specialty inpatient | 75,343 | 31,675 | 84,274 | 55,569 | 23,975 | 52,888 | 36,823 | 6,812 | 31,083 | 30,812 |
| Critical/intensive care | 7,944 | 21,976 | 6,981 | 8,231 | 9,169 | 7,576 | 28,703 | 5,384 | 16,550 | |
| Step down/transitional/progressive/telemetry | | 18,346 | 3,416 | 10,525 | 9,565 | 6,635 | 30,887 | | 15,918 | 4,764 |
| Subacute care | | 3,931 | | 6,040 | 4,308 | 3,552 | 9,019 | | 4,388 | 2,860 |
| Emergency | 3,812 | 6,510 | 3,774 | 4,123 | 5,326 | 8,166 | 9,317 | | 4,276 | 4,717 |
| Urgent care | | | | | 3,960 | | 4,015 | | | |
| Rehabilitation | | 5,718 | | | 6,170 | 3,972 | 6,194 | | | 4,346 |
| Surgery | 11,784 | 17,531 | 12,856 | 12,914 | 9,856 | 3,781 | 17,251 | 6,355 | 9,410 | 29,576 |
| Ambulatory care | 4,064 | 5,280 | 9,050 | 19,968 | 19,207 | 6,503 | 10,108 | 5,618 | 6,807 | 14,437 |
| Education | 3,037 | 4,061 | 4,509 | 3,918 | 5,109 | 2,806 | 5,761 | | | 2,917 |
| Other*** | 2,919 | 4,960 | 4,738 | 8,506 | 7,206 | 8,220 | 7,350 | 7,582 | 3,523 | 8,477 |

*The number of reported clinical specialties and levels of care will exceed the total number of employed RNs in hospital settings because each RN may report more than one clinical speciality and level of care.

**Other clinical specialty includes dermatology, endocrinology, infusion, occupational health, ophthalmology, orthopedic, otolaryngology, plastic/cosmetic surgery, urology, no specific area, and other specialties.

***Other level of care/type of work includes ancillary care, long-term care, home health, public/community health, business/administration/review, research, informatics, flight nurse, and other.

| | | Patient population | | | | | | | | |
|---------------------------------|-----------------------------|--------------------|-------------------|---------------------|-------------------------------|----------------------------|--|--|--|--|
| | - | | | | Pediatric/ | | | | | |
| Clinical Specialty | Total estimated number** | Adult only | Geriatric only | Adult and geriatric | maternal-child/ newborn*** | Multiple ages/ all ages | | | | |
| All clinical specialties* | 1,479,993 | 882,787 | 187,144 | 46,718 | 245,428 | 107,460 | | | | |
| Critical care | 301,271 | 172,475 | 30,073 | 10,959 | 72,978 | 14,545 | | | | |
| Cardiac care | 262,175 | 176,040 | 47,921 | 14,738 | 12,618 | 10,626 | | | | |
| Chronic care | 60,237 | 27,863 | 18,180 | 3,274 | 5,038 | 5,287 | | | | |
| Emergency/trauma care | 186,451 | 108,475 | 13,671 | 5,265 | 10,682 | 48,073 | | | | |
| Gastrointestinal | 85,806 | 54,470 | 11,287 | 5,334 | 7,268 | 7,447 | | | | |
| General medical surgical | 459,315 | 292,976 | 82,231 | 17,632 | 32,292 | 33,946 | | | | |
| Gynecology (women's health) | 60,739 | 36,762 | | | 11,169 | 9,042 | | | | |
| Hospice | 22,459 | 9,240 | 8,917 | | | | | | | |
| Infectious/communicable disease | 50,647 | 27,949 | 10,920 | | 3,517 | 5,102 | | | | |
| Labor and delivery | 86,271 | 32,889 | | | 47,607 | 4,918 | | | | |
| Neurological | 67,151 | 42,764 | 10,244 | 3,256 | 5,735 | 5,152 | | | | |
| Obstetrics | 102,300 | 36,361 | | | 57,685 | 7,495 | | | | |
| Oncology | 89,611 | 61,976 | 9,500 | 3,796 | 9,479 | 4,860 | | | | |
| Primary care | 57,861 | 28,259 | 11,409 | | 9,578 | 5,536 | | | | |
| Psychiatric/mental health | 79,638 | 51,133 | 12,217 | | 7,235 | 6,571 | | | | |
| Pulmonary/respiratory | 85,602 | 45,944 | 19,694 | 5,127 | 8,181 | 6,383 | | | | |
| Radiology | 19,215 | 12,993 | | | | | | | | |
| Renal/dialysis | 57,094 | 35,127 | 11,115 | 4,012 | | 3,448 | | | | |
| Other specialty**** | 78,318 | 50,569 | 9,475 | | 6,726 | 8,658 | | | | |

Table 26. Clinical specialty by patient population in principal nursing position for registered nurses who provide patient care in hospital settings: 2008

*The number of reported clinical specialties will exceed the total number of employed RNs in hospital settings because each RN may report more than one specialty.

**Includes an estimated 10,455 RNs for whom patient population for principal nursing position was not known.

***An estimated 180,521 RNs provide patient care in a hospital setting to a pediatric population, including newborn, neonatal, pediatric, and adolescent.

****Other clinical specialty includes dermatology, endocrinology, infusion, occupational health, ophthalmology, orthopedic,

otolaryngology, plastic/cosmetic surgery, urology, no specific area, and other specialties.

| | Functional area | | | | | | | | | | | | |
|---|--|--|---|---------------------------------|---|--|---------------------------------|-------------------------------------|--------------------------------------|------------------------------------|------------------------------------|-------------------------------------|--|
| Percentage of time spent during usual work week | Number in sample patient care/charting | Estimated number patient care/charting | Estimated percent patient care/charting | | Estimated number non- nursing tasks | Estimated percent non- nursing tasks | Number in sample consultation | Estimated number consultation | Estimated percent consultation | Number in sample supervision | Estimated number supervision | Estimated percent supervision | |
| Total | 28,554 | 2,596,599 | 100.0 | 28,554 | 2,596,599 | 100.0 | 28,554 | 2,596,599 | 100.0 | 28,554 | 2,596,599 | 100.0 | |
| None | 3,857 | 327,192 | 12.6 | 9,740 | 841,860 | 32.4 | 11,153 | 1,009,762 | 38.9 | 15,686 | 1,420,385 | 54.7 | |
| 1-24 | 2,837 | 258,136 | 9.9 | 17,885 | 1,665,891 | 64.2 | 15,342 | 1,404,757 | 54.1 | 8,777 | 804,392 | 31.0 | |
| 25-49 | 2,357 | 213,727 | 8.2 | 814 | 76,440 | 2.9 | 1,332 | 118,526 | 4.6 | 1,902 | 174,450 | 6.7 | |
| 50-74 | 6,314 | 582,913 | 22.4 | 103 | 11,189 | 0.4 | 425 | 36,721 | 1.4 | 1,279 | 115,217 | 4.4 | |
| 75-100 | 13,189 | 1,214,630 | 46.8 | 12 | | | 302 | 26,832 | 1.0 | 910 | 82,155 | 3.2 | |
| | | | | | | Function | al area | | | | | | |
| Percentage of time spent during usual work week | Number in sample administration | Estimated number administration | Estimated percent administration | Number in sample research | Estimated number research | Estimated percent research | Number in sample teaching | Estimated number teaching | Estimated percent teaching | Number in sample other | Estimated number other | Estimated percent other | |
| Total | 28,554 | 2,596,599 | 100.0 | 28,554 | 2,596,599 | 100.0 | 28,554 | 2,596,599 | 100.0 | 28,554 | 2,596,599 | 100.0 | |
| None | 21,440 | 1,967,927 | 75.8 | 22,759 | 2,070,299 | 79.7 | 13,687 | 1,230,338 | 47.4 | 22,840 | 2,089,570 | 80.5 | |
| 1-24 | 5,427 | 482,419 | 18.6 | 5,312 | 484.723 | 18.7 | 12,824 | 1,187,142 | 45.7 | 4,000 | 364,273 | 14.0 | |
| 25-49 | 873 | 75,047 | 2.9 | 258 | 21,654 | 0.8 | 1,010 | 88,717 | 3.4 | 491 | 43,536 | 1.7 | |
| 50-74 | 448 | 39,132 | 1.5 | 122 | 10,901 | 0.4 | 462 | 41,637 | 1.6 | 462 | 38,537 | 1.5 | |
| 75-100 | 366 | 32,073 | 1.2 | 103 | 9,022 | 0.3 | 571 | 48,765 | 1.9 | 761 | 60,683 | 2.3 | |

Table 27. Percent of time in each functional area during usual work week in principal nursing position: 2008

--Too few cases to report estimated number and estimated percent (fewer than 30 cases unweighted). NOTE: Estimated numbers may not equal totals, and percents may not add to 100, because of rounding.

Table 28. Average percent of time in each functional area during usual work week in principal nursing position, by highest nursing or nursing-related education: 2008

| Highest education preparation | Average percent time in patient care/charting | Average percent time in non-nursing tasks | Average percent time in consultation | Average percent time in supervision |
|-------------------------------|---|--|--|---|
| Total | 58.1 | 6.1 | 7.5 | 10.2 |
| Diploma | 60.3 | 6.0 | 7.2 | 10.3 |
| Associate degree | 61.4 | 6.9 | 6.7 | 9.9 |
| Bachelor's | 58.1 | 6.5 | 7.9 | 10.2 |
| Master's | 48.2 | 3.0 | 8.9 | 10.7 |
| Doctorate | 18.1 | 1.6 | 8.8 | 8.7 |
| Not known* | 66.3 | 5.8 | 7.7 | 6.5 |
| Highest education preparation | Average percent | Average | Average | Average |
| | time in | percent time in | percent time in | percent time in |
| | administration | research | teaching | other |
| Total | 4.4 | 1.9 | 7.3 | 4.6 |
| Diploma | 3.3 | 1.5 | 5.6 | 6.0 |
| Associate degree | 3.2 | 1.7 | 6.1 | 4.1 |
| Bachelor's | 3.9 | 1.9 | 7.0 | 4.6 |
| Master's | 9.7 | 2.5 | 12.4 | 4.7 |
| Doctorate | 19.6 | 9.7 | 26.7 | 6.7 |
| Not known* | 9.0 | 1.2 | 2.4 | 1.2 |

*Not known represents registered nurses for whom education preparation information was not provided.

| Employment setting | Number in sample | Total estimated number | Total estimated percent | Estimated number employee of organization | | Estimated number employment agency | Estimated percent employment agency | Estimated number traveling nurse | Estimated percent traveling nurse | Estimated number self- employed/ per-diem | Estimated percent self- employed/ per-diem |
|--|---------------------|------------------------------|-------------------------------|--|------|---|--|---|--|--|---|
| Total | 28,554 | 2,596,599 | 100.0 | 2,412,775 | 92.9 | 36,863 | 1.4 | 26,182 | 1.0 | 120,778 | 4.7 |
| Hospital | 16,957 | 1,601,831 | 100.0 | 1,507,944 | 94.1 | 18,244 | 1.1 | 23,049 | 1.4 | 52,594 | 3.3 |
| Nursing home/extended care facility | 1,636 | 135,514 | 100.0 | 127,229 | 93.9 | | | | | 5,746 | 4.2 |
| Academic education program | 1,137 | 98,268 | 100.0 | 91,906 | 93.5 | | | | | 4,895 | 5.0 |
| Home health setting | 1,828 | 165,697 | 100.0 | 141,077 | 85.1 | 7,104 | 4.3 | | | 16,140 | 9.7 |
| Public or community health setting | 1,114 | 97,210 | 100.0 | 93,412 | 96.1 | | | | | 3,050 | 3.1 |
| School health service | 913 | 84,418 | 100.0 | 77,440 | 91.7 | | | | | 5,543 | 6.6 |
| Occupational health | 246 | 18,840 | 100.0 | 14,881 | 79.0 | 1,844 | 9.8 | | | | |
| Ambulatory care setting (not hospital) | 2,904 | 270,556 | 100.0 | 252,081 | 93.2 | | | | | 16,174 | 6.0 |
| Insurance claims/benefits/utilization review | 631 | 49,441 | 100.0 | 45,590 | 92.2 | | | | | | |
| Other | 921 | 51,947 | 100.0 | 40,404 | 77.8 | | | | | 11,137 | 21.4 |
| Not known | 267 | 22,875 | 100.0 | 20,811 | 91.0 | | | | | | |

Table 29. Employment setting in principal nursing position, by employment relationship: 2008

--Too few cases to report estimated number and estimated percent (fewer than 30 cases unweighted).

| Job title | Overall | Hospital | Nonhospital |
|--|------------|-----------|-------------|
| | average | setting | setting |
| | (dollars)* | (dollars) | (dollars) |
| Total | 66,973 | 69,079 | 63,491 |
| Staff nurse | 61,706 | 63,809 | 55,851 |
| Management/administration | 78,356 | 85,431 | 72,493 |
| Certified registered nurse anesthetist | 154,221 | 157,184 | 139,179 |
| Clinical nurse specialist | 72,856 | 76,386 | 65,177 |
| Nurse midwife | 82,111 | | 78,777 |
| Nurse practitioner | 85,025 | 89,362 | 82,018 |
| Patient educator | 59,421 | 62,759 | 56,438 |
| Instruction | 65,844 | 72,163 | 62,758 |
| Patient coordinator | 62,978 | 67,463 | 60,603 |
| Informatics nurse | 75,242 | 76,886 | 71,186 |
| Consultant | 76,473 | | 75,872 |
| Researcher | 67,491 | 68,278 | 66,695 |
| Surveyor/auditor/regulator | 65,009 | | 65,768 |
| Other** | 64,003 | 67,682 | 60,778 |

Table 30. Average annual earnings for registered nurses working full-timein their principal nursing position, by job title: 2008

* Excludes an estimated 467 RNs for whom annual earnings amount was not known.

**Other job title includes RNs for whom job title was not known.

--Too few cases to compute average salaries (fewer than 30 cases unweighted).

| Employment setting | Overall average (dollars)* | Staff nurse (dollars) | Management/ administration (dollars) | Patient coordinator (dollars) | Other job titles (dollars)*** |
|--|----------------------------------|--------------------------|--|-------------------------------------|----------------------------------|
| Total | 66,973 | 61,706 | 78,356 | 62,978 | 77,904 |
| Hospital | 69,079 | 63,809 | 85,431 | 67,463 | 87,162 |
| Nursing home/extended care facility | 61,507 | 54,239 | 65,351 | 58,387 | 67,177 |
| Academic education program | 66,701 | 66,542 | 82,387 | | 63,399 |
| Home health setting | 63,490 | 59,289 | 72,882 | 59,555 | 64,465 |
| Public or community health setting | 60,580 | 55,708 | 70,352 | 58,404 | 65,732 |
| School health service | 47,851 | 46,416 | 53,343 | | 64,723 |
| Occupational health | 65,947 | 62,086 | 72,716 | | |
| Ambulatory care setting (not hospital) | 65,976 | 55,657 | 76,412 | 53,811 | 80,105 |
| Insurance claims/benefits/utilization review | 67,897 | | 81,787 | 64,680 | 64,547 |
| Other** | 72,750 | 67,660 | 87,051 | 64,577 | 71,229 |

Table 31. Average annual earnings for registered nurses working full-time in their principal nursing position, by employment setting: 2008

*Excludes an estimated 467 RNs for whom annual earnings amount was not known.

**Other employment setting includes RNs for whom employment setting was not known.

***Other job titles include consultant, instruction, NP, NM, CNS, NA, researcher, informatics, surveyor/auditor/regulator, patient educator, and RNs with no job title.

--Too few cases to compute average salaries (fewer than 30 cases unweighted).

Table 32. Average annual earnings for registered nurses working full-time in their principal nursing position, by job title and highest nursing or nursing-related education: 2008

| | Overall | | Associate | | Master's/ |
|--|------------|-----------|-----------|------------|-------------|
| Job title | average | Diploma | degree | Bachelor's | doctorate |
| | (dollars)* | (dollars) | (dollars) | (dollars) | (dollars)** |
| Total | 66,973 | 65,349 | 60,890 | 66,316 | 87,363 |
| Staff nurse | 61,706 | 63,027 | 59,310 | 63,382 | 69,616 |
| Management/administration | 78,356 | 71,970 | 68,533 | 78,155 | 98,817 |
| Certified registered nurse anesthetist | 154,221 | | | 153,314 | 154,912 |
| Clinical nurse specialist | 72,856 | | 59,192 | 68,960 | 80,195 |
| Nurse midwife | 82,111 | | | | 81,824 |
| Nurse practitioner | 85,025 | | | 85,298 | 85,037 |
| Patient educator | 59,421 | 56,776 | 57,458 | 59,576 | |
| Instruction | 65,844 | 58,551 | 61,379 | 63,598 | 69,077 |
| Patient coordinator | 62,978 | 62,693 | 60,240 | 64,068 | 71,516 |
| Informatics nurse | 75,242 | | 69,950 | 78,046 | |
| Consultant | 76,473 | | 67,151 | 75,446 | 93,576 |
| Researcher | 67,491 | | 63,016 | 70,634 | 71,084 |
| Surveyor/auditor/regulator | 65,009 | | 64,308 | 62,675 | |
| Other*** | 64,003 | 60,756 | 57,287 | 64,628 | 79,618 |

* Excludes an estimated 467 RNs for whom annual earnings amount was not known.

Average annual earnings for registered nurses across all job titles who reported working full-time in their principal nursing position and holding a doctorate degree was an estimated \$86,173. *Other employment setting includes nurses for whom employment setting was not known.

--Too few cases to compute average salaries (fewer than 30 cases unweighted).

| | | | | Employment settin | g | |
|------------------------------------|--------------------------|------------------------|--------------------------------|----------------------------------|-------------|----------------------------|
| Level of job satisfaction | All settings* | Hospital | Nursing home/ extended care | Academic education program | Home health | Public/community health |
| Total | 2,573,723 | 1,601,831 | 135,514 | 98,268 | 165,697 | 97,210 |
| Extremely satisfied | 29.2 | 25.9 | 26.0 | 34.4 | 32.5 | 31.8 |
| Moderately satisfied | 51.8 | 53.9 | 48.5 | 52.2 | 50.3 | 49.5 |
| Neither satisfied nor dissatisfied | 7.8 | 8.3 | 8.6 | 6.0 | 6.9 | 7.4 |
| Moderately dissatisfied | 8.6 | 9.3 | 12.1 | 6.1 | 7.7 | 7.9 |
| Extremely dissatisfied | 2.5 | 2.5 | 4.8 | | 2.6 | |
| | | | Employment setting | | | _ |
| Level of job satisfaction | School health service | Occupational health | Ambulatory care | Insurance claims/benefits | Other | |
| Total | 84,418 | 18,840 | 270,556 | 49,441 | 51,947 | |
| Extremely satisfied | 39.7 | 37.8 | 38.5 | 38.4 | 39.3 | |
| Moderately satisfied | 48.2 | 49.8 | 47.0 | 42.4 | 42.9 | |
| Neither satisfied nor dissatisfied | 6.2 | | 5.8 | 10.1 | 8.6 | |
| Moderately dissatisfied | 4.6 | | 7.0 | 6.4 | 6.8 | |
| Extremely dissatisfied | | | 1.6 | | | |

Table 33. Job satisfaction, by employment setting in principal nursing position: 2008

*Includes an estimated 22,875 RNs for whom employment setting was not known.

--Too few cases to report estimated percent (fewer than 30 cases unweighted).

| | | Job title | | | | | | | | | |
|-------------------------------------|----------------------------|-------------|-------------------------------|-------------------------|------------------------------|---------------|-----------------------|--------|--|--|--|
| Level of ick action (normal) | _ | | | Certified registered | | | | | | | |
| Level of job satisfaction (percent) | Total estimated number* | Staff nurse | Management/ administration | nurse anesthetist | Clinical nurse specialist | Nurse midwife | Nurse practitioner | | | | |
| Total | 2,580,201 | 1,711,271 | 322,790 | 29,538 | 22,070 | 6,455 | 98,487 | | | | |
| Extremely satisfied | 29.3 | 25.3 | 36.5 | 52.0 | 37.8 | 41.1 | 40.9 | | | | |
| Moderately satisfied | 51.8 | 53.8 | 46.9 | 42.0 | 51.3 | 52.5 | 48.8 | | | | |
| Neither satisfied nor dissatisfied | 7.8 | 8.9 | 5.5 | | | | 3.8 | | | | |
| Moderately dissatisfied | 8.6 | 9.4 | 8.3 | | | | 5.0 | | | | |
| Extremely dissatisfied | 2.5 | 2.6 | 2.7 | | | | | | | | |
| | | | | | o title | | | | | | |
| Level of job satisfaction (percent) | Patient educator | Instruction | Patient coordinator | Informatics nurse | Consultant | Researcher | Surveyor/ auditor | Other | | | |
| Total | 18,405 | 94,946 | 140,060 | 8,952 | 23,115 | 17,136 | 10,652 | 76,323 | | | |
| Extremely satisfied | 48.5 | 37.8 | 31.3 | 49.7 | 41.3 | 29.7 | 39.2 | 37.1 | | | |
| Moderately satisfied | 41.0 | 52.6 | 50.9 | 30.7 | 44.4 | 55.6 | 42.4 | 43.7 | | | |
| Neither satisfied nor dissatisfied | | 2.9 | 7.5 | | | | | 8.7 | | | |
| Moderately dissatisfied | | 5.4 | 8.0 | | | | | 7.7 | | | |
| Extremely dissatisfied | | | 2.3 | | | | | | | | |

Table 34. Job satisfaction, by job title in principal nursing position: 2008

*Includes an estimated 16,397 RNs for whom job title was not known. --Too few cases to report estimated percent (fewer than 30 cases unweighted).

NOTE: Estimated numbers may not equal totals, and percents may not add to 100, because of rounding.

| | | | | Estimated | | |
|--------------|-------------------------------|--|---|---|--|---|
| Age group | Total estimated percent | Estimated percent extremely satisfied | Estimated percent moderately satisfied | percent neither satisfied nor dissatisfied | Estimated percent moderately dissatisfied | Estimated percent extremely dissatisfied |
| Total | 2,596,599 | 761,178 | 1,344,064 | 202,618 | 224,573 | 64,165 |
| Less than 25 | 100.0 | 18.6 | 59.9 | 9.9 | 9.8 | |
| 25-29 | 100.0 | 22.1 | 56.1 | 11.1 | 9.1 | 1.5 |
| 30-34 | 100.0 | 22.2 | 56.8 | 9.1 | 9.3 | 2.6 |
| 35-39 | 100.0 | 25.7 | 55.5 | 8.5 | 8.0 | 2.3 |
| 40-44 | 100.0 | 25.2 | 54.0 | 8.5 | 9.7 | 2.7 |
| 45-49 | 100.0 | 30.2 | 50.2 | 7.1 | 9.6 | 2.9 |
| 50-54 | 100.0 | 31.4 | 50.6 | 6.8 | 8.9 | 2.3 |
| 55-59 | 100.0 | 34.3 | 47.7 | 6.7 | 8.3 | 3.0 |
| 60-64 | 100.0 | 37.8 | 46.7 | 6.3 | 6.8 | 2.5 |
| 65-69 | 100.0 | 46.8 | 42.5 | 5.4 | 3.8 | |
| 70-74 | 100.0 | 57.2 | 29.6 | | | |
| 75 and over | 100.0 | 54.7 | 38.1 | | | |

Table 35. Job satisfaction of registered nurses employed in nursing, by age group: 2008

--Too few cases to report estimated percent (fewer than 30 cases unweighted). NOTE: Estimated numbers may not equal totals, and percents may not add to 100, because of rounding.

| | Highest education preparation | | | | | | | | | | |
|------------------------------------|-------------------------------|---------|------------------|------------|--------------------|--|--|--|--|--|--|
| Level of job satisfaction | Total* | Diploma | Associate degree | Bachelor's | Master's/doctorate | | | | | | |
| Total | 2,593,301 | 315,071 | 973,815 | 966,974 | 337,441 | | | | | | |
| Extremely satisfied | 29.3 | 31.4 | 26.1 | 28.4 | 39.2 | | | | | | |
| Moderately satisfied | 51.8 | 51.9 | 51.9 | 52.9 | 48.1 | | | | | | |
| Neither satisfied nor dissatisfied | 7.8 | 6.9 | 8.9 | 8.1 | 4.5 | | | | | | |
| Moderately dissatisfied | 8.7 | 7.3 | 10.1 | 8.5 | 6.1 | | | | | | |
| Extremely dissatisfied | 2.5 | 2.5 | 3.0 | 2.1 | 2.0 | | | | | | |

Table 36. Job satisfaction of registered nurses employed in nursing,by highest nursing or nursing-related education: 2008

*Includes an estimated 3,297 RNs for whom highest nursing or nursing-related education preparation was not known.

| | | | | | Er | nployment sta | atus in 2007 | | | | | |
|---|---------------------------|---------------------------------|-------------------------------|---|---|----------------------|---|-----------------------------|--------------------|-----------------------|------------------------------|--|
| Employment status in 2008 | Total number in sample | Total estimated number | Total estimated percent | Number in sample employed in nursing full- time | Estimated number employed in nursing full- time | | Number in sample employed in nursing part- time | | | | | Estimated percent not employed in nursing |
| Total | 33,179 | 3,063,162 | 100.0 | 21,323 | 1,916,670 | 62.6 | 6,823 | 619,490 | 20.2 | 5,033 | 527,002 | 17.2 |
| Employed in nursing full-time Employed in nursing part-time Not employed in nursing | 21,307 7,247 4,625 | 1,937,405 659,194 466,564 | 100.0 100.0 100.0 | 19,453 1,255 615 | 1,753,305 110,737 52,628 | 90.5 16.8 11.3 | 778 5,625 420 | 69,166 513,551 36,773 | 3.6 77.9 7.9 | 1,076 367 3,590 | 114,934 34,905 377,163 | 5.9 5.3 80.8 |

Table 37. Employment status in 2007 by employment status in 2008: 2008

| | | | | | Er | nployment set | ting in 200 | 7 | | | | |
|----------------------------|---------------------|-------------------------------|-------------------------------|----------|--------------------------------------|-----------------------|----------------|----------------------------------|-----------------|----------|-----------------|-----------------|
| Employment setting in 2008 | Number in sample | Total estimated number* | Total estimated percent | Hospital | Nursing home/ extended care | Academic education | Home health | Public/ community health** | Ambulatory care | Other*** | Not employed | New graduate |
| Total | 33,179 | 3,063,162 | 100.0 | 58.9 | 5.2 | 3.5 | 5.9 | 7.2 | 9.8 | 3.7 | 2.8 | 2.9 |
| Hospital | 16,957 | 1,601,831 | 100.0 | 90.8 | 0.6 | 0.2 | 0.4 | 0.3 | 0.7 | 0.2 | 2.4 | 4.3 |
| Nursing home/extended care | 1,636 | 135,514 | 100.0 | 5.5 | 85.3 | | 1.5 | | | | 3.1 | |
| Academic education | 1,137 | 98,268 | 100.0 | 4.9 | | 85.8 | | | | | 4.4 | |
| Home health | 1,828 | 165,697 | 100.0 | 7.9 | | | 82.5 | | | | 3.4 | |
| Public/community health** | 2,273 | 200,469 | 100.0 | 4.4 | | | | 86.4 | 2.2 | | 4.3 | |
| Ambulatory care | 2,904 | 270,556 | 100.0 | 8.2 | | | | 0.7 | 85.5 | | 2.5 | |
| Other*** | 1,552 | 101,388 | 100.0 | 6.4 | | | | | | 85.5 | 4.2 | |
| Not employed in nursing | 4,625 | 466,564 | 100.0 | 9.5 | 1.4 | 1.6 | 2.3 | 2.2 | 1.2 | 1.0 | 79.9 | |

Table 38. Employment setting in 2007 by employment setting in 2008: 2008

*Includes an estimated 22,875 RNs whose employment setting was not known for 2008 and an estimated 16,295 RNs for whom employment setting was not known for 2007.

**Public/community health setting includes school health and occupational health.

***Other includes insurance, benefits, and utilization review.

--Too few cases to report estimated percent (fewer than 30 cases unweighted).

| Reasons for employer or position change | Number in sample | Estimated number | Estimated |
|--|------------------|---------------------|-----------|
| Reasons for employer of position change | Sample | number | percent |
| Total | 5,670 | 473,248 | 100.0 |
| Relocated to different area | 1059 | 70,427 | 14.9 |
| Personal/family reasons | 882 | 72,756 | 15.4 |
| Disability | 89 | 6,265 | 1.3 |
| Illness | 103 | 7,837 | 1.7 |
| Interested in another position/job | 1618 | 139,413 | 29.5 |
| Opportunity to do the nursing that I like | 634 | 54,987 | 11.6 |
| Went back to school | 211 | 20,524 | 4.3 |
| Burnout | 881 | 78,617 | 16.6 |
| Stressful work environment | 1599 | 140,013 | 29.6 |
| Lack of advancement opportunities | 696 | 59,613 | 12.6 |
| _ack of collaboration/communication | 696 | 61,934 | 13.1 |
| Lack of good management/leadership | 1511 | 131,686 | 27.8 |
| Career advancement/promotion | 862 | 83,094 | 17.6 |
| Inadequate staffing | 1029 | 94,941 | 20.1 |
| Interpersonal differences with colleagues | 579 | 49,070 | 10.4 |
| Physical demands of job | 507 | 44,958 | 9.5 |
| Pay/benefits better | 1057 | 90,955 | 19.2 |
| Scheduling/inconvenient hours/too many hours | 909 | 82,614 | 17.5 |
| Reorganization that shifted positions | 244 | 21,029 | 4.4 |
| Laid off/downsizing/contract | 295 | 23,019 | 4.9 |
| Sign-on bonus offered | 98 | 8,947 | 1.9 |
| Retired | 238 | 20,814 | 4.4 |
| Other | 15 | | |
| Summary of reasons for employment change | | | |
| Personal/family reasons | 1877 | 139,360 | 29.4 |
| Personal career reasons | 2,067 | 177,651 | 37.5 |
| Workplace reasons | 3,963 | 346,179 | 73.1 |

Table 39. Reasons for employer or position change among registered nursesemployed in nursing in 2007 and 2008: 2008

--Too few cases to report estimated number and estimated percent (fewer than 30 cases unweighted). NOTE: Estimated numbers and percents may not add up to totals because registered nurses may have answered to more than one reason.

| | Number in | Estimated | Estimated |
|---|-----------|-----------|-----------|
| Reasons for other occupation | sample* | number | percent |
| Total | 1,112 | 111,062 | 100.0 |
| Retired | 94 | 8,966 | 8.1 |
| Taking care of home and family | 237 | 23,422 | 21.1 |
| Burnout | 175 | 17,122 | 15.4 |
| Stressful work environment | 251 | 24,694 | 22.2 |
| Scheduling/inconvenient hours/too many hours | 293 | 29,918 | 26.9 |
| Physical demands of job | 158 | 15,246 | 13.7 |
| Disability/Illness | 45 | 5,091 | 4.6 |
| Inadequate staffing | 221 | 22,726 | 20.5 |
| Salaries too low/better pay elsewhere | 248 | 24,291 | 21.9 |
| Skills are out of date | 141 | 13,956 | 12.6 |
| Liability concerns | 94 | 9,265 | 8.3 |
| Lack of collaboration/communication | 111 | 11,348 | 10.2 |
| Inability to practice nursing on professional level | 60 | 6,318 | 5.7 |
| Lack of advancement opportunities | 96 | 9,186 | 8.3 |
| Lack of good management/leadership | 147 | 14,747 | 13.3 |
| Career change | 526 | 53,201 | 47.9 |
| Difficult to find nursing position | 33 | 3,924 | 3.5 |
| Went back to school | 57 | 5,369 | 4.8 |
| Other** | 50 | 4,442 | 4.0 |
| Summary of reasons for other occupations | | | |
| Personal/family reasons | 286 | 28,783 | 25.9 |
| Personal career reasons | 704 | 71,722 | 64.6 |
| Workplace reasons | 594 | 59,418 | 53.5 |
| Retirement reasons | 94 | 8,966 | 8.1 |

Table 40. Reasons for registered nurses to have occupationother than nursing: 2008

*The number in sample excludes RNs who are employed in nursing, but have additional employment outside of nursing.

**Other includes travel, volunteering in nursing, and other reasons.

NOTE: Estimated numbers and percents may not add up to totals because registered nurses may have answered to more than one reason.

| Type of non-nursing | Number in sample | Estimated | Estimated |
|-------------------------------|------------------|-----------|-----------|
| employment | | number* | percent |
| Total | 1,112 | 111,062 | 100.0 |
| Health-related occupation | 605 | 60,716 | 54.7 |
| Full-time | 434 | 44,770 | 40.3 |
| Part-time | 171 | 15,947 | 14.4 |
| Not health-related occupation | 507 | 50,346 | 45.3 |
| Full-time | 252 | 25,488 | 22.9 |
| Part-time | 255 | 24,858 | 22.4 |

Table 41. Type of employment of registered nurses who are innon-nursing occupations and not employed in nursing:2008

*Includes an estimated 5,073 RNs who are employed in non-nursing and are actively seeking nursing employment. Excludes an estimated 97,365 RNs who are employed in nursing and have other employment in non-nursing.

| | | | | Ту | pe of position | outside of nurs | ursing | |
|---|-----------|--------------------|--------------------|-------------------|--------------------|-----------------------|------------------------|--|
| | | | | Estimated | Estimated | Estimated | Estimated | |
| Type of non-nursing employment | Number in | Total estimated | Total estimated | number health- | percent health- | number not health- | percent not health- | |
| | sample* | number | percent | related | related | related | related | |
| Total | 1,112 | 111,062 | 100.0 | 60,716 | 100.0 | 50,346 | 100.0 | |
| Health-related services, outside nursing | 354 | 34,891 | 31.4 | 31,892 | 52.5 | | | |
| Retail sales/services and food services | 160 | 16,040 | 14.4 | | | 14,587 | 29.0 | |
| Education (elementary and secondary) | 120 | 11,517 | 10.4 | 2,789 | 4.6 | 8,728 | 17.3 | |
| Administration/HR/consulting organization/legal | 92 | 9,767 | 8.8 | 5,960 | 9.8 | 3,806 | 7.6 | |
| Pharmaceutical/biotechnology/medical equipment | 89 | 9,955 | 9.0 | 9,563 | 15.8 | | | |
| Financial/accounting/insurance services | 68 | 6,931 | 6.2 | | | 4,794 | 9.5 | |
| Computer services | 49 | 4,768 | 4.3 | 3,659 | 6.0 | | | |
| Government | 38 | 3,300 | 3.0 | | | | | |
| Real estate | 33 | 3,241 | 2.9 | | | 3,241 | 6.4 | |
| Other** | 109 | 10,652 | 9.6 | | | 8,991 | 17.9 | |

Table 42. Detailed type of employment of registered nurses who are innon-nursing occupations and not employed in nursing: 2008

*The number in sample excludes RNs who are employed in nursing, but have other employment outside of nursing.

**Other includes agriculture/farming/ranching, fine arts/performing arts, emergency response, fitness/sport, religious organizations, etc.

--Too few cases to report estimated number and estimated percent (fewer than 30 cases unweighted).

| | | | | Ту | pe of position | outside of nurs | ing |
|-------------------------------------|-----------|-----------|-----------|-----------|----------------|-----------------|-------------|
| | | | - | Estimated | Estimated | Estimated | Estimated |
| lob title of non purging accuration | | Total | Total | number | percent | number not | percent not |
| Job title of non-nursing occupation | Number in | estimated | estimated | health- | health- | health- | health- |
| | sample* | number | percent | related | related | related | related |
| Total | 1,112 | 111,062 | 100.0 | 60,716 | 100.0 | 50,346 | 100.0 |
| Management | 247 | 25,892 | 23.3 | 19,594 | 32.3 | 6,299 | 12.5 |
| Business owner/proprietor | 188 | 17,914 | 16.1 | 4,807 | 7.9 | 13,107 | 26.0 |
| Administrative/clerical support | 171 | 17,410 | 15.7 | 9,145 | 15.1 | 8,265 | 16.4 |
| Instructor/professor | 130 | 12,527 | 11.3 | 4,899 | 8.1 | 7,629 | 15.2 |
| Consultant | 116 | 11,761 | 10.6 | 8,839 | 14.6 | 2,921 | 5.8 |
| Sales | 111 | 11,695 | 10.5 | 4,622 | 7.6 | 7,073 | 14.0 |
| Health practitioner | 54 | 5,094 | 4.6 | 4,254 | 7.0 | | |
| Other** | 95 | 8,770 | 7.9 | 4,557 | 7.5 | 4,212 | 8.4 |

Table 43. Job title of registered nurses who are in non-nursing occupationsand not employed in nursing: 2008

*The number in sample excludes RNs who are employed in nursing, but have other employment outside of nursing.

**Other includes such job titles as emergency response, clinical research, construction, clergy, chef, IT, etc.

--Too few cases to report estimated number and estimated percent (fewer than 30 cases unweighted).

| Advanced practice preparation combinations | Total number in sample | Total estimated number* | Total estimated percent | Estimated number employed in nursing | Estimated percent employed in nursing | Estimated number not employed in nursing | Estimated percent not employed in nursing |
|--|------------------------------|-------------------------------|-------------------------------|---|--|---|--|
| Total nurses with advanced practice preparation | 2,897 | 250,527 | 100.0 | 220,494 | 100.0 | 30,033 | 100.0 |
| Nurse anesthetists only | 489 | 33,950 | 13.6 | 30,997 | 14.1 | 2,953 | 9.8 |
| Clinical nurse specialists only | 509 | 42,414 | 16.9 | 34,987 | 15.9 | 7,427 | 24.7 |
| Nurse practitioners only | 1,495 | 138,558 | 55.3 | 123,759 | 56.1 | 14,799 | 49.3 |
| Nurse practitioners/ clinical nurse specialists | 186 | 16,370 | 6.5 | 14,427 | 6.5 | | |
| Nurse midwives only | 169 | 15,327 | 6.1 | 12,737 | 5.8 | | |
| Nurse midwives/ nurse practitioners | 30 | 2,764 | 1.1 | | | | |

Table 44. Advanced practice registered nurses, by employment status: 2008

*Total includes an estimated 1,144 RNs with combinations of advanced practice preparation other than those listed in the table.

| | | | | Employed i | n nursing | | |
|--|-------------------------------|--|--|---|--|---|---|
| Type of advanced practice nurse and employment status | Total estimated number* | Estimated number employed in nursing* | National certification or State Board recognition | Estimated number national certification | National certification required for job | Estimated number State Board recognition | State Board recognition required for job |
| Total | 250,527 | 220,494 | 194,661 | 177,190 | 126,050 | 176,783 | 142,228 |
| Clinical nurse specialists | 59,242 | 49,792 | 27,205 | 19,786 | 7,814 | 20,699 | 9,763 |
| With position title | | 9,327 | 7,962 | 6,409 | 3,881 | 6,610 | 5,338 |
| Without position title | | 40,466 | 19,243 | 13,377 | 3,933 | 14,089 | 4,424 |
| Nurse practitioners | 158,348 | 141,286 | 131,612 | 118,462 | 84,778 | 124,487 | 103,095 |
| With position title | | 97,876 | 97,463 | 91,518 | 72,417 | 93,320 | 86,621 |
| Without position title | | 43,410 | 34,149 | 26,944 | 12,361 | 31,167 | 16,473 |
| Nurse anesthetists** | 34,821 | 31,868 | 31,636 | 31,636 | 28,360 | 28,593 | 25,589 |
| With position title | | 29,429 | 29,429 | 29,429 | 27,252 | 26,699 | 24,462 |
| Nurse midwives | 18,492 | 15,581 | 14,256 | 13,983 | 8,097 | 10,913 | 7,841 |
| With position title | | 6,455 | 6,455 | 6,455 | 5,926 | 6,287 | 5,775 |
| Without position title | | 9,126 | 7,801 | 7,527 | | 4,626 | |

Table 45. Advanced practice registered nurses and job title, by national certification and State Board recognition: 2008

*Estimated numbers of individual specialties of advanced practice registered nurses and their percents add to more than the total of RNs who have achieved advanced practice registered nurse status because registered nurses may have had preparation in more than one advanced practice specialty.

**Nurse anesthetists without position title are not reported due to too few cases (fewer than 30 cases unweighted).

--Too few cases to report estimated number (fewer than 30 cases unweighted).

| Advanced practice | All adv | anced praction | ce RNs | Nurse pra | ctitioners** | Clinical nurs | e specialists** |
|---------------------------|-----------|----------------|-----------|-----------|--------------|---------------|-----------------|
| study focus | Number in | Estimated | Estimated | Estimated | Estimated | Estimated | Estimated |
| | sample | number* | percent | number* | percent | number* | percent |
| Total number of | | | | | | | |
| advanced practice | 2,897 | 250,527 | 100.0 | 158,348 | 100.0 | 59,242 | 100.0 |
| registered nurses | | | | | | | |
| Acute care | 239 | 21,723 | 8.7 | 11,406 | 7.2 | 7,488 | 12.6 |
| Adult health | 577 | 54,617 | 21.8 | 40,715 | 25.7 | 15,625 | 26.4 |
| Anesthesia | 475 | 33,302 | 13.3 | | | | |
| Cardiac care | 156 | 14,035 | 5.6 | 4,598 | 2.9 | 5,839 | 9.9 |
| Community health | 128 | 10,744 | 4.3 | 5,970 | 3.8 | 4,192 | 7.1 |
| Critical care | 159 | 13,779 | 5.5 | 3,359 | 2.1 | 6,440 | 10.9 |
| Family care | 810 | 68,949 | 27.5 | 66,377 | 41.9 | 4,249 | 7.2 |
| General medical surgical | 160 | 13,988 | 5.6 | 4,623 | 2.9 | 5,672 | 9.6 |
| Geriatrics/gerontology | 241 | 21,535 | 8.6 | 12,540 | 7.9 | 6,768 | 11.4 |
| Maternal-child health | 196 | 16,089 | 6.4 | 6,402 | 4.0 | 4,363 | 7.4 |
| Neonatal | 124 | 11,230 | 4.5 | 5,334 | 3.4 | | |
| Nurse-midwifery | 160 | 14,228 | 5.7 | | | | |
| Obstetrics/gynecology | 249 | 22,483 | 9.0 | 11,333 | 7.2 | | |
| Oncology | 67 | 6,550 | 2.6 | | | 3,929 | 6.6 |
| Pediatrics | 340 | 32,194 | 12.9 | 24,773 | 15.6 | 5,152 | 8.7 |
| Psychiatric/mental health | 273 | 22,471 | 9.0 | 10,792 | 6.8 | 14,265 | 24.1 |
| Women's health | 297 | 25,393 | 10.1 | 19,436 | 12.3 | | |
| Other*** | 161 | 14,436 | 5.8 | 6,755 | 4.3 | 5,919 | 10.0 |

Table 46. Advanced practice study focus: 2008

*The number of advanced practice study foci will exceed the total number of advanced practice registered nurses because each APRN may have more than one focus.

**Nurse practitioners and clinical nurse specialists represent subsets of the total of all advanced practice registered nurses. These subset groups are not mutually exclusive. An APRN who reported being trained as both nurse practitioner and clinical nurse specialist will be counted in both sets of columns.

***Other advanced practice study foci include home health, occupational health, palliative care, rehabilitation, school health, and others.

--Too few cases to report estimated number and estimated percent (fewer than 30 cases unweighted).

| | Advanced practice | e registered nurses | with certification |
|---|----------------------|---------------------|----------------------|
| Type of certification | Number in sample* | Estimated number | Estimated percent |
| Total number of advanced practice registered nurses | 2,953 | 254,711 | 100.0 |
| Acute care/critical care | 74 | 6,624 | 2.6 |
| Adult | 284 | 27,113 | 10.6 |
| Anesthesia | 389 | 26,941 | 10.6 |
| Family | 645 | 52,039 | 20.4 |
| Gerontological | 54 | 4,963 | 1.9 |
| Midwifery | 110 | 9,427 | 3.7 |
| Neonatal | 38 | 3,521 | 1.4 |
| Pediatric | 143 | 13,373 | 5.3 |
| Psychiatric/mental health | 147 | 11,163 | 4.4 |
| Women's health care | 141 | 13,198 | 5.2 |
| Other | 87 | 7,133 | 2.8 |

Table 47. Certifications in advanced practice registered nursing: 2008

*Advanced practice registered nurses may have completed educational requirements or certification in one or more specialties in advanced practice nursing. If specialty education certifications were received across disciplines, both specialty subjects are included in these distributions.

NOTE: Estimated numbers may not equal totals because of rounding, and percents will not total 100 because advanced practice registered nurses may not have reported a certification or may have reported multiple certifications.

| Age groups | Number in sample | Total estimated number* | Estimated number up to 5 years not working in nursing | Estimated percent up to 5 years not working in nursing | Estimated number 6-10 years not working in nursing | Estimated percent 6-10 years not working in nursing | Estimated number 11-15 years not working in nursing | Estimated percent 11-15 years not working in nursing | Estimated number 16 or more years not working in nursing | Estimated percent 16 or more years not working in nursing |
|--------------|---------------------|-------------------------------|---|--|--|---|---|--|--|---|
| Total | 4,071 | 415,450 | 230,432 | 100.0 | 87,496 | 100.0 | 44,814 | 100.0 | 47,733 | 100.0 |
| Less than 30 | 77 | 7,511 | 6,302 | 2.7 | | | | | | |
| 30-39 | 376 | 43,488 | 31,189 | 13.5 | 8,499 | 9.7 | | | | |
| 40-44 | 267 | 28,883 | 15,551 | 6.7 | 8,943 | 10.2 | 3,142 | 7.0 | | |
| 45-49 | 349 | 36,875 | 17,433 | 7.6 | 8,976 | 10.3 | 5,664 | 12.6 | 4,748 | 9.9 |
| 50-54 | 558 | 54,778 | 23,626 | 10.3 | 11,077 | 12.7 | 7,904 | 17.6 | 11,594 | 24.3 |
| 55-59 | 572 | 52,897 | 30,710 | 13.3 | 9,513 | 10.9 | 4,655 | 10.4 | 7,580 | 15.9 |
| 60-64 | 718 | 69,001 | 42,571 | 18.5 | 11,677 | 13.3 | 6,298 | 14.1 | 8,164 | 17.1 |
| 65-69 | 667 | 67,573 | 39,142 | 17.0 | 15,121 | 17.3 | 6,586 | 14.7 | 6,261 | 13.1 |
| 70-74 | 302 | 32,962 | 14,949 | 6.5 | 9,387 | 10.7 | 5,351 | 11.9 | | |
| 75 and over | 185 | 21,482 | 8,961 | 3.9 | 4,152 | 4.7 | 3,778 | 8.4 | 4,592 | 9.6 |

Table 48. Age distribution of registered nurses not employed in nursing,
by length of time not working in nursing: 2008

*Includes an estimated 4,975 licensed registered nurses who never worked in nursing.

--Too few cases to report estimated number and estimated percent (fewer than 30 cases unweighted).

| Table 49. | Age and presence of children, by employment status |
|-----------|--|
| | in principal nursing position: 2008 |

| Age and presence of children | Total number in sample | Total estimated number | Total estimated percent | Number in sample employed in nursing full- time | Estimated number employed in nursing full- time | Estimated percent employed in nursing full- time | Number in sample employed in nursing part- time | Estimated number employed in nursing part- time | Estimated percent employed in nursing part- time | Number in sample not employed in nursing | Estimated number not employed in nursing | Estimated percent not employed in nursing |
|------------------------------|---------------------------|------------------------------|-------------------------------|---|---|--|---|---|--|---|---|--|
| Total | 33,179 | 3,063,162 | 100.0 | 21,307 | 1,937,405 | 63.2 | 7,247 | 659,194 | 21.5 | 4,625 | 466,564 | 15.2 |
| Less than 40 years old | 8,522 | 902,155 | 100.0 | 6,108 | 644,826 | 71.5 | 1,802 | 190,941 | 21.2 | 612 | 66,388 | 7.4 |
| Children under 6 only | 2,249 | 243,121 | 100.0 | 1,333 | 143,139 | 58.9 | 708 | 77,565 | 31.9 | 208 | 22,417 | 9.2 |
| Children 6-18 only | 1,980 | 212,340 | 100.0 | 1,482 | 158,906 | 74.8 | 377 | 39,674 | 18.7 | 121 | 13,761 | 6.5 |
| Children in both age groups | 1,423 | 149,485 | 100.0 | 863 | 91,462 | 61.2 | 421 | 43,229 | 28.9 | 139 | 14,794 | 9.9 |
| No children | 2,870 | 297,209 | 100.0 | 2,430 | 251,319 | 84.6 | 296 | 30,474 | 10.3 | 144 | 15,416 | 5.2 |
| 40-49 years old | 8,386 | 791,932 | 100.0 | 5,805 | 537,400 | 67.9 | 1,821 | 174,519 | 22.0 | 760 | 80,013 | 10.1 |
| Children under 6 only | 318 | 30,981 | 100.0 | 194 | 18,330 | 59.2 | 79 | 7,229 | 23.3 | 45 | 5,423 | 17.5 |
| Children 6-18 only | 4,386 | 428,646 | 100.0 | 2,843 | 272,088 | 63.5 | 1,146 | 113,919 | 26.6 | 397 | 42,639 | 9.9 |
| Children in both age groups | 414 | 38,151 | 100.0 | 238 | 20,520 | 53.8 | 114 | 10,801 | 28.3 | 62 | 6,830 | 17.9 |
| No children | 3,268 | 294,154 | 100.0 | 2,530 | 226,462 | 77.0 | 482 | 42,570 | 14.5 | 256 | 25,122 | 8.5 |
| 50 years and older | 16,271 | 1,369,074 | 100.0 | 9,394 | 755,179 | 55.2 | 3,624 | 293,734 | 21.5 | 3,253 | 320,162 | 23.4 |
| Children under 6 only | 145 | 13,006 | 100.0 | 101 | 9,121 | 70.1 | 19 | | | 25 | | |
| Children 6-18 only | 2,026 | 174,062 | 100.0 | 1,283 | 105,956 | 60.9 | 492 | 41,532 | 23.9 | 251 | 26,574 | 15.3 |
| Children in both age groups | 81 | 7,771 | 100.0 | 54 | 4,937 | 63.5 | 13 | | | 14 | | |
| No children | 14,019 | 1,174,235 | 100.0 | 7,956 | 635,165 | 54.1 | 3,100 | 249,822 | 21.3 | 2,963 | 289,248 | 24.6 |

--Too few cases to report estimated number and estimated percent (fewer than 30 cases unweighted). NOTE: Estimated numbers may not equal total, and percents may not add to 100, because of rounding.

| Age group | Number in sample | Total estimated number | Total estimated percent | Estimated number resident State same in 2007 and 2008 | Estimated percent resident State same in 2007 and 2008 | Estimated number resident State different in 2007 and 2008* | Estimated percent resident State different in 2007 and 2008* |
|--------------|---------------------|------------------------------|-------------------------------|--|---|--|---|
| Total | 33,179 | 3,063,162 | 100.0 | 2,965,809 | 96.8 | 97,353 | 3.2 |
| Less than 25 | 730 | 79,641 | 100.0 | 71,838 | 90.2 | 7,803 | 9.8 |
| 25-29 | 1,966 | 208,543 | 100.0 | 192,028 | 92.1 | 16,515 | 7.9 |
| 30-34 | 2,677 | 281,999 | 100.0 | 267,969 | 95.0 | 14,030 | 5.0 |
| 35-39 | 3,149 | 331,972 | 100.0 | 319,042 | 96.1 | 12,930 | 3.9 |
| 40-44 | 3,623 | 350,101 | 100.0 | 340,459 | 97.2 | 9,642 | 2.8 |
| 45-49 | 4,763 | 441,831 | 100.0 | 433,489 | 98.1 | 8,342 | 1.9 |
| 50-54 | 6,035 | 495,985 | 100.0 | 486,055 | 98.0 | 9,930 | 2.0 |
| 55-59 | 4,858 | 396,967 | 100.0 | 387,513 | 97.6 | 9,454 | 2.4 |
| 60-64 | 2,995 | 249,533 | 100.0 | 243,671 | 97.7 | 5,862 | 2.3 |
| 65-69 | 1,509 | 140,263 | 100.0 | 138,174 | 98.5 | 2,089 | 1.5 |
| 70-74 | 562 | 53,670 | 100.0 | 52,979 | 98.7 | | |
| 75 and over | 312 | 32,657 | 100.0 | 32,591 | 99.8 | | |

Table 50. Resident State in 2007 and 2008, by age group: 2008

*Resident State different includes cases where either 2007 or 2008 residence was a foreign country or a U.S. territory.

--Too few cases to report estimated number and estimated percent (fewer than 30 cases unweighted).

| Length of time since worked | Number in sample | Total estimated number* | Total estimated percent | Estimated number seeking nursing employment | Estimated percent seeking nursing employment | Estimated number with health-related employment | Estimated percent with health-related employment | Estimated number with nonhealth- related employment | Estimated percent with nonhealth- related employment |
|-----------------------------|---------------------|-------------------------------|-------------------------------|---|--|--|---|---|--|
| Total | 4,071 | 415,450 | 100.0 | 22,770 | 100.0 | 57,983 | 100.0 | 45,824 | 100.0 |
| Less than 1 year | 517 | 46,204 | 11.1 | 6,582 | 28.9 | 5,324 | 9.2 | 3,744 | 8.2 |
| 1-4 years | 1,692 | 163,595 | 39.4 | 8,320 | 36.5 | 16,827 | 29.0 | 12,345 | 26.9 |
| 5-9 years | 776 | 83,417 | 20.1 | 3,344 | 14.7 | 12,524 | 21.6 | 11,935 | 26.0 |
| 10-19 years | 738 | 84,677 | 20.4 | | | 15,577 | 26.9 | 10,122 | 22.1 |
| 20 years or more | 313 | 32,583 | 7.8 | | | 6,451 | 11.1 | 6,642 | 14.5 |

Table 51. Registered nurses not employed in nursing and length of time since last worked as a nurse, by whether or not registered nurse was seeking nursing position or had other occupation outside of nursing: 2008

*Includes an estimated 4,975 licensed registered nurses who never worked in nursing. Also includes an estimated 2,147 registered nurses who were seeking nursing employment while employed in a non-nursing, health-related occupation, and an estimated 2,926 registered nurses who were seeking nursing employment while employed in a nonhealth-related occupation.

--Too few cases to report estimated number and estimated percent (fewer than 30 cases unweighted).

| • | | • | |
|---|---------------------|----------------------------|----------------------------|
| Type of employment sought and months looking | Number in sample | Total estimated number* | Total estimated percent |
| Total | 231 | 22,770 | 100.0 |
| Type of employment | | | |
| Full-time | 60 | 5878 | 25.8 |
| Part-time | 94 | 8,950 | 39.3 |
| Either | 77 | 7,943 | 34.9 |
| Number of months looking | | | |
| Up to 1 month | 96 | 8,608 | 37.8 |
| 2-3 months | 40 | 3,750 | 16.5 |
| 4-9 months | 66 | 7,206 | 31.6 |
| More than 9 months | 29 | | |

Table 52. Registered nurses not employed in nursing andactively seeking nursing employment, by type of employmentsought and number of months looking: 2008

*Of the estimated 22,770 RNs not employed in nursing and actively seeking nursing employment, and an estimated 5,073 were employed in non-nursing occupations and an estimated 17,697 were not employed for pay in any occupation.

--Too few cases to report estimated number and estimated percent (fewer than 30 cases unweighted).

Table 53. Age and caring for adults at home or others elsewhere, by employment status in principal nursing position: 2008

| Age and presence of adults and dependents | Total number in sample | Total estimated number | Total estimated percent | Number in sample employed in nursing full- time | Estimated number employed in nursing full- time | | Number in sample employed in nursing part- time | | | | Estimated number not employed in nursing | |
|---|---------------------------|------------------------------|-------------------------------|---|---|------|---|---------|------|-------|---|------|
| Total | 33,179 | 3,063,162 | 100.0 | 21,307 | 1,937,405 | 63.2 | 7,247 | 659,194 | 21.5 | 4,625 | 466,564 | 15.2 |
| Less than 50 years old | 16,908 | 1,694,088 | 100.0 | 11,913 | 1,182,226 | 69.8 | 3,623 | 365,460 | 21.6 | 1,372 | 146,402 | 8.6 |
| Adult dependents at home | 1,745 | 183,626 | 100.0 | 1,325 | 139,417 | 75.9 | 284 | 30,034 | 16.4 | 136 | 14,175 | 7.7 |
| Other dependents living elsewhere | 1,657 | 164,062 | 100.0 | 1,177 | 113,370 | 69.1 | 326 | 33,589 | 20.5 | 154 | 17,103 | 10.4 |
| Both adults at home and others living elsewhere | 390 | 38,298 | 100.0 | 296 | 29,033 | 75.8 | 76 | 7,555 | 19.7 | 18 | | |
| No adult dependents | 13,116 | 1,308,102 | 100.0 | 9,115 | 900,407 | 68.8 | 2,937 | 294,282 | 22.5 | 1,064 | 113,413 | 8.7 |
| 50-59 years old | 10,893 | 892,952 | 100.0 | 7,416 | 596,847 | 66.8 | 2,186 | 174,919 | 19.6 | 1,291 | 121,186 | 13.6 |
| Adult dependents at home | 1,741 | 154,432 | 100.0 | 1,228 | 107,568 | 69.7 | 313 | 27,071 | 17.5 | 200 | 19,793 | 12.8 |
| Other dependents living elsewhere | 1,900 | 159,702 | 100.0 | 1,268 | 104,286 | 65.3 | 394 | 32,938 | 20.6 | 238 | 22,478 | 14.1 |
| Both adults at home and others living elsewhere | 537 | 45,880 | 100.0 | 374 | 31,906 | 69.5 | 100 | 7,798 | 17.0 | 63 | 6,177 | 13.5 |
| No adult dependents | 6,715 | 532,937 | 100.0 | 4,546 | 353,087 | 66.3 | 1,379 | 107,113 | 20.1 | 790 | 72,738 | 13.6 |
| 60 years and older | 5,378 | 476,123 | 100.0 | 1,978 | 158,332 | 33.3 | 1,438 | 118,815 | 25.0 | 1,962 | 198,976 | 41.8 |
| Adult dependents at home | 563 | 49,720 | 100.0 | 254 | 20,725 | 41.7 | 121 | 9,663 | 19.4 | 188 | 19,332 | 38.9 |
| Other dependents living elsewhere | 583 | 51,510 | 100.0 | 229 | 17,486 | 33.9 | 148 | 12,137 | 23.6 | 206 | 21,886 | 42.5 |
| Both adults at home and others living elsewhere | 96 | 8,538 | 100.0 | 38 | 2,770 | 32.4 | 20 | | | 38 | 4,149 | 48.6 |
| No adult dependents | 4,136 | 366,355 | 100.0 | 1,457 | 117,351 | 32.0 | 1,149 | 95,396 | 26.0 | 1,530 | 153,608 | 41.9 |

--Too few cases to report estimated number and estimated percent (fewer than 30 cases unweighted).

| | | | | Number in | Estimated | Estimated | Number in | Estimated | Estimated | | | |
|--------------------------------|-----------|-----------|-----------|---------------|---------------|---------------|---------------|----------------|---------------|-------------|-------------|-------------|
| | | | | sample | number | percent | sample | number | percent | Number in | Estimated | Estimated |
| | Total | Total | Total | employed in | employed in | sample not | number not | percent not |
| Marital status and presence of | number in | estimated | estimated | nursing full- | nursing full- | nursing full- | nursing part- | -nursing part- | nursing part- | employed in | employed in | employed ir |
| children | sample | number | percent | time | time | time | time | time | time | nursing | nursing | nursing |
| larried/domestic partnership | 24,380 | 2,259,553 | 73.8 | 15,020 | 1,368,085 | 70.6 | 5,926 | 543,404 | 82.4 | 3,434 | 348,065 | 74.6 |
| Children under 6 only | 2,464 | 261,699 | 8.5 | 1,434 | 151,025 | 7.8 | 771 | 82,475 | 12.5 | 259 | 28,199 | 6.0 |
| Children 6-18 only | 6,932 | 670,920 | 21.9 | 4,441 | 422,643 | 21.8 | 1,826 | 176,325 | 26.7 | 665 | 71,952 | 15.4 |
| Children in both age groups | 1,735 | 177,082 | 5.8 | 1,016 | 103,650 | 5.3 | 519 | 51,645 | 7.8 | 200 | 21,787 | 4.7 |
| No children | 13,249 | 1,149,853 | 37.5 | 8,129 | 690,767 | 35.7 | 2,810 | 232,959 | 35.3 | 2,310 | 226,127 | 48.5 |
| Vidowed/separated/divorced | 5,854 | 515,455 | 16.8 | 3,993 | 344,117 | 17.8 | 957 | 82,303 | 12.5 | 904 | 89,035 | 19.1 |
| Children under 6 only | 163 | 15,587 | 0.5 | 122 | 11,205 | 0.6 | 26 | | | 15 | | |
| Children 6-18 only | 1,221 | 118,390 | 3.9 | 961 | 91,771 | 4.7 | 169 | 16,883 | 2.6 | 91 | 9,736 | 2.1 |
| Children in both age groups | 157 | 15,651 | 0.5 | 118 | 11,215 | 0.6 | 26 | | | 13 | | |
| No children | 4,313 | 365,827 | 11.9 | 2,792 | 229,926 | 11.9 | 736 | 59,704 | 9.1 | 785 | 76,196 | 16.3 |
| lever married | 2,945 | 288,153 | 9.4 | 2,294 | 225,203 | 11.6 | 364 | 33,486 | 5.1 | 287 | 29,464 | 6.3 |
| Children under 6 only | 85 | 9,822 | 0.3 | 72 | 8,360 | 0.4 | 9 | | | 4 | | |
| Children 6-18 only | 239 | 25,738 | 0.8 | 206 | 22,535 | 1.2 | 20 | | | 13 | | |
| Children in both age groups | 26 | | | 21 | | | 3 | | | 2 | | |
| No children | 2,595 | 249,918 | 8.2 | 1,995 | 192,252 | 9.9 | 332 | 30,202 | 4.6 | 268 | 27,464 | 5.9 |

6 - I. 11 -_ . . _

| Table 55. Marital status and caring for adults at home or others elsewhere, by employment status in principal nursing position: 2008 | | | | | | | | | | | |
|--|---|--|--|---|--|--|--|--|--|---|--|
| | | | Number in | Estimated | Estimated | Number in | Estimated | Estimated | | Estimated | Estimated |
| | | | sample | number | percent | sample | number | percent | Number in | number | percent |
| | | | 1 2 | | | | 1 2 | | | not | not |
| | | | 0 | - | • | 0 | 0 | 0 | | 1 2 | employed |
| | | • | | | | | | • | <u> </u> | 0 | in nursing |
| 33,179 | 3,063,162 | 100.0 | 21,307 | 1,937,405 | 100.0 | 7,247 | 659,194 | 100.0 | 4,625 | 466,564 | 100.0 |
| | | | | | | | | | | | |
| 24,380 | 2,259,553 | 73.8 | 15,020 | 1,368,085 | 70.6 | 5,926 | 543,404 | 82.4 | 3,434 | 348,065 | 74.6 |
| 3,596 | 338,937 | 11.1 | 2,414 | 225,503 | 11.6 | 720 | 65,619 | 10.0 | 462 | 47,815 | 10.2 |
| 20,784 | 1,920,616 | 62.7 | 12,606 | 1,142,582 | 59.0 | 5,206 | 477,785 | 72.5 | 2,972 | 300,249 | 64.4 |
| 5,854 | 515,455 | 16.8 | 3,993 | 344,117 | 17.8 | 957 | 82,303 | 12.5 | 904 | 89,035 | 19.1 |
| 1,073 | 96,929 | 3.2 | 805 | 72,170 | 3.7 | 135 | 12,430 | 1.9 | 133 | 12,329 | 2.6 |
| 4,781 | 418,526 | 13.7 | 3,188 | 271,948 | 14.0 | 822 | 69,873 | 10.6 | 771 | 76,706 | 16.4 |
| 2,945 | 288,153 | 9.4 | 2,294 | 225,203 | 11.6 | 364 | 33,486 | 5.1 | 287 | 29,464 | 6.3 |
| 403 | 44,629 | 1.5 | 296 | 33,746 | 1.7 | 59 | 5,690 | 0.9 | 48 | 5,192 | 1.1 |
| 2,542 | 243,525 | 8.0 | 1,998 | 191,457 | 9.9 | 305 | 27,796 | 4.2 | 239 | 24,272 | 5.2 |
| | | | | | | | | | | | |
| 24,380 | 2,259,553 | 73.8 | 15,020 | 1,368,085 | 70.6 | 5,926 | 543,404 | 82.4 | 3,434 | 348,065 | 74.6 |
| 4,028 | 367,445 | 12.0 | 2,525 | 224,110 | 11.6 | 919 | 82,252 | 12.5 | 584 | 61,083 | 13.1 |
| 20,352 | 1,892,109 | 61.8 | 12,495 | 1,143,975 | 59.0 | 5,007 | 461,151 | 70.0 | 2,850 | 286,982 | 61.5 |
| 5,854 | 515,455 | 16.8 | 3,993 | 344,117 | 17.8 | 957 | 82,303 | 12.5 | 904 | 89,035 | 19.1 |
| 952 | 81,603 | 2.7 | 711 | 60,183 | 3.1 | 121 | 10,945 | 1.7 | 120 | 10,475 | 2.2 |
| 4,902 | 433,853 | 14.2 | 3,282 | 283,935 | 14.7 | 836 | 71,358 | 10.8 | 784 | 78,560 | 16.8 |
| 2,945 | 288,153 | 9.4 | 2,294 | 225,203 | 11.6 | 364 | 33,486 | 5.1 | 287 | 29,464 | 6.3 |
| 183 | 18,942 | 0.6 | 146 | 14,558 | 0.8 | 24 | | | 13 | | |
| 2,762 | 269,211 | 8.8 | 2,148 | 210,644 | 10.9 | 340 | 31,049 | 4.7 | 274 | 27,518 | 5.9 |
| | Total number in sample 33,179 24,380 3,596 20,784 5,854 1,073 4,781 2,945 403 2,542 24,380 4,028 20,352 5,854 952 4,902 2,945 183 | Total number in sample Total estimated number 33,179 3,063,162 24,380 2,259,553 3,596 338,937 20,784 1,920,616 5,854 515,455 1,073 96,929 4,781 418,526 2,945 288,153 403 44,629 2,542 243,525 24,380 2,259,553 4,028 367,445 20,352 1,892,109 5,854 515,455 952 81,603 4,902 433,853 2,945 288,153 183 18,942 | $\begin{array}{c c c c c c c c c c c c c c c c c c c $ | Total number in sample Total estimated number in sample Total estimated percent Number in sample in nursing percent 33,179 3,063,162 100.0 21,307 24,380 2,259,553 73.8 15,020 3,596 338,937 11.1 2,414 20,784 1,920,616 62.7 12,606 5,854 515,455 16.8 3,993 1,073 96,929 3.2 805 4,781 418,526 13.7 3,188 2,945 288,153 9.4 2,294 403 44,629 1.5 296 2,542 243,525 8.0 1,998 24,380 2,259,553 73.8 15,020 4,028 367,445 12.0 2,525 20,352 1,892,109 61.8 12,495 5,854 515,455 16.8 3,993 952 81,603 2.7 711 4,902 433,853 14.2 3,282 2,945 28 | $\begin{array}{c c c c c c c c c c c c c c c c c c c $ | $\begin{array}{c c c c c c c c c c c c c c c c c c c $ | $\begin{array}{c c c c c c c c c c c c c c c c c c c $ | $\begin{array}{c c c c c c c c c c c c c c c c c c c $ | Total number in sample Total number Total number Total number Total estimated number Number in sample Estimated number Estimated employed in nursing Estimated number Estimated number Estimated number Estimated employed in nursing Estimated mumber Estimated employed in nursing Estimated mumber Estimated employed in nursing Estimated percent Estimated employed in nursing 24,380 2,259,553 73.8 15,020 1,368,085 70.6 5,926 543,404 82.4 3,596 338,937 11.1 2,414 225,503 11.6 720 65,619 10.0 20,784 1,920,616 62.7 12,606 1,142,582 59.0 5,206 477,785 72.5 5,854 515,455 16.8 3,993 344,117 17.8 957 82,303 12.5 <td< td=""><td>Total number Total estimated sample Number in employed Estimated number Number in employed Estimated number Number in employed Estimated mumber Number in employed Estimated mumber Number in employed Estimated mumber Number in employed Estimated mumber Number in employed 33,179 3,063,162 100.0 21,307 1,937,405 100.0 7,247 659,194 100.0 4,625 24,380 2,259,553 73.8 15,020 1,368,085 70.6 5,926 543,404 82.4 3,434 3,596 338,937 11.1 2,414 225,503 11.6 720 65,619 10.0 462 20,784 1,920,616 62.7 12,606 1,142,582 59.0 5,206 477,785 72.5 2,972 5,854 515,455 16.8 3,993 344,117 17.8 957 82,303 12.5 904 1,073 96,929 3.2 805 72,170 3.7 135 12,430 1.9 133</td></td<> <td>Total number Total estimated number Number in sample employed Estimated number in number Estimated employed Number in sample employed Estimated percent Sample sample in nursing Estimated number Estimated employed Estimated percent Sample employed Estimated percent Estimated sample Estimated employed Estimated percent Sample employed Estimated percent Estimated percent Estimated percent Estimated percent Estimated percent Sample employed Inumber employed Estimated percent Estimated percent Estimated percent Estimated percent Sample employed Inumber employed Inumber employed Inumber part-time Inumbe</td> | Total number Total estimated sample Number in employed Estimated number Number in employed Estimated number Number in employed Estimated mumber Number in employed Estimated mumber Number in employed Estimated mumber Number in employed Estimated mumber Number in employed 33,179 3,063,162 100.0 21,307 1,937,405 100.0 7,247 659,194 100.0 4,625 24,380 2,259,553 73.8 15,020 1,368,085 70.6 5,926 543,404 82.4 3,434 3,596 338,937 11.1 2,414 225,503 11.6 720 65,619 10.0 462 20,784 1,920,616 62.7 12,606 1,142,582 59.0 5,206 477,785 72.5 2,972 5,854 515,455 16.8 3,993 344,117 17.8 957 82,303 12.5 904 1,073 96,929 3.2 805 72,170 3.7 135 12,430 1.9 133 | Total number Total estimated number Number in sample employed Estimated number in number Estimated employed Number in sample employed Estimated percent Sample sample in nursing Estimated number Estimated employed Estimated percent Sample employed Estimated percent Estimated sample Estimated employed Estimated percent Sample employed Estimated percent Estimated percent Estimated percent Estimated percent Estimated percent Sample employed Inumber employed Estimated percent Estimated percent Estimated percent Estimated percent Sample employed Inumber employed Inumber employed Inumber part-time Inumbe |

Table 55. Marital status and caring for adults at home or others elsewhere, by employment status in principal nursing position: 2008

--Too few cases to report estimated number and estimated percent (fewer than 30 cases unweighted).

| | | | Number in | Estimated | Estimated | Number in | Estimated | Estimated | | | |
|--------------|---|--|--|--|--|---|---|---|---|---|--|
| | | | sample | number | percent | sample | number | percent | Number in | Estimated | Estimated |
| | Total | Total | employed in | employed in | employed in | employed in | employed in | employed in | sample not | number not | percent not |
| Total number | estimated | estimated | nursing full- | nursing full- | nursing full- | nursing part- | nursing part- | - nursing part- | employed in | employed in | employed in |
| in sample | number | percent | time | time | time | time | time | time | nursing | nursing | nursing |
| 33,179 | 3,063,162 | 100.0 | 21,307 | 1,937,405 | 100.0 | 7,247 | 659,194 | 100.0 | 4,625 | 466,564 | 100.0 |
| 260 | 25,673 | 0.8 | 3 | | | 76 | 7,162 | 1.1 | 181 | 18,118 | 3.9 |
| 382 | 35,099 | 1.1 | 21 | | | 112 | 9,326 | 1.4 | 249 | 24,436 | 5.2 |
| 768 | 71,485 | 2.3 | 119 | 9,925 | 0.5 | 246 | 21,801 | 3.3 | 403 | 39,760 | 8.5 |
| 2,754 | 250,131 | 8.2 | 1,375 | 123,373 | 6.4 | 751 | 64,691 | 9.8 | 628 | 62,067 | 13.3 |
| 7,796 | 705,874 | 23.0 | 5,363 | 480,099 | 24.8 | 1,523 | 133,354 | 20.2 | 910 | 92,422 | 19.8 |
| 8,188 | 747,704 | 24.4 | 5,764 | 522,626 | 27.0 | 1,682 | 153,799 | 23.3 | 742 | 71,279 | 15.3 |
| 8,386 | 786,510 | 25.7 | 5,861 | 542,026 | 28.0 | 1,797 | 169,533 | 25.7 | 728 | 74,951 | 16.1 |
| 2,787 | 266,890 | 8.7 | 1,848 | 172,746 | 8.9 | 575 | 55,748 | 8.5 | 364 | 38,396 | 8.2 |
| 1,858 | 173,795 | 5.7 | 953 | 84,880 | 4.4 | 485 | 43,781 | 6.6 | 420 | 45,134 | 9.7 |
| 24,380 | 2,259,553 | 100.0 | 15,020 | 1,368,085 | 100.0 | 5,926 | 543,404 | 100.0 | 3,434 | 348,065 | 100.0 |
| 92 | 8,369 | 0.4 | 2 | | | 31 | 2,392 | 0.4 | 59 | 5,596 | 1.6 |
| 109 | 9,688 | 0.4 | 6 | | | 37 | 3,201 | 0.6 | 66 | 6,263 | 1.8 |
| 334 | 31,673 | 1.4 | 37 | 2,526 | 0.2 | 102 | 9,709 | 1.8 | 195 | 19,438 | 5.6 |
| 1,249 | 111,828 | 4.9 | 460 | 39,381 | 2.9 | 402 | 34,147 | 6.3 | 387 | 38,300 | 11.0 |
| 4,427 | 404,574 | 17.9 | 2,620 | 237,795 | 17.4 | 1,121 | 97,695 | 18.0 | 686 | 69,084 | 19.8 |
| 6,291 | 574,884 | 25.4 | 4,159 | 376,732 | 27.5 | 1,500 | 137,232 | 25.3 | 632 | 60,920 | 17.5 |
| 7,510 | 702,926 | 31.1 | 5,137 | 472,550 | 34.5 | 1,712 | 162,104 | 29.8 | 661 | 68,272 | 19.6 |
| 2,597 | 248,852 | 11.0 | 1,699 | 158,186 | 11.6 | 551 | 53,790 | 9.9 | 347 | 36,877 | 10.6 |
| 1,771 | 166,759 | 7.4 | 900 | 80,312 | 5.9 | 470 | 43,134 | 7.9 | 401 | 43,314 | 12.4 |
| | in sample 33,179 260 382 768 2,754 7,796 8,188 8,386 2,787 1,858 24,380 92 109 334 1,249 4,427 6,291 7,510 2,597 | Total numberestimated number33,1793,063,16226025,67338235,09976871,4852,754250,1317,796705,8748,188747,7048,386786,5102,787266,8901,858173,79524,3802,259,553928,3691099,68833431,6731,249111,8284,427404,5746,291574,8847,510702,9262,597248,852 | $\begin{array}{c c c c c c c c c c c c c c c c c c c $ | $\begin{array}{c c c c c c c c c c c c c c c c c c c $ | $\begin{array}{c c c c c c c c c c c c c c c c c c c $ | $ \begin{array}{c c c c c c c c c c c c c c c c c c c $ | $ \begin{array}{c c c c c c c c c c c c c c c c c c c $ | $ \begin{array}{ c c c c c c c c c c c c c c c c c c c$ | $ \begin{array}{c c c c c c c c c c c c c c c c c c c $ | $ \begin{array}{c c c c c c c c c c c c c c c c c c c $ | Total Total Sample number percent sample number percent number fermployed number numsing numsing numsing numsing numsing numsing numsing |

Table 56. Household income and marital status, by employment status in principal nursing position: 2008

--Too few cases to report estimated number and estimated percent (fewer than 30 cases unweighted). NOTE: Estimated numbers may not equal total, and percents may not add to 100, because of rounding.

| State and geographic area | Number in sample | Total estimated number | Estimated number employed in nursing | Estimated percent employed in nursing | Estimated number not employed in nursing | Estimated percent not employed in nursing | Employed nurses per 100,000* | State licensure rate, per 100,000* |
|---------------------------|------------------|---------------------------|--------------------------------------|---------------------------------------|--|---|---------------------------------|---------------------------------------|
| United States | 33,179 | 3,063,162 | 2,596,599 | 84.8 | 466,564 | 15.2 | 854 | 1,007 |
| New England | 3,614 | 187,542 | 161,627 | 86.2 | 25,915 | 13.8 | 1,130 | 1,311 |
| Connecticut | 584 | 42,215 | 35,343 | 83.7 | 6,871 | 16.3 | 1,009 | 1,206 |
| Maine | 517 | 19,249 | 16,279 | 84.6 | 2,971 | 15.4 | 1,237 | 1,462 |
| Massachusetts | 868 | 89,434 | 77,574 | 86.7 | 11,860 | 13.3 | 1,194 | 1,376 |
| New Hampshire | 628 | 15,468 | 13,562 | 87.7 | 1,906 | 12.3 | 1,031 | 1,176 |
| Rhode Island | 494 | 12,887 | 11,531 | 89.5 | 1,356 | 10.5 | 1,097 | 1,226 |
| Vermont | 523 | 8,290 | 7,338 | 88.5 | 951 | 11.5 | 1,181 | 1,334 |
| Middle Atlantic | 3,178 | 462,263 | 377,938 | 81.8 | 84,325 | 18.2 | 930 | 1,138 |
| New Jersey | 745 | 89,314 | 73,634 | 82.4 | 15,679 | 17.6 | 848 | 1,029 |
| New York | 1,246 | 203,597 | 167,427 | 82.2 | 36,170 | 17.8 | 859 | 1,045 |
| Pennsylvania | 1,187 | 169,352 | 136,876 | 80.8 | 32,476 | 19.2 | 1,100 | 1,360 |
| South Atlantic | 6,604 | 576,337 | 481,050 | 83.5 | 95,287 | 16.5 | 824 | 987 |
| Delaware | 605 | 10,547 | 9,169 | 86.9 | 1,377 | 13.1 | 1,050 | 1,208 |
| District of Columbia | 434 | 11,487 | 11,053 | 96.2 | | | 1,868 | 1,941 |
| Florida | 1,227 | 186,349 | 150,321 | 80.7 | 36,027 | 19.3 | 820 | 1,017 |
| Georgia | 634 | 84,489 | 68,254 | 80.8 | 16,235 | 19.2 | 705 | 872 |
| Maryland | 848 | 55,276 | 50,159 | 90.7 | 5,117 | 9.3 | 890 | 981 |
| North Carolina | 825 | 96,864 | 82,104 | 84.8 | 14,760 | 15.2 | 890 | 1,050 |
| South Carolina | 469 | 41,371 | 37,472 | 90.6 | 3,899 | 9.4 | 836 | 923 |
| Virginia | 1,077 | 70,499 | 54,988 | 78.0 | 15,511 | 22.0 | 708 | 907 |
| West Virginia | 485 | 19,456 | 17,529 | 90.1 | 1,927 | 9.9 | 966 | 1,072 |
| East South Central | 1,987 | 195,713 | 171,020 | 87.4 | 24,693 | 12.6 | 946 | 1,082 |
| Alabama | 534 | 49,780 | 41,488 | 83.3 | 8,291 | 16.7 | 890 | 1,068 |
| Kentucky | 487 | 46,473 | 41,520 | 89.3 | 4,953 | 10.7 | 973 | 1,089 |
| Mississippi | 430 | 30,801 | 27,414 | 89.0 | 3,387 | 11.0 | 933 | 1,048 |
| Tennessee | 536 | 68,660 | 60,598 | 88.3 | 8,063 | 11.7 | 975 | 1,105 |
| West South Central | 2,697 | 290,304 | 249,889 | 86.1 | 40,415 | 13.9 | 709 | 824 |
| Arkansas | 503 | 26,096 | 22,443 | 86.0 | 3,653 | 14.0 | 786 | 914 |
| Louisiana | 491 | 41,863 | 36,859 | 88.0 | 5,004 | 12.0 | 836 | 949 |
| Oklahoma | 490 | 32,522 | 27,256 | 83.8 | 5,266 | 16.2 | 748 | 893 |
| Texas | 1,213 | 189,823 | 163,331 | 86.0 | 26,492 | 14.0 | 671 | 780 |

Table 57. Registered nurses in each State and geographic area, by activity status: 2008

Footnotes at end of table.

| State and geographic area | Number in sample | Total estimated number | Estimated number employed in nursing | Estimated percent employed in nursing | Estimated number not employed in nursing | Estimated percent not employed in nursing | Employed nurses per 100,000* | State licensure rate, per 100,000* |
|---------------------------|------------------|---------------------------|---|---------------------------------------|--|---|---------------------------------|------------------------------------|
| East North Central | 3,882 | 516.080 | 435,942 | 84.5 | 80,138 | 15.5 | 940 | 1,112 |
| Illinois | 922 | 136,213 | 111,939 | 82.2 | 24,275 | 17.8 | 868 | 1,056 |
| Indiana | 579 | 69,672 | 58,089 | 83.4 | 11,584 | 16.6 | 911 | 1,093 |
| Michigan | 733 | 105,222 | 88,449 | 84.1 | 16,773 | 15.9 | 884 | 1,052 |
| Ohio | 1,032 | 138,743 | 120,642 | 87.0 | 18,100 | 13.0 | 1,050 | 1,208 |
| Wisconsin | 616 | 66,229 | 56,823 | 85.8 | 9,407 | 14.2 | 1,010 | 1,177 |
| West North Central | 3,814 | 251,281 | 221,392 | 88.1 | 29,889 | 11.9 | 1,098 | 1,246 |
| lowa | 525 | 38,208 | 33,929 | 88.8 | 4,279 | 11.2 | 1,130 | 1,273 |
| Kansas | 553 | 32,165 | 27,032 | 84.0 | 5,134 | 16.0 | 965 | 1,148 |
| Minnesota | 650 | 67,551 | 58,391 | 86.4 | 9,160 | 13.6 | 1,119 | 1,294 |
| Missouri | 658 | 72,698 | 64,064 | 88.1 | 8,634 | 11.9 | 1,084 | 1,230 |
| Nebraska | 479 | 20,501 | 19,086 | 93.1 | 1,415 | 6.9 | 1,070 | 1,150 |
| North Dakota | 500 | 8,566 | 8,169 | 95.4 | | | 1,273 | 1,335 |
| South Dakota | 449 | 11,591 | 10,720 | 92.5 | 871 | 7.5 | 1,333 | 1,441 |
| Mountain | 3,780 | 187,159 | 161,591 | 86.3 | 25,568 | 13.7 | 742 | 859 |
| Arizona | 528 | 55,001 | 49,325 | 89.7 | 5,676 | 10.3 | 759 | 846 |
| Colorado | 668 | 47,805 | 39,461 | 82.5 | 8,344 | 17.5 | 799 | 968 |
| Idaho | 502 | 12,922 | 10,808 | 83.6 | 2,114 | 16.4 | 709 | 848 |
| Montana | 423 | 10,373 | 8,877 | 85.6 | 1,496 | 14.4 | 918 | 1,072 |
| Nevada | 454 | 17,902 | 16,069 | 89.8 | 1,833 | 10.2 | 618 | 688 |
| New Mexico | 345 | 18,949 | 16,240 | 85.7 | 2,709 | 14.3 | 818 | 955 |
| Utah | 487 | 19,200 | 16,364 | 85.2 | 2,835 | 14.8 | 598 | 702 |
| Wyoming | 373 | 5,008 | 4,446 | 88.8 | 561 | 11.2 | 835 | 940 |
| Pacific | 3,623 | 396,484 | 336,150 | 84.8 | 60,334 | 15.2 | 685 | 808 |
| Alaska | 450 | 5,883 | 5,247 | 89.2 | 636 | 10.8 | 765 | 857 |
| California | 1,541 | 277,575 | 234,530 | 84.5 | 43,045 | 15.5 | 638 | 755 |
| Hawaii | 363 | 12,061 | 10,195 | 84.5 | 1,866 | 15.5 | 791 | 936 |
| Oregon | 585 | 37,170 | 33,773 | 90.9 | 3,397 | 9.1 | 891 | 981 |
| Washington | 684 | 63,795 | 52,405 | 82.1 | 11,390 | 17.9 | 800 | 974 |

Table 57. Registered nurses in each State and geographic area, by activity status: 2008 (continued)

*Population data were based on July 1, 2008 estimates of resident population of States from Census Bureau Press Release NST-EST2008-01.

--Too few cases to report estimated number and estimated percent (fewer than 30 cases unweighted)

| State and geographic area | Number in sample | Total estimated number employed | Total estimated percent employed | Estimated number employed full-time | Estimated percent employed full-time | Estimated number employed part-time | Estimated percent employed part-time | Estimated full-time equivalent* | Estimated full-time equivalent per 100,000** |
|---------------------------|---------------------|------------------------------------|-------------------------------------|--|---|--|--------------------------------------|---------------------------------|--|
| United States | 28,554 | 2,596,599 | 100.0 | 1,937,405 | 74.6 | 659,194 | 25.4 | 2,267,002 | 746 |
| New England | 3,198 | 161,627 | 100.0 | 108,388 | 67.1 | 53,239 | 32.9 | 135,007 | 944 |
| Connecticut | 487 | 35,343 | 100.0 | 23,616 | 66.8 | 11,727 | 33.2 | 29,480 | 842 |
| Maine | 444 | 16,279 | 100.0 | 11,667 | 71.7 | 4,612 | 28.3 | 13,973 | 1,061 |
| Massachusetts | 757 | 77,574 | 100.0 | 51,172 | 66.0 | 26,402 | 34.0 | 64,373 | 991 |
| New Hampshire | 586 | 13,562 | 100.0 | 9,791 | 72.2 | 3,771 | 27.8 | 11,676 | 887 |
| Rhode Island | 445 | 11,531 | 100.0 | 7,539 | 65.4 | 3,992 | 34.6 | 9,535 | 907 |
| Vermont | 479 | 7,338 | 100.0 | 4,603 | 62.7 | 2,735 | 37.3 | 5,971 | 961 |
| Middle Atlantic | 2,604 | 377,938 | 100.0 | 281,527 | 74.5 | 96,411 | 25.5 | 329,732 | 812 |
| New Jersey | 616 | 73,634 | 100.0 | 54,042 | 73.4 | 19,592 | 26.6 | 63,838 | 735 |
| New York | 1,016 | 167,427 | 100.0 | 130,502 | 77.9 | 36,925 | 22.1 | 148,964 | 764 |
| Pennsylvania | 972 | 136,876 | 100.0 | 96,982 | 70.9 | 39,894 | 29.1 | 116,929 | 939 |
| South Atlantic | 5,617 | 481,050 | 100.0 | 381,517 | 79.3 | 99,534 | 20.7 | 431,283 | 739 |
| Delaware | 556 | 9,169 | 100.0 | 6,999 | 76.3 | 2,170 | 23.7 | 8,084 | 926 |
| District of Columbia | 417 | 11,053 | 100.0 | 9,061 | 82.0 | 1,992 | 18.0 | 10,057 | 1,699 |
| Florida | 981 | 150,321 | 100.0 | 122,499 | 81.5 | 27,822 | 18.5 | 136,410 | 744 |
| Georgia | 520 | 68,254 | 100.0 | 53,873 | 78.9 | 14,381 | 21.1 | 61,063 | 630 |
| Maryland | 753 | 50,159 | 100.0 | 37,765 | 75.3 | 12,393 | 24.7 | 43,962 | 780 |
| North Carolina | 701 | 82,104 | 100.0 | 65,997 | 80.4 | 16,107 | 19.6 | 74,051 | 803 |
| South Carolina | 427 | 37,472 | 100.0 | 30,098 | 80.3 | 7,373 | 19.7 | 33,785 | 754 |
| Virginia | 825 | 54,988 | 100.0 | 41,911 | 76.2 | 13,077 | 23.8 | 48,450 | 624 |
| West Virginia | 437 | 17,529 | 100.0 | 13,312 | 75.9 | 4,217 | 24.1 | 15,421 | 850 |
| East South Central | 1,725 | 171,020 | 100.0 | 135,139 | 79.0 | 35,881 | 21.0 | 153,079 | 846 |
| Alabama | 448 | 41,488 | 100.0 | 33,559 | 80.9 | 7,929 | 19.1 | 37,524 | 805 |
| Kentucky | 431 | 41,520 | 100.0 | 31,939 | 76.9 | 9,581 | 23.1 | 36,730 | 860 |
| Mississippi | 380 | 27,414 | 100.0 | 23,434 | 85.5 | 3,980 | 14.5 | 25,424 | 865 |
| Tennessee | 466 | 60,597 | 100.0 | 46,207 | 76.3 | 14,391 | 23.7 | 53,402 | 859 |
| West South Central | 2,311 | 249,889 | 100.0 | 213,406 | 85.4 | 36,483 | 14.6 | 231,647 | 657 |
| Arkansas | 428 | 22,443 | 100.0 | 19,129 | 85.2 | 3,314 | 14.8 | 20,786 | 728 |
| Louisiana | 433 | 36,859 | 100.0 | 31,808 | 86.3 | 5,051 | 13.7 | 34,334 | 778 |
| Oklahoma | 410 | 27,256 | 100.0 | 23,008 | 84.4 | 4,248 | 15.6 | 25,132 | 690 |
| Texas | 1,040 | 163,331 | 100.0 | 139,461 | 85.4 | 23,870 | 14.6 | 151,396 | 622 |

Table 58. State and geographic area of registered nurses, by employment status in principal nursing position: 2008

Footnotes at end of table.

| | | | | | | | | | Estimated full-time |
|---------------------------|-----------|-----------------|------------------|--------------------|--------------------|--------------------|--------------------|---------------------|---------------------|
| | Number in | Total estimated | Total estimated | Estimated number | Estimated percent | Estimated number | Estimated percent | Estimated full-time | equivalent per |
| State and geographic area | sample | number employed | percent employed | employed full-time | employed full-time | employed part-time | employed part-time | equivalent* | 100,000** |
| | 0.000 | 405.040 | 400.0 | 200.000 | 74.0 | 400.040 | 00.0 | 070 047 | 004 |
| East North Central | 3,280 | 435,942 | 100.0 | 309,693 | 71.0 | 126,249 | 29.0 | 372,817 | 804 |
| Illinois | 765 | 111,939 | 100.0 | 78,700 | 70.3 | 33,239 | 29.7 | 95,319 | 739 |
| Indiana | 484 | 58,089 | 100.0 | 42,207 | 72.7 | 15,881 | 27.3 | 50,148 | 786 |
| Michigan | 613 | 88,449 | 100.0 | 63,171 | 71.4 | 25,278 | 28.6 | 75,810 | 758 |
| Ohio | 895 | 120,642 | 100.0 | 87,280 | 72.3 | 33,363 | 27.7 | 103,961 | 905 |
| Wisconsin | 523 | 56,822 | 100.0 | 38,334 | 67.5 | 18,489 | 32.5 | 47,578 | 845 |
| West North Central | 3,410 | 221,392 | 100.0 | 155,961 | 70.4 | 65,431 | 29.6 | 188,677 | 936 |
| lowa | 464 | 33,929 | 100.0 | 25,788 | 76.0 | 8,142 | 24.0 | 29,859 | 994 |
| Kansas | 469 | 27,032 | 100.0 | 21,322 | 78.9 | 5,709 | 21.1 | 24,177 | 863 |
| Minnesota | 558 | 58,391 | 100.0 | 32,831 | 56.2 | 25,560 | 43.8 | 45,611 | 874 |
| Missouri | 585 | 64,064 | 100.0 | 47,209 | 73.7 | 16,854 | 26.3 | 55,637 | 941 |
| Nebraska | 446 | 19,086 | 100.0 | 14,727 | 77.2 | 4,360 | 22.8 | 16,907 | 948 |
| North Dakota | 474 | 8,169 | 100.0 | 5,935 | 72.6 | 2,234 | 27.4 | 7,052 | 1,099 |
| South Dakota | 414 | 10,720 | 100.0 | 8,149 | 76.0 | 2,571 | 24.0 | 9,435 | 1,173 |
| Mountain | 3,281 | 161,591 | 100.0 | 120,919 | 74.8 | 40,673 | 25.2 | 141,255 | 648 |
| Arizona | 472 | 49,325 | 100.0 | 37,834 | 76.7 | 11,491 | 23.3 | 43,580 | 670 |
| Colorado | 565 | 39,461 | 100.0 | 28,024 | 71.0 | 11,437 | 29.0 | 33,743 | 683 |
| Idaho | 420 | 10,808 | 100.0 | 7,962 | 73.7 | 2,846 | 26.3 | 9,385 | 616 |
| Montana | 363 | 8,877 | 100.0 | 6,580 | 74.1 | 2,297 | 25.9 | 7,729 | 799 |
| Nevada | 413 | 16,069 | 100.0 | 13,076 | 81.4 | 2,992 | 18.6 | 14,573 | 560 |
| New Mexico | 298 | 16,240 | 100.0 | 12,172 | 75.0 | 4,068 | 25.0 | 14,206 | 716 |
| Utah | 418 | 16,364 | 100.0 | 11,835 | 72.3 | 4,530 | 27.7 | 14,100 | 515 |
| Wyoming | 332 | 4,446 | 100.0 | 3,435 | 77.3 | 1,011 | 22.7 | 3,940 | 740 |
| Pacific | 3,128 | 336,150 | 100.0 | 230,857 | 68.7 | 105,293 | 31.3 | 283,503 | 578 |
| Alaska | 404 | 5,247 | 100.0 | 3,911 | 74.5 | 1,336 | 25.5 | 4,579 | 667 |
| California | 1,312 | 234,530 | 100.0 | 163,664 | 69.8 | 70,866 | 30.2 | 199,097 | 542 |
| Hawaii | 308 | 10,195 | 100.0 | 7,742 | 75.9 | 2,453 | 24.1 | 8,968 | 696 |
| Oregon | 539 | 33,773 | 100.0 | 22,044 | 65.3 | 11,729 | 34.7 | 27,908 | 736 |
| Washington | 565 | 52,405 | 100.0 | 33,496 | 63.9 | 18,909 | 36.1 | 42,951 | 656 |

Table 58. State and geographic area of registered nurses, by employment status in principal nursing position: 2008 (continued)

*Nurses working full-time plus one-half of working part-time. **Population data were based on July 1, 2008 estimates of resident population of States from Census Bureau Press Release NST-EST2008-01. NOTE: Estimated numbers may not equal total, and percents may not add to 100, because of rounding.

| | | | | | Geograp | hic area | | | | |
|--|--------------|---------|----------|----------|---------|----------|------------|---------|----------|---------|
| | | | | | East | West | | West | | |
| | Total United | New | Middle | South | South | South | East North | North | | |
| Employment setting of principal nursing position | States* | England | Atlantic | Atlantic | Central | Central | Central | Central | Mountain | Pacific |
| | | | | | | | | | | |
| Estimated registered nurses in area | 2,596,599 | 161,627 | 377,938 | 481,050 | 171,020 | 249,889 | 435,942 | 221,392 | 161,591 | 336,150 |
| | 4 004 004 | 04.054 | 000 700 | 004 400 | 400 500 | 450.070 | 000 400 | 407.004 | 00.000 | 045 404 |
| Hospital | 1,601,831 | 91,254 | 223,786 | 301,463 | 109,506 | 156,273 | 268,423 | 137,024 | 98,668 | 215,434 |
| Nursing home/extended care | 135,514 | 14,747 | 25,033 | 21,643 | 7,236 | 7,859 | 27,356 | 15,313 | 7,688 | 8,640 |
| Academic education | 98,268 | 5,187 | 12,942 | 18,883 | 6,877 | 8,756 | 17,447 | 7,997 | 6,384 | 13,795 |
| Home health | 165,697 | 13,767 | 27,298 | 30,842 | 10,767 | 21,597 | 26,478 | 12,732 | 9,909 | 12,307 |
| Community/public health** | 116,051 | 6,347 | 14,785 | 22,992 | 7,765 | 9,172 | 18,898 | 11,362 | 6,971 | 17,760 |
| School health | 84,418 | 9,376 | 21,023 | 12,848 | 3,174 | 9,533 | 9,445 | 6,716 | 4,834 | 7,468 |
| Ambulatory care | 270,556 | 12,657 | 36,111 | 46,496 | 17,225 | 24,130 | 50,209 | 21,305 | 18,677 | 43,746 |
| Insurance claims/benefits/utilization review | 49,441 | 3,125 | 7,618 | 9,392 | 3,425 | 4,419 | 8,216 | 3,007 | 2,998 | 7,241 |
| Other | 51,947 | 3,445 | 6,080 | 12,205 | 3,539 | 5,974 | 6,813 | 4,184 | 3,741 | 5,966 |

Table 59. Employment setting of principal nursing position, by geographic area: 2008

*Includes an estimated 22,875 RNs for whom employment setting was not known. **Community/public health includes occupational health. NOTE: Estimated numbers may not equal totals because of rounding.

| | F | Full-time staff nurses | |
|---|---|---|--|
| Geographic area of employment | Number in sample | Estimated number* | Annual earnings (dollars) |
| Total | 12,808 | 1,212,525 | 61,706 |
| New England Middle Atlantic South Atlantic East South Central West South Central East North Central West North Central Mountain Pacific | 1,235 1,172 2,598 831 1,198 1,386 1,490 1,525 1,373 | 66,117 175,327 235,974 83,316 135,373 192,610 96,531 77,386 149,890 | 65,799 66,228 59,022 53,514 59,612 57,356 52,627 61,823 76,665 |

Table 60. Average annual earnings for registered nurses working full-time as staff nurses in their principal nursing position, by geographical area: 2008

*Excludes an estimated 467 full-time staff nurses who did not report annual earnings. NOTE: Estimated numbers may not equal totals because of rounding.

| | | | Highest nursing or nursing-related education preparation | | | | | | | |
|---------------------------|-----------|-----------|--|------------|-----------|------------|------------|------------|-----------|------------|
| | | | | | Estimated | Estimated | | | Estimated | Estimated |
| | | Total | Estimated | Estimated | number of | percent of | Estimated | Estimated | number of | percent of |
| | Number in | estimated | number of | percent of | associate | associate | number of | percent of | master's/ | master's/ |
| State and geographic area | sample | number* | diploma | diploma | degree | degree | bachelor's | bachelor's | doctorate | doctorate |
| East North Central | 3,280 | 435,942 | 61,780 | 14.2 | 159,091 | 36.5 | 165,053 | 37.9 | 49,491 | 11.4 |
| Illinois | 765 | 111,939 | 16,922 | 15.1 | 36,073 | 32.3 | 43,374 | 38.8 | 15,392 | 13.8 |
| Indiana | 484 | 58,089 | 5,448 | 9.4 | 24,395 | 42.0 | 22,464 | 38.7 | 5,782 | 10.0 |
| Michigan | 613 | 88,449 | 9,194 | 10.4 | 36,130 | 40.8 | 33,021 | 37.3 | 10,104 | 11.4 |
| Ohio | 895 | 120,642 | 23,065 | 19.2 | 43,106 | 35.8 | 41,705 | 34.7 | 12,469 | 10.4 |
| Wisconsin | 523 | 56,822 | 7,150 | 12.6 | 19,387 | 34.1 | 24,489 | 43.1 | 5,744 | 10.1 |
| West North Central | 3,410 | 221,392 | 28,462 | 12.9 | 81,047 | 36.7 | 86,827 | 39.3 | 24,771 | 11.2 |
| lowa | 464 | 33,929 | 5,260 | 15.5 | 15,721 | 46.3 | 9,709 | 28.6 | 3,239 | 9.5 |
| Kansas | 469 | 27,032 | 3,771 | 14.0 | 9,947 | 36.8 | 10,685 | 39.5 | 2,628 | 9.7 |
| Minnesota | 558 | 58,391 | 5,661 | 9.7 | 22,948 | 39.5 | 22,987 | 39.6 | 6,518 | 11.2 |
| Missouri | 585 | 64,064 | 7,499 | 11.7 | 22,021 | 34.4 | 25,666 | 40.1 | 8,878 | 13.9 |
| Nebraska | 446 | 19,086 | 3,838 | 20.1 | 4,427 | 23.2 | 9,040 | 47.4 | 1,782 | 9.3 |
| North Dakota | 474 | 8,169 | 1,271 | 15.6 | 1,493 | 18.3 | 4,567 | 56.0 | 831 | 10.2 |
| South Dakota | 414 | 10,720 | 1,162 | 10.8 | 4,490 | 41.9 | 4,172 | 38.9 | 897 | 8.4 |
| Mountain | 3,281 | 161,591 | 13,116 | 8.1 | 61,196 | 38.0 | 64,507 | 40.0 | 22,424 | 13.9 |
| Arizona | 472 | 49,325 | 4,889 | 10.0 | 19,638 | 40.1 | 17,039 | 34.8 | 7,455 | 15.2 |
| Colorado | 565 | 39,461 | 3,809 | 9.7 | 11,010 | 27.9 | 18,128 | 45.9 | 6,514 | 16.5 |
| Idaho | 420 | 10,808 | | | 5,220 | 48.3 | 3,958 | 36.6 | 1,180 | 10.9 |
| Montana | 363 | 8,877 | 699 | 7.9 | 2,735 | 30.8 | 4,424 | 49.8 | 1,019 | 11.5 |
| Nevada | 413 | 16,069 | 1,587 | 9.9 | 5,653 | 35.2 | 6,355 | 39.6 | 2,445 | 15.2 |
| New Mexico | 298 | 16,240 | | | 7,171 | 44.2 | 6,196 | 38.2 | 2,034 | 12.5 |
| Utah | 418 | 16,364 | | | 7,426 | 45.4 | 7,074 | 43.2 | 1,313 | 8.0 |
| Wyoming | 332 | 4,446 | | | 2,344 | 52.9 | 1,333 | 30.1 | 463 | 10.5 |
| Pacific | 3,128 | 336,150 | 29,271 | 8.7 | 121,665 | 36.2 | 135,468 | 40.3 | 49,477 | 14.7 |
| Alaska | 404 | 5,247 | 403 | 7.7 | 1,639 | 31.2 | 2,345 | 44.7 | 860 | 16.4 |
| California | 1,312 | 234,530 | 19,970 | 8.5 | 82,739 | 35.3 | 94,673 | 40.4 | 36,879 | 15.7 |
| Hawaii | 308 | 10,195 | | | 2,654 | 26.0 | 5,345 | 52.4 | 1,540 | 15.1 |
| Oregon | 539 | 33,773 | 2,842 | 8.4 | 13,226 | 39.2 | 13,829 | 40.9 | 3,876 | 11.5 |
| Washington | 565 | 52,405 | 5,400 | 10.3 | 21,408 | 40.9 | 19,275 | 36.8 | 6,321 | 12.1 |

Table 61. Registered nurses employed in nursing in each State and geographic area, by highest nursing or nursing-related education: 2008 (continued)

*Includes an estimated 3,297 RNs for whom highest nursing education was not known.

--Too few cases to report estimated number and estimated percent (fewer than 30 cases unweighted).

| | | | | | Geograp | hic area | | | | |
|--------------------------------------|--------------|---------|----------|----------|---------|----------|------------|---------|----------|---------|
| | | | | | East | West | | West | | |
| | Total United | New | Middle | South | South | South | East North | North | | |
| Age group | States | England | Atlantic | Atlantic | Central | Central | Central | Central | Mountain | Pacific |
| Estimated registered nurses in area | 3,063,162 | 187,542 | 462,263 | 576,337 | 195,713 | 290,304 | 516,080 | 251,281 | 187,159 | 396,484 |
| Less than 30 | 9.4 | 7.9 | 6.9 | 9.4 | 12.0 | 10.7 | 9.2 | 12.6 | 10.5 | 8.6 |
| 30-34 | 9.2 | 7.0 | 8.6 | 9.4 | 11.4 | 10.8 | 8.6 | 9.6 | 8.6 | 9.3 |
| 35-39 | 10.8 | 9.3 | 10.2 | 11.0 | 13.2 | 13.0 | 10.7 | 9.2 | 10.3 | 10.8 |
| 40-44 | 11.4 | 11.5 | 10.9 | 12.4 | 12.4 | 11.1 | 11.6 | 11.7 | 10.2 | 10.5 |
| 45-49 | 14.4 | 15.1 | 14.4 | 13.7 | 14.9 | 14.2 | 14.8 | 15.3 | 14.8 | 13.9 |
| 50-54 | 16.2 | 16.8 | 17.6 | 15.3 | 14.0 | 16.2 | 16.5 | 16.1 | 15.7 | 16.5 |
| 55-59 | 13.0 | 13.8 | 12.9 | 12.7 | 11.1 | 12.0 | 13.6 | 12.2 | 14.2 | 13.8 |
| 60-64 | 8.1 | 10.1 | 8.8 | 8.8 | 6.6 | 7.1 | 7.1 | 7.2 | 9.0 | 8.6 |
| 65 years and over | 7.4 | 8.6 | 9.7 | 7.3 | 4.5 | 5.0 | 7.9 | 6.0 | 6.8 | 7.9 |
| Average age | 47.0 | 48.5 | 48.4 | 46.9 | 44.6 | 45.5 | 47.2 | 45.9 | 47.0 | 47.6 |
| Median age | 48 | 49 | 49 | 48 | 45 | 46 | 48 | 47 | 48 | 49 |
| Estimated registered nurses employed | | | | | | | | | | |
| in nursing in area | 2,596,599 | 161,627 | 377,938 | 481,050 | 171,020 | 249,889 | 435,942 | 221,392 | 161,591 | 336,150 |
| Less than 30 | 10.6 | 8.7 | 7.9 | 10.7 | 13.5 | 11.4 | 10.6 | 13.9 | 11.8 | 9.7 |
| 30-34 | 9.9 | 7.9 | 9.4 | 10.0 | 11.8 | 11.7 | 9.4 | 10.1 | 9.0 | 10.0 |
| 35-39 | 11.7 | 10.3 | 11.7 | 11.6 | 13.1 | 14.2 | 11.3 | 9.7 | 10.8 | 11.8 |
| 40-44 | 12.1 | 12.0 | 11.9 | 13.2 | 13.2 | 11.6 | 12.4 | 11.9 | 10.5 | 11.2 |
| 45-49 | 15.3 | 15.8 | 15.6 | 14.6 | 15.7 | 14.5 | 15.8 | 16.2 | 15.9 | 14.7 |
| 50-54 | 16.7 | 17.3 | 18.9 | 15.6 | 14.3 | 16.3 | 17.0 | 16.7 | 16.3 | 17.1 |
| 55-59 | 13.0 | 13.8 | 13.3 | 12.7 | 10.4 | 11.9 | 13.8 | 12.3 | 14.3 | 13.7 |
| 60-64 | 6.8 | 9.4 | 7.2 | 7.5 | 5.1 | 5.9 | 5.7 | 6.2 | 7.8 | 7.′ |
| 65 years and over | 3.9 | 4.8 | 4.1 | 4.2 | 2.9 | 2.5 | 4.0 | 3.0 | 3.7 | 4.7 |
| Average age | 45.5 | 47.1 | 46.4 | 45.5 | 43.6 | 44.3 | 45.5 | 44.7 | 45.8 | 46.1 |
| Median age | 46 | 48 | 47 | 46 | 44 | 45 | 46 | 46 | 47 | 47 |

Table 62. Age distribution and average age of registered nurses, by geographic areafor all registered nurses and registered nurses employed in nursing: 2008

| | Geographic area | | | | | | | | | |
|---|------------------------|-------------|--------------------|-------------------|-----------------------|-----------------------|-----------------------|-----------------------|----------|---------|
| Racial/ethnic background | Total United States | New England | Middle Atlantic | South Atlantic | East South Central | West South Central | East North Central | West North Central | Mountain | Pacific |
| Estimated registered nurses in area | 3,063,162 | 187,542 | 462,263 | 576,337 | 195,713 | 290,304 | 516,080 | 251,281 | 187,159 | 396,484 |
| Racial/ethnic background | | | | | | | | | | |
| White (non-Hispanic) | 83.2 | 93.1 | 82.9 | 80.7 | 89.0 | 75.4 | 89.4 | 94.2 | 85.8 | 69.3 |
| Black/African American (non-Hispanic) | 5.4 | 2.0 | 6.2 | 9.5 | 7.7 | 8.8 | 3.7 | 1.6 | 1.3 | 3.0 |
| Asian (non-Hispanic) | 5.5 | 2.1 | 7.3 | 4.2 | | 5.6 | 3.2 | 1.0 | 3.7 | 16.0 |
| Native Hawaiian/Pacific Islander (non-Hispanic) | 0.3 | | | | | | | | | 1.3 |
| American Indian/Alaska Native (non-Hispanic) | 0.3 | | | | | | | | 1.3 | |
| Hispanic/Latino (any race) | 3.6 | 1.4 | 2.2 | 3.6 | | 7.8 | 1.9 | 1.3 | 6.1 | 6.7 |
| Two or more races (non-Hispanic) | 1.7 | 1.2 | 1.1 | 1.5 | 1.4 | 1.7 | 1.5 | 1.7 | 1.3 | 3.5 |

Table 63. Racial/ethnic background of registered nurses, by geographic area: 2008

--Too few cases to report estimated percent (fewer than 30 cases unweighted).

| | Total estimated number* | Metro area employed in nursing | Metro area not employed in nursing | Nonmetro area employed in nursing | Nonmetro area not employed in nursing |
|--------------------|----------------------------|--------------------------------------|--|---|---|
| Total | 3,063,162 | 1,971,644 | 358,212 | 501,480 | 95,927 |
| New England | 187,542 | 103,196 | 19,386 | 46,012 | 6,029 |
| Middle Atlantic | 462,263 | 319,350 | 72,102 | 39,995 | 10,450 |
| South Atlantic | 576,337 | 384,972 | 74,898 | 72,148 | 18,143 |
| East South Central | 195,713 | 116,486 | 15,986 | 49,539 | 7,435 |
| West South Central | 290,304 | 198,862 | 30,379 | 37,606 | 8,947 |
| East North Central | 516,080 | 302,975 | 56,922 | 116,369 | 21,604 |
| West North Central | 251,281 | 146,712 | 19,925 | 66,131 | 9,515 |
| Mountain | 187,159 | 124,230 | 19,079 | 31,103 | 5,800 |
| Pacific | 396,484 | 274,861 | 49,535 | 42,576 | 8,005 |

 Table 64. Geographic location of registered nurses, by employment status: 2008

*Includes an estimated 135,899 RNs for whom metro/nonmetro area status was not known.

NOTE: Estimated numbers may not equal totals because of rounding.

| Predominant countries where internationally | Number in | Total estimated | Total estimated | sample | Estimated number | Estimated percent employed | Estimated percent of all employed internationally |
|--|-----------|--------------------|--------------------|------------------------|---------------------|----------------------------------|---|
| educated nurses received initial nursing education | sample | number | percent | employed in nursing | employed in nursing | in nursing | educated nurses |
| Total (excluding U.S. territories) | 1,461 | 165,539 | 100.0 | 1,296 | 146,097 | 88.3 | 100.0 |
| Asia/Australia | 937 | 114,299 | 69.0 | 846 | 102,792 | 89.9 | 70.4 |
| Philippines | 681 | 82,988 | 50.1 | 620 | 76,021 | 91.6 | 52.0 |
| India | 120 | 15,827 | 9.6 | 113 | 14,593 | 92.2 | 10.0 |
| Korea | 36 | 4,359 | 2.6 | 28 | | | |
| Other Asia/Australia | 100 | 11,124 | 6.7 | 85 | 8,983 | 80.8 | 6.1 |
| North/South America/Caribbean | 290 | 28,942 | 17.5 | 254 | 25,140 | 86.9 | 17.2 |
| Canada | 218 | 19,619 | 11.9 | 192 | 17,152 | 87.4 | 11.7 |
| Other North/South America/Caribbean | 72 | 9,322 | 5.6 | 62 | 7,988 | 85.7 | 5.5 |
| Europe | 150 | 15,410 | 9.3 | 115 | 11,573 | 75.1 | 7.9 |
| United Kingdom | 95 | 9,921 | 6.0 | 74 | 7,206 | 72.6 | 4.9 |
| Other Europe | 55 | 5,489 | 3.3 | 41 | 4,366 | 79.5 | 3.0 |
| Africa | 84 | 6,888 | 4.2 | 81 | 6,592 | 95.7 | 4.5 |
| Nigeria | 36 | 3,409 | 2.1 | 33 | 3,113 | 91.3 | 2.1 |
| Other Africa | 48 | 3,479 | 2.1 | 48 | 3,479 | 100.0 | 2.4 |
| U.S. territories* | 39 | 4,696 | | 37 | 4,405 | 93.8 | |

Table 65. Predominant countries where internationally educated nurses received initial nursing education, by employment status: 2008

*Counts for U.S. territories are provided to complement the information for internationally educated nurses, but are not included in the column totals.

--Too few cases to report estimated number and estimated percent (fewer than 30 cases unweighted).

| State employing IENs | Number in sample | Estimated number IENs employed in nursing | Estimated percent IENs employed in nursing | Percent of employed IENs in the State |
|----------------------|---------------------|---|--|---|
| Total | 1,296 | 146,097 | 100.0 | 5.6 |
| California | 202 | 38,615 | 26.4 | 16.5 |
| New York | 108 | 16,803 | 11.5 | 10.0 |
| Texas | 91 | 14,922 | 10.2 | 9.1 |
| Florida | 86 | 14,009 | 9.6 | 9.3 |
| New Jersey | 70 | 8,552 | 5.9 | 11.6 |
| Illinois | 51 | 7,947 | 5.4 | 7.1 |
| Maryland | 80 | 4,019 | 2.8 | 8.0 |
| Virginia | 53 | 2,215 | 1.5 | 4.0 |
| Nevada | 50 | 1,971 | 1.3 | 12.3 |
| All other states | 505 | 37,043 | 25.4 | 2.4 |

Table 66. Predominant States employing internationally educated nurses (IENs): 2008

| Table 67. Employment setting of internationally educated nurses (IENs), |
|---|
| by selected job titles: 2008 |

| Employment setting of IENs | Total estimated number of employed IENs* | Total estimated percent of employed IENs | Estimated number of IENs employed as staff nurse | Estimated percent of IENs employed as staff nurse | Estimated number of IENs employed in all other positions | Estimated percent of IENs employed in all other positions |
|-------------------------------------|--|--|--|---|--|---|
| Total | 146,097 | 100.0 | 113,274 | 100.0 | 30,461 | 100.0 |
| Hospital | 105,320 | 72.1 | 87,245 | 77.0 | 16,561 | 54.4 |
| Nursing home/extended care facility | 9,559 | 6.5 | 6,163 | 5.4 | 3,359 | 11.0 |
| Academic education program | 9,490 | 6.5 | 6,694 | 5.9 | | |
| Home health setting | 5,785 | 4.0 | | | | |
| Public/community health** | 4,985 | 3.4 | 3,835 | 3.4 | | |
| Ambulatory care setting | 6,276 | 4.3 | 4,323 | 3.8 | | |
| Other *** | 3,042 | 2.1 | | | | |

*Includes an estimated 15 IENs for whom employment setting was not known.

Public/community health includes school health and occupational health. *Other includes insurance, benefits, and utilization review.

--Too few cases to report estimated number and estimated percent (fewer than 30 cases unweighted).

Appendix B

Survey Methodology

Appendix B: Survey Methodology

B.1 Sampling

Since the inception of the National Sample Survey of Registered Nurses (NSSRN), the sample of Registered Nurses (RNs) was based on a nested cluster design, clustering RNs based on last names. This last name cluster design is referred to as the alpha-segment design. While the alpha-segment design helped produce good estimates of the nursing workforce over the years, it has some shortcomings. Since people of the same racial/ethnic background tended to be clustered within the same alphabetic clusters of last names, the clustered design resulted in larger than desired variation for some estimates, most notably the racial and ethnic composition of the RN workforce. Additionally, constructing the alpha-segment frame and implementing the sample design were tedious and time-consuming tasks. Sorting to achieve implicit stratification was not feasible, and the alpha-segment design made adjustments to account for differential nonresponse a bit more cumbersome.

For the 2008 NSSRN, the Health Resources and Service Administration (HRSA) implemented a sample design based on independent systematic random samples selected from State-based strata with equal probability of selection within each stratum. An equal probability systematic random sample can be thought of as randomly selecting a record from the beginning of a list with probability 1/n and then every nth record after that until all records on the list have had a chance of being selected. This kind of design was not feasible in 1976, but technological advances and the advent of electronic recordkeeping have largely eliminated the barriers that existed previously. This design was straightforward to implement and eliminated the clustering that could contribute undesirably to the variability of sample estimates, particularly for those associated with race/ethnicity.

B.1.1 Sample Selection

The 2008 NSSRN included a sample of 55,151 RN records.

Systematic random samples of RNs were selected from each of the sample frames established. A sample frame was developed for each of the 50 States (and the District of Columbia) of those RNs with currently active licenses in the State. The sample frames were developed from listings obtained from each of the State licensure boards.

State frames were first partitioned into explicit strata for sample selection purposes. RNs appearing in sample frame of licensed nurses for more than one State had multiple chances of selection for the 2008 NSSRN. Hence multi-State strata were formed for several groups of States where a relatively high degree of interstate commuting was expected. Probability matching was used to form such strata, using only a single record for those RNs found on listings from two or more States in a given group (for example, a stratum was formed for all RNs with licenses found in New York and New Jersey, each such RN having only a single record in that stratum). Following are the groupings of States where such strata were formed:

- District of Columbia, Maryland, and Virginia
- New York, New Jersey, and Pennsylvania
- Arizona, California, and Nevada
- Massachusetts, New Hampshire, and Vermont
- Delaware, Pennsylvania, and New Jersey

For most of the remaining States, four geographic sampling strata were used: one stratum for the State itself, and three additional, much smaller, strata. These three strata were for RNs whose respective residence was listed as Alaska, Hawaii, or New Mexico. These additional strata were created because nurses working in Federal facilities need not be licensed in the State in which they work, and such RNs are found at disproportionately high rates in Alaska, Hawaii, and New Mexico. When sampling for Alaska, Hawaii, and New Mexico, only three total strata were used. Alaska, Hawaii, and New Mexico strata were also formed for the "multi-State strata" groups listed above.

Prior to sample selection, the sample frame for each stratum was sorted on variables such as age group, ZIP Code, or other variables. The choice of variables depended on the variables available on a State's frame and which among them were of greatest analytic interest or were related to variables of analytic interest. If a frame was sorted only on one variable, say ZIP Code, then the sample would have a proportionate or approximately proportionate distribution across the ZIP Codes appearing on the frame. If a second variable was added, say age group, then within each ZIP Code a proportionate distribution across age groups would be obtained. Thus, sorting was done to help achieve a proportionate or approximately proportionate sample distribution across important variables related to analytic variables of interest, helping to increase the precision of the Survey estimates.

B.1.2 Final Sample Allocation

Tables B1-1 and B1-2 show the final allocation of sampled RNs across the States where the multi-State strata mentioned earlier were not established. Table B1-1 is arranged in the order of size of frame from largest to smallest while Table B1-2 is arranged alphabetically. Table B1-3 shows the final allocation of RNs across the States where multiple strata were established associated with neighboring States.

Four columns of numbers are provided for each State or strata: the 2008 State and strata frame size after unduplication; 2008 sample size after assigning Alaska, Hawaii, and Mexico samples from different State or strata back to Alaska, Hawaii, and New Mexico; targeted sample size used to set the sampling rate for the State or stratum; and the difference between the targeted sample size and the actual sample size.¹ Note that for the District of Columbia/Maryland/Virginia strata it was decided to form strata taking into account whether or not the RN lived in the DC metro area regardless of which records matched. So summing across strata does not produce unduplicated totals for the District of Columbia, Maryland, or Virginia separately. Had the Nurse Licensure Compact (the Compact) State specifications been applied perfectly, there would have been no RNs found on both the Virginia and Maryland listings. However, there were a few (perhaps due to very recent residential moves from one State to the other) where synchronization of the licensing to eliminate the earlier compact license had not yet been achieved.

B.2 Sample Performance

Response rates serve to help assess the quality or unbiasedness of survey results and are often reported both unweighted and weighted. An unweighted response rate provides an indication of the degree to which the RNs actually selected for the sample decided to participate in the survey. A weighted response rate provides an estimate of the propensity to respond to the survey among the members of the target population generally. A weighted response rate accounts for factors such as oversampling. For example, for the 2008 NSSRN, RNs have been allocated to the smaller States at disproportionately high rates (i.e., they have been "oversampled") in order to help achieve adequate precision for State level estimates for all States.

¹ Departures arose due to the relatively few RNs in each State with a residence address of Alaska, Hawaii, or New Mexico

| | | | | Difference |
|------------------------|------------------|-----------------------|-----------------|--------------------|
| | | | | between final and |
| | | 2008 sample sizes | | targeted sample |
| | | after assigning AK, | | sizes (differences |
| | | HI, and NM samples | 2008 allocation | due to sampling |
| | Final 2008 frame | from different States | on which | RNs living in AK, |
| Single State | size after | or strata to AK, HI, | sampling rates | HI, and NM |
| samples | unduplication | and NM | were based | separately) |
| Florida | 221,482 | 1,896 | 1,900 | -4 |
| Texas | 203,595 | 1,796 | 1,800 | -4 |
| Illinois | 160,909 | 1,598 | 1,600 | -2 |
| Ohio | 153,487 | 1,449 | 1,450 | -1 |
| Michigan | 126,007 | 1,198 | 1,200 | -2 |
| North Carolina | 105,317 | 1,100 | 1,100 | 0 |
| Georgia | 99,795 | 1,099 | 1,100 | -1 |
| Missouri | 89,035 | 899 | 900 | -1 |
| Indiana | 80,779 | 850 | 850 | 0 |
| Minnesota | 76,708 | 849 | 850 | -1 |
| Washington | 76,364 | 900 | 900 | 0 |
| Wisconsin | 74,891 | 849 | 850 | -1 |
| Tennessee | 72,754 | 800 | 800 | 0 |
| Alabama | 55,862 | 849 | 850 | -1 |
| Colorado | 53,871 | 896 | 900 | -4 |
| Connecticut | 52,898 | 999 | 1,000 | -1 |
| Kentucky | 51,229 | 749 | 750 | -1 |
| South Carolina | 47,808 | 849 | 850 | -1 |
| Louisiana | 47,419 | 799 | 800 | -1 |
| Oregon | 42,995 | 845 | 850 | -5 |
| Iowa | 42,419 | 800 | 800 | 0 |
| Kansas | 39,801 | 899 | 900 | -1 |
| Oklahoma | 38,914 | 946 | 950 | -4 |
| Mississippi | 38,406 | 949 | 950 | -1 |
| Arkansas | 30,677 | 899 | 900 | -1 |
| New Mexico | 24,658 | 882 | 800 | 82 |
| West Virginia | 24,184 | 949 | 950 | -1 |
| Maine | 23,210 | 898 | 900 | -2 |
| Utah | 23,019 | 797 | 800 | -3 |
| Nebraska | 22,358 | 699 | 700 | -1 |
| Hawaii | 17,155 | 964 | 900 | 64 |
| Rhode Island | 16,706 | 999 | 1,000 | -1 |
| Idaho | 15,059 | 798 | 800 | -2 |
| Montana | 13,835 | 793 | 800 | -7 |
| South Dakota | 13,833 | 748 | 750 | -7 -2 |
| North Dakota | 9,395 | 748 | 750 | -2 |
| Alaska | 9,393 8,478 | 980 | 800 | 180 |
| Wyoming | 7,826 | 980 794 | 800 | -6 |
| Subtotal single States | 2,303,132 | 36,813 | 36,550 | -6 263 |
| Suntotal single states | 2,303,132 | 30,813 | 30,000 | 203 |

Table B1-1.Sample information for States where the strata were based only on Alaska, Hawaii,
and New Mexico (sorted by frame size)

| Table B1-2. | Sample information for States where strata were based only on Alaska, Hawaii, and |
|-------------|---|
| | New Mexico (sorted alphabetically) |

| Single State | Final 2008 frame size after | 2008 sample sizes after assigning AK, HI, and NM samples from different States or strata to AK, HI, and | 2008 allocation on which sampling | Difference between final and targeted sample sizes (differences due to sampling RNs living in AK, HI, and NM |
|--------------------|-----------------------------------|---|-----------------------------------|---|
| Samples Alabama | | NM 849 | rates were based 850 | separately) -1 |
| Alaska | 55,862 | 980 | 800 | -1 180 |
| Arkansas | 8,478 30,677 | 899 | 900 | -1 |
| Colorado | 53,871 | 899 | 900 | -1 -4 |
| Connecticut | 52,898 | 999 999 | 1,000 | -4 -1 |
| Florida | 221,482 | 999 1,896 | 1,900 | -1 |
| Georgia | 99,795 | 1,099 | 1,100 | -4 -1 |
| Hawaii | 17,155 | 964 | 900 | 64 |
| Idaho | 15,059 | 798 | 800 | -2 |
| Illinois | 160,909 | 1,598 | 1,600 | -2 |
| Indiana | 80,779 | 850 | 850 | 0 |
| lowa | 42,419 | 800 | 800 | 0 |
| Kansas | 39,801 | 899 | 900 | -1 |
| Kentucky | 51,229 | 749 | 750 | -1 |
| Louisiana | 47,419 | 799 | 800 | -1 |
| Maine | 23,210 | 898 | 900 | -2 |
| Michigan | 126,007 | 1,198 | 1,200 | -2 |
| Minnesota | 76,708 | 849 | 850 | -1 |
| Mississippi | 38,406 | 949 | 950 | -1 |
| Missouri | 89,035 | 899 | 900 | -1 |
| Montana | 13,835 | 793 | 800 | -7 |
| Nebraska | 22,358 | 699 | 700 | -1 |
| New Mexico | 24,658 | 882 | 800 | 82 |
| North Carolina | 105,317 | 1,100 | 1,100 | 0 |
| North Dakota | 9,395 | 750 | 750 | 0 |
| Ohio | 153,487 | 1,449 | 1,450 | -1 |
| Oklahoma | 38,914 | 946 | 950 | -4 |
| Oregon | 42,995 | 845 | 850 | -5 |
| Rhode Island | 16,706 | 999 | 1,000 | -1 |
| South Carolina | 47,808 | 849 | 850 | -1 |
| South Dakota | 13,827 | 748 | 750 | -2 |
| Tennessee | 72,754 | 800 | 800 | 0 |
| Texas | 203,595 | 1,796 | 1,800 | -4 |
| Utah | 23,019 | 797 | 800 | -3 |
| Washington | 76,364 | 900 | 900 | 0 |
| West Virginia | 24,184 | 949 | 950 | -1 |
| Wisconsin | 74,891 | 849 | 850 | -1 |
| Wyoming | 7,826 | 794 | 800 | -6 |
| Subtotal single | 2,303,132 | 36,813 | 36,550 | 263 |
| States | | | | |

| Description | ns | Final 2008 frame size after unduplication | 2008 sample sizes after assigning AK, HI, NM samples from different States or strata to AK, HI, and NM | 2008 allocation on which sampling rates were based | Difference between final and targeted sample sizes (differences due to sampling RNs living in AK, HI, and NM separately) |
|--|----------------------------|--|---|---|---|
| Cumulative State totals or estimated Stat | e totals across Strata | | | | |
| California | CA | 349,916 | 3,257 | 3,260 | -3 |
| Arizona | AZ | 65,387 | 924 | 930 | -6 |
| Nevada | NV | 24,412 | 1,065 | 1,070 | -5 |
| Strata based on address in: | | | | | |
| Alaska | AK | 263 | 25 | х | x |
| Hawaii | HI | 672 | 34 | х | x |
| New Mexico | NM | 531 | 17 | Х | X |
| Strata descriptions | | | | | |
| Licensed in both NV and CA, no match with AZ | NVCA | 6,782 | 288 | 290 | -2 |
| Licensed in both NV and AZ, no match with CA | NVAZ | 572 | 20 | 20 | 0 |
| Licensed in NV, AZ, and CA | NVCAAZ | 742 | 20 | 20 | 0 |
| Remainder of those licensed in NV | NV_REMAINDER | 16,316 | 737 | 740 | -3 |
| Licensed in both CA and AZ, no match with NV | AZCA | 6,328 | 69 | 70 | -1 |
| Remainder of those licensed in AZ | AZ_REMAINDER | 57,745 | 815 | 820 | -5 |
| Remainder of those licensed in CA | CA_REMAINDER (half sample) | 168,032 | 2,880 | 2,880 | 0 |
| (after doubling CA remainder total to obtain appropriate estimate) | Subtotal est., CA NV AZ | 425,537 | 4,905 | 4,840 | -11 |
| Cumulative State totals or estimated Stat | e totals across Strata | | | | |
| New York | NY | 255,459 | 2,644 | 2,650 | -6 |
| Pennsylvania | PA | 204,371 | 1,887 | 1,890 | -3 |
| New Jersey | NJ | 111,117 | 1,418 | 1,420 | -2 |

Table B1-3. Sampling States where strata are based on RNs licensed in neighboring States as well

| Descriptio | ons | Final 2008 frame size after unduplication | 2008 sample sizes after assigning AK, HI, NM samples from different States or strata to AK, HI, and NM | 2008 allocation on which sampling rates were based | Difference between final and targeted sample sizes (differences due to sampling RNs living in AK, HI, and NM separately) |
|--|-------------------------|--|---|---|---|
| Strata based on address in | | | | | |
| Alaska | AK | 135 | 13 | х | Х |
| Hawaii | HI | 228 | 11 | х | х |
| New Mexico | NM | 295 | 9 | х | x |
| Strata descriptions | | | | | |
| Licensed in both NJ and NY, no match with PA | NJ_NY | 12,663 | 160 | 160 | 0 |
| Licensed in both NJ and PA, no match with NY | NJ_PA | 12,872 | 170 | 170 | 0 |
| Licensed in NJ, NY, and PA | NJ_NY_PA | 2,152 | 30 | 30 | 0 |
| Remainder of those licensed in NJ | NJ_REMAINDER | 83,430 | 1,058 | 1,060 | -2 |
| Licensed in both NY and PA, no match with NJ | NY_PA | 4,979 | 60 | 60 | 0 |
| Remainder of those licensed in PA | PA_REMAINDER | 184,368 | 1,627 | 1,630 | -3 |
| Remainder of those licensed in NY | NY_REMAINDER | 235,665 | 2,394 | 2,400 | -6 |
| | Subtotal, NY PA NJ | 536,787 | 5,532 | 5,510 | -11 |
| Cumulative State totals or estimated Sta | te totals across Strata | | | | |
| Massachusetts | MA | 111,845 | 1,382 | 1,385 | -3 |
| New Hampshire | NH | 18,447 | 1,029 | 1,030 | -1 |
| Vermont | VT | 15,494 | 958 | 960 | -2 |
| Strata based on address in | | | | | |
| Alaska | AK | 44 | 4 | х | x |
| Hawaii | HI | 59 | 3 | х | x |
| New Mexico | NM | 84 | 2 | х | x |

Table B1-3. Sampling States where strata are based on RNs licensed in neighboring States as well (continued)

| Description | S | Final 2008 frame size after unduplication | 2008 sample sizes after assigning AK, HI, NM samples from different States or strata to AK, HI, and NM | 2008 allocation on which sampling rates were based | Difference between final and targeted sample sizes (differences due to sampling RNs living in AK, HI, and NM separately) |
|--|------------------------|--|---|---|---|
| Strata descriptions | | - | | | |
| Licensed in VT, matched with record in MA, resides in VT, MA, or NH | VT_MTCH_MA_IN3 | 418 | 30 | 30 | 0 |
| Licensed in VT, matched with record in NH, resides in VT, MA, or NH | VT_MTCH_NH_IN3 | 863 | 70 | 70 | 0 |
| Licensed in VT, did not match with records from either MA and NH, resides in VT, MA, or NH | VT_MTCH_MANH_IN3 | 354 | 15 | 15 | 0 |
| Licensed in VT, matched with records in both MA and NH, resides in VT, MA, or NH | VT_NOMTCH_IN3 | 6,014 | 723 | 725 | -2 |
| Licensed in VT, foreign residential address | VT_FOREIGN | 2,059 | 25 | 25 | 0 |
| Licensed in VT, NY residential address | VT_NY | 749 | 40 | 40 | 0 |
| Remainder in VT | VT_REMAINDER | 4,994 | 55 | 55 | 0 |
| Licensed in NH, matched with a record from MA but not one from VT | NH_MTCHMA | 4,161 | 169 | 170 | -1 |
| Licensed in NH, did not match with a MA or VT record | NH_NOMTCH | 12,866 | 775 | 775 | 0 |
| Licensed in MA, did not match with a NH or VT record | MA_NOMTCH | 105,797 | 1,168 | 1,170 | -2 |
| - | Subtotal, MA VT NH | 138,462 | 3,079 | 3,075 | -5 |
| Cumulative State totals or estimated State | e totals across strata | | · | · · · · | |
| Virginia | VA | 80,553* | 1,787 | 1,787.5 | -0.5 |
| Maryland | MD | 63,645* | 1,523 | 1,523.9 | -0.9 |
| District of Columbia | DC | 20,289* | 1,330 | 1,330.7 | -0.7 |

Table B1-3. Sampling States where strata are based on RNs licensed in neighboring States as well (continued)

| Description | IS | Final 2008 frame size after unduplication | 2008 sample sizes after assigning AK, HI, NM samples from different States or strata to AK, HI, and NM | 2008 allocation on which sampling rates were based | Difference between final and targeted sample sizes (differences due to sampling RNs living in AK, HI, and NM separately) |
|--|--------------------|--|---|---|---|
| Strata based on address in | | | | | |
| Alaska | AK | 49 | 5 | х | Х |
| Hawaii | HI | 87 | 5 | х | Х |
| New Mexico | NM | 26 | 1 | x | Х |
| Strata descriptions | | | | | |
| Licensed in MD, matched a record from DC or VA, living in DC metro area | MD_MTCH_METRO | 5,449 | 447 | 446.9 | 0.1 |
| Licensed in MD, matched a record from DC or VA, living outside DC metro area | MD_MTCH_NOMET | 2,455 | 105 | 104.9 | 0.1 |
| Licensed in MD, did not match a record from DC or VA, living in DC metro area | MD_NOMTCH_METRO | 7,874 | 541 | 541.3 | -0.3 |
| Licensed in MD, did not match a record from DC or VA, living outside DC metro area | MD_NOMTCH_NOMET | 47,813 | 430 | 430.7 | -0.7 |
| Licensed in VA, matched a record from DC but not MD, living in DC metro area | VA_MTCH_METRO | 2,400 | 197 | 196.8 | 0.2 |
| Licensed in VA, matched a record from DC but not MD, living outside DC metro area | VA_MTCH_NOMET | 1,080 | 46 | 46.2 | -0.2 |
| Licensed in VA, did not match a record from DC or MD, living in DC metro area | VA_NOMTCH_METRO | 11,087 | 762 | 762.2 | -0.2 |
| Licensed in VA, did not match a record from DC or MD, living outside DC metro area | VA_NOMTCH_NOMET | 65,094 | 782 | 782.3 | -0.3 |
| Licensed in DC, did not match VA or MD, living in Metro area | DC_NOMTCH_METRO | 5,571 | 383 | 383 | 0 |
| Licensed in DC, did not match VA or MD, living outside Metro area | DC_NOMTCH_NOMET | 3,802 | 152 | 152.9 | -0.9 |
| - | Subtotal, VA MD DC | 152,787 | 3,856 | 3,847.3 | -2.3 |

Table B1-3. Sampling States where strata are based on RNs licensed in neighboring States as well (continued)

| Descript | tions | Final 2008 frame size after unduplication | 2008 sample sizes after assigning AK, HI, NM samples from different States or strata to AK, HI, and NM | 2008 allocation on which sampling rates were based | Difference between final and targeted sample sizes (differences due to sampling RNs living in AK, HI, and NM separately) |
|---|---------------------------|--|---|---|---|
| Cumulative State totals or estimated St | tate totals across strata | | | | |
| Delaware | DE | 12,235 | 1,115 | 1,115 | 0 |
| Strata based on address in | | | | | |
| Alaska | AK | 1 | 0 | х | X |
| Hawaii | HI | 0 | 0 | х | X |
| New Mexico | NM | 0 | 0 | х | X |
| Strata descriptions | | | | | |
| Licensed in DE and NJ, but not PA | DE_NJ | 371 | 20 | 20 | 0 |
| Licensed in DE and PA, but not NJ | DE_PA | 2,234 | 160 | 160 | 0 |
| Licensed in DE, NJ, and PA | DE_NJ_PA | 646 | 35 | 35 | 0 |
| Remainder of those licensed in DE | DE_REMAINDER | 8,983 | 900 | 900 | 0 |
| Subtotal, DE | | 12,235 | 1,115 | 1,115 | 0 |
| Subtotal, Multistrata States | | 1,265,808 | 18,487 | 18,387 | -29.3 |
| Total | | 3,568,940 | 55,171 | 54,937.3 | 233.7 |

Table B1-3. Sampling States where strata are based on RNs licensed in neighboring States as well (continued)

* Total prior to unduplication-a small number of RNs licensed in VA but living in MD or vice versa appear in both lists though they should not since they are Compact States- as a result, strata, taking into account residence in the DC metro area, were not formed in a fashion that would permit the sum of the stratum totals to provide unduplicated totals for VA, MD, and DC separately.

The overall NSSRN unweighted response rate was 62.41 percent while the overall weighted response rate was 61.48 percent.² Response rates varied by State (Table B2-1). Unweighted response rates ranged from 49.62 percent (New Mexico) to 74.79 percent (Wisconsin), while weighted response rates ranged from 49.54 percent to 74.87 percent (also for New Mexico and Wisconsin, respectively). State is defined here as the State licensure file(s) on which the given RN could be found; RNs appearing in more than one State's licensure files can contribute to more than one State's (weighted and unweighted) response rate.

B.2.1 Vermont Sample Performance

Vermont's listing of RNs with currently active licenses was partitioned into a number of strata in order to deal with a variety of issues. These include: RNs licensed in Vermont and either Massachusetts or New Hampshire, or both; a large number of RNs licensed in Vermont, but living in foreign countries; and RNs licensed in Vermont but with addresses indicating they are living in States but not in the immediate vicinity of Vermont. Vermont appears to be somewhat of a gateway for RNs from foreign countries seeking to become licensed in the United States.

RNs licensed in Vermont with addresses indicating they are living in the United States but not in the immediate vicinity of Vermont were sampled at a disproportionately low rate for Vermont. It was considered highly unlikely that this group of RNs would either work or live in Vermont and as a result, would not contribute to the Vermont estimates. In fact, no RNs who responded from this latter group did work or live in Vermont, so this sampling strategy appears to have accomplished what it was intended to: reduce the expenditure of survey resources on records not likely to contribute to estimates for groups of analytic interest for Vermont estimates.

Among RNs licensed in Vermont, living in the United States but not in the immediate vicinity, the response rates were low. From among 49 sampled RNs, only 13 completed the 2008 NSSRN, yielding a response rate of roughly 25 percent. The reason for the low response rate for this particular group is uncertain. While this portion of the Vermont sample was small, it had a dramatic impact on the weighted response rate for RNs sampled from among those licensed in Vermont.

² The American Association for Public Opinion Research Response Rate #3 calculation was used; it is defined as the number of completed surveys divided by the number of returned surveys (complete plus partial) plus an estimate of the number of eligible cases among those who did not respond and whose eligibility could not be ascertained.

| | Sample | Responding | Unweighted | Weighted |
|----------------|--------|-------------|---------------|---------------|
| State | size | sample size | response rate | response rate |
| Total | 54,805 | 34,219 | 62.44 | 61.47 |
| Alaska | 777 | 479 | 61.65 | 61.43 |
| Alabama | 843 | 516 | 61.21 | 61.31 |
| Arkansas | 894 | 561 | 62.75 | 62.78 |
| Arizona | 952 | 505 | 53.05 | 53.03 |
| California | 3,283 | 1,744 | 53.12 | 52.91 |
| Colorado | 899 | 606 | 67.41 | 67.52 |
| Connecticut | 992 | 663 | 66.83 | 66.77 |
| DC | 1,331 | 761 | 57.18 | 57.20 |
| Delaware | 1,114 | 716 | 64.27 | 64.19 |
| Florida | 1,902 | 1,060 | 55.73 | 55.58 |
| Georgia | 1,096 | 596 | 54.38 | 54.49 |
| Hawaii | 845 | 483 | 57.16 | 57.03 |
| lowa | 799 | 550 | 68.84 | 68.92 |
| Idaho | 796 | 558 | 70.10 | 70.12 |
| Illinois | 1,603 | 921 | 57.45 | 57.56 |
| Indiana | 845 | 568 | 67.22 | 67.21 |
| Kansas | 893 | 598 | 66.97 | 66.99 |
| Kentucky | 750 | 474 | 63.20 | 63.23 |
| Louisiana | 796 | 482 | 60.55 | 60.45 |
| Massachusetts | 1,384 | 931 | 67.27 | 66.87 |
| Maryland | 1,525 | 971 | 63.67 | 65.88 |
| Maine | 898 | 589 | 65.59 | 65.63 |
| Michigan | 1,204 | 819 | 68.02 | 68.21 |
| Minnesota | 848 | 613 | 72.29 | 72.21 |
| Missouri | 898 | 616 | 68.60 | 68.46 |
| Mississippi | 950 | 533 | 56.11 | 56.21 |
| Montana | 784 | 528 | 67.35 | 67.39 |
| North Carolina | 1,100 | 721 | 65.55 | 65.64 |
| North Dakota | 747 | 526 | 70.41 | 70.41 |
| Nebraska | 696 | 496 | 71.26 | 71.24 |
| New Hampshire | 1,031 | 705 | 68.38 | 68.57 |
| New Jersey | 1,423 | 862 | 60.58 | 60.55 |
| New Mexico | 798 | 396 | 49.62 | 49.54 |
| Nevada | 1,067 | 580 | 54.36 | 54.34 |
| New York | 2,647 | 1,427 | 53.91 | 53.79 |
| Ohio | 1,447 | 983 | 67.93 | 67.98 |
| Oklahoma | 941 | 541 | 57.49 | 57.37 |
| Oregon | 849 | 595 | 70.08 | 69.97 |
| Pennsylvania | 1,897 | 1,274 | 67.16 | 67.25 |
| Rhode Island | 980 | 614 | 62.65 | 62.68 |
| South Carolina | 848 | 500 | 58.96 | 59.02 |
| South Dakota | 744 | 533 | 71.64 | 71.61 |
| Tennessee | 801 | 477 | 59.55 | 59.47 |
| Texas | 1,815 | 1,012 | 55.76 | 55.93 |
| Utah | 799 | 548 | 68.59 | 68.47 |
| Virginia | 1,792 | 1,144 | 63.84 | 62.86 |

 Table B2-1.
 Response rates, unweighted and weighted by State and for the United States as a whole

| State | Sample size | Responding sample size | Unweighted response rate | Weighted response rate |
|---|----------------|------------------------|-----------------------------|---------------------------|
| Vermont (all) | 937 | 610 | 65.10 | 52.04 |
| Vermont (excluding the "outlier" stratum) | 888 | 597 | 67.23 | 63.34 |
| Washington | 922 | 642 | 69.63 | 69.67 |
| Wisconsin | 853 | 638 | 74.79 | 74.87 |
| West Virginia | 936 | 567 | 60.58 | 60.56 |
| Wyoming | 785 | 559 | 71.21 | 71.28 |

 Table B2-1.
 Response rates, unweighted and weighted by State and for the United States as a whole (continued)

To provide a clear picture of the Vermont sample performance, two sets of response rates are provided. The response rates for Vermont including the RNs sampled from the stratum of Vermont licensees living in States not in the immediate vicinity were 65.10 percent unweighted but 52.04 percent weighted. The response rates presented for Vermont were 67.23 percent unweighted and 63.34 percent weighted when excluding the RNs sampled from this stratum.

B.3 Imputation

Statistical imputation is the process by which question-level missing values are assigned likely or plausible values based on other respondent-provided data elements. Imputation is widely used, and a lot of research has been done on the topic. However, one should not consider imputation as the panacea for nonresponse phenomenon in surveys, but rather as the last resort after trying to get respondent answers. The goal of imputation is to help data users avoid complications that could arise with other options addressing the problem of item missing values.

There are two types of nonresponse in surveys: unit nonresponse and item nonresponse. Unit nonresponse occurs when a sampled RN does not complete all of the questions in the survey. Unit nonresponses were reflected in the calculation of the response rate and in weighting through nonresponse adjustment.

Item nonresponse refers to nonresponse to individual questions, which result in a missing value for an individual question. Item missing values in the 2008 NSSRN were filled in, if available, from the State sample frame files. The variables available in the frames included demographic characteristics (date of birth, race/ethnicity, and gender), resident address, and first RN license date. Not all States provided these variables, and data could be very sparse even when available (e.g., race/ethnicity). If any critical item nonresponse existed after review of the State sample files, respondents were called and queried about the missing data item(s). The process of calling respondents to fill in item nonresponse is referred to as data retrieval. For item nonresponse that persisted after review of the sample frame file, data cleaning and data retrieval, as a last resort, statistical imputation was used to assign values.

To improve response rates, a short form of the questionnaire was adapted for the telephone interview. The short form included all the questionnaire items deemed necessary to obtain a completed survey. Questions not asked in the short form were designated as missing and imputed. However, it should be noted that these contribute to a small percentage of missing values and do not compromise the integrity of the data.

B.3.1 Imputation Methods

In using an imputation procedure, one tries to obtain the best value for the missing case instead of picking an arbitrary value to fill in the missing. Regression modeling is very useful in this process. If a strong regression model is available for a particular variable with missing values, good imputed values can be generated using the model. For example, a registered nurse's year of graduation from initial nursing education and year of first licensure are highly correlated. If the year of licensure is missing, but not the year of graduation, the missing year of licensure value can be imputed using the predicted value of the year of licensure. This is done by a regression model with the year of licensure as the dependent variable and the year of graduation as the independent (or predictor) variable—the possible value for the year of licensure is predicted using the regression model and the predictor variable, year of graduation. The regression model can be estimated from respondent cases with nonmissing year of licensure and year of graduation. The regression model can be improved by using more predictor variables.

To avoid nonsensical values, donor imputation is used. Regression modeling is used to facilitate the donor imputation procedure. Donor imputation is a popular technique whereby a donor cell is selected and the donor's value is donated or copied to the missing (recipient) case on a cell by cell basis. If donor and recipients are selected randomly, the procedure is known as hot-deck imputation. Hot-deck imputation was used extensively for the 2008 NSSRN data.

B.3.2 Procedures

Sometimes, missing values can be precisely deduced. For example, if variables A and B are percentages that are supposed to be summed to 100. If A is present and B is missing, then B should be (100 - A). Such imputation is called deterministic imputation. This step was included in data cleaning and not discussed here.

The majority of missing values however, cannot be deduced with certainty. These missing values are prime candidates for statistical imputation.

Imputation was not attempted for some variables because imputation was unnecessary or impractical (such as the "Other Specify" variables). The "other-specify" variables were not imputed, not because their imputation was too difficult, but because nonmissing answers were unique in the sense that the respondents could not find appropriate answers among the common categories provided in the questionnaire and their answers were individualistic so that using them as donors for missing cases does not make sense. Further, some variables were simply too difficult to impute coherently while maintaining complex inter-relationships with other variables, such as employment setting, job title, upgrade education, and preparation for advanced practice RNs.

Variables that were not statistically imputed for 2008 NSSRN include the following:

- Survey eligibility and nursing employment status (questions 1, 19, and 21). These items were considered essential for survey completion and therefore did not have missing values;
- Address fields except State in questions 55, 59, and 61;
- Licensure and certification detail (questions 70A, 70B, 70C, and 70D);
- All "other-specify" variables and their leading questions;
- Education upgrades (questions13E_A through 13E_J);
- All subitems in the question series about preparation as a nurse practitioner, clinical nurse specialist, nurse midwife and nurse anesthetist (questions 14, 14A, 14B, 14C, 14D, 14E, 14E2, 14F, 14F2, and 14G);
- Employment setting (question 23);
- Job title (question 24);

- Level of care (question 29A);
- Patient population served (question 29B); and
- Clinical specialty; (question 29C).

Lastly, while questions 30, 37, and 49 (the three salary variables) were imputed, some cases still have missing values for these three variables. Imputation could not provide a logical value for these cases. So, even though they were imputed, there are still some missing values for these three variables.

The imputation procedures can be summarized as follows:

- 1. Impute data for items using a preliminary hot deck imputation that respects skip patterns;
- 2. Build a regression model for each variable to be imputed through stepwise regression modeling starting with all available predictors;
- 3. Re-impute data using predictive means matching, through grouping similar regression predicted values, again respecting skip patterns; and
- 4. Cycle through steps 2 and 3 until convergence criteria are met for all the variables or the maximum number of iterations specified is reached.

B.3.3 Results and Implications on Variance Estimates

For each imputed variable a conditional imputation rate was computed. Conditional rate is defined as the ratio of the number of imputed applicable cases by the total number of applicable cases in the data set. The conditional rate allows the analyst to see impact of imputation on data analysis.

Eighty percent of the imputed variables for the 2008 NSSRN have imputation rates below 10 percent (Table B3-1), 19 percent are borderline (imputation rate of 10-19 percent), and three variables (1 percent) have an imputation rate greater than 20 percent. The three variables are: question 11B—total number of hours provided for emergency preparedness training in the past year (imputation rate: 46 percent); question 13D_A—year receiving an upgrade Associate Degree in Nursing (ADN) (imputation rate: 36 percent); and question 41—years until planned return to paid work in nursing (imputation rate: 21 percent). When these variables are analyzed, the analyst should be aware of the potential underestimation of the variance and bias in the point estimates.

| Imputation Rate | Number of Variables | Percent of Variables |
|-----------------|---------------------|----------------------|
| Less than 5 | 126 | 37.6 |
| 5 to 9.9 | 135 | 42.0 |
| 10 to 14.9 | 57 | 17.8 |
| 15 to 19.9 | 6 | 1.6 |
| 20 to 24.9 | 1 | 0.3 |
| 25 to 29.9 | 0 | 0.0 |
| 30 to 34.9 | 0 | 0.0 |
| 35 to 39.9 | 1 | 0.3 |
| 40 to 44.9 | 0 | 0.0 |
| 45 to 49.9 | 1 | 0.3 |

Table B3-1. Distribution of imputation rates

There is a component of variation associated with the generation of imputed values that is generally not reflected in survey estimates of variance. As a result, use of imputation can result in an understatement of the variance of survey estimates. The degree to which estimates of variability have been understated is generally considered negligible if the imputation rate is low (less than 10 percent). However, if the imputation rate is over 20 percent, the degree of underestimation may be nontrivial. When these variables are analyzed, the analyst should be aware of the potential underestimation of the variance and bias in the point estimates.

B.4 Weighting and Estimation

The 2008 NSSRN used a stratified, single-stage list frame sample of registered nurses in the United States. Sample weights have been established to permit the development of national or State-level estimates for RNs licensed, working, or residing in the United States. The final weight assigned to a participating registered nurse may be interpreted as the number of RNs with a currently active license in at least one State that the RN represents.

This section describes the methods used for computing the sampling weights and for calculating estimates and their variances. The final sample weights incorporate a number of different components:

- Assignment of initial weights to responding and ineligible RNs in each sample stratum;
- Adjustments for survey nonresponse;

- Adjustments to reflect multiple chances for selection; and
- Adjustments to achieve agreement with State totals of number of RNs licensed.

Survey estimates can be obtained by applying the final sample weights to records representing the eligible responding RNs. Corresponding estimates of variability (standard errors or confidence intervals) can be obtained through the use of 100 jackknife replicate weights developed for this purpose.

B.4.1 Sample Weighting

The 2008 NSSRN weighting process consisted of several steps. The first few steps involved the use of records at the "sampled license" level. The sampled license level reflects the original unit of sample selection that was a record on the list of RNs with currently active licenses. Thus, for these initial steps, any RN sampled multiple times (due to the holding of licenses in multiple States) will have several records associated with her/him. After completion of the adjustments at the sampled license level, any RN sampled multiple times had her/his records "folded" into a single record, producing a database with a single record representing each RN who participated in the 2008 NSSRN.

The weighting process can be described as follows:

- Determination of the initially assigned weights at the sampled license level;
- Adjustment for survey nonresponse by age group (for strata where age was available);
- Assignment of factors to account for multiple opportunities for selection (determined as part of a matching effort to determine the number of sample frames on which an RN appeared);
- Establishing an initial weight at the RN level; and
- Iterative Raking procedures to converge on State and national totals.

Each step is described in more detail below.

B.4.1.1 Initially Assigned Weights

In past implementations of the NSSRN, a single weight was assigned to each RN. This weight was designed to incorporate both the probability of selection and attain a global adjustment for nonresponse for each State separately. No information was used to try to address issues of differential nonresponse by subgroups (e.g., by age). In part, this may have been due to the cumbersome nature of the weighting process employed. Moreover, there were relatively few variables available on both respondents and nonrespondents from which adjustment classes could be formed. Nevertheless, there have been concerns that younger RNs were underrepresented in the survey effort.

As a result of concerns about the representativeness of the various age groups, plans were made to incorporate age into the process for nonresponse adjustment of the weights for the 2008 NSSRN where possible. It was operationally most efficient to initially assign a "global nonresponse adjustment weight" to those sampled RN records with a response disposition of "eligible respondent" or "ineligible" (similar to what has been done for previous implementations of the NSSRN). Then, for those States where age was available for nonresponse adjustment purposes, an additional adjustment was made to account for differential propensities to respond by age group. This approach is mathematically equivalent to assigning weights reflecting the reciprocal of the probability of selection to all sampled records, including nonrespondents, within a stratum and then undertaking the same adjustments for nonresponse, but it requires far less effort.

The nonresponse adjustment process (for both the global and the age-group approaches) was actually a poststratification with population totals obtained from the corresponding sample frame for a given stratum. The initially assigned weight for all eligible respondents and ineligibles in stratum h was calculated as follows:

$$w_{hi} = \frac{N_h}{n_{h,resp \ or \ inelig}}$$

Where

 w_{hi} = the initially assigned weight for an RN sampled record *i* from stratum *h* that had been assigned a response disposition indicating an eligible respondent or ineligible

 N_h = the frame count in stratum *b*

 $n_{h,resp or inelig}$ = the number of eligible responding or ineligible sampled RN records selected from stratum b

Note that the sum of the initially assigned weights for all responding or ineligible RN records within stratum h is thus equal to the frame count for the stratum.

B.4.1.2 Adjustment for Differential Nonresponse by Age Groups

For those States where age of the RN at the time of sample selection could be determined, response rates by age group were examined. Date of birth was not provided on the sample frames for the following States: Georgia, Hawaii, Illinois, Louisiana, New York, Pennsylvania, and Rhode Island. While New York did not provide date of birth in their sample frame, the State subsequently provided an age distribution of RNs appearing on the New York listing. Age was provided by virtually all RNs who responded to the survey and was imputed for the remaining few cases where age was not obtained. Age was imputed for roughly 3 percent of all respondents—those respondents who did not provide their age during data collection, plus some ineligible cases.

Examination of frequency distributions by age showed that propensity to respond to the 2008 NSSRN varied by age. Younger RNs (chiefly, those under 40 though this could vary by State) generally had lower response rates than did older RNs (those aged 50 or older). Thus, nonresponse adjustments were undertaken by age group for those States where age was available. Again, the adjustment process was handled through poststratification: the adjustment factors were computed within sample strata as the ratio of a population (frame) total associated with a given age group to the sum of the weights of the eligible respondents and the ineligibles in that same age group. The resulting poststratified weights were thus computed as:

$$w_{hci} = \left(\frac{N_{hc}}{\sum_{i \in c} w_{hi}}\right) w_{hi}$$

Where

 w_{hci} = the poststratified weight calculated for RN record *i* in poststratification cell *c* of stratum *h*

 N_{hc} = the frame count in for adjustment cell *c* within stratum *b*

 w_{hi} = the initially assigned weight for RN record *i* in stratum *h*

The poststratification cells *c* within each stratum were defined by age.

For those States where age was not available (with one exception discussed below), the initially assigned weight was simply retained.

Note that for the purposes of computing adjustments for differential nonresponse by age, RN records selected from strata associated with residence in Alaska, Hawaii, or New Mexico (regardless of the State frame from which they were sampled) were treated as if they were sampled from a single stratum associated with Alaska, Hawaii, and New Mexico, respectively. (Please see Section B.1 for a discussion of the strata associated with Alaska, Hawaii and New Mexico).

As described in Section B.1 on Sampling, the New York sample frame was partitioned into strata representing overlap between the New Jersey and Pennsylvania frames. New Jersey records included date of birth, so those RNs licensed in both New York and New Jersey already had their weights adjusted for nonresponse. The focus of the nonresponse adjustment here was for those RNs who were licensed in New York but not New Jersey, which represented about 92 percent of all records on the New York frame.

For such RNs who responded or were ineligible and that were sampled from a stratum which included New York licensees but not those from New Jersey, estimates of the number of RNs in two age-based adjustment cells ("under 40" and "41 or older") were developed and served as control totals for the poststratification process so that the weights of eligible responding and ineligible RNs were calibrated to match the cell estimates. Since there was a bit more uncertainty associated with the New York poststratification (estimates of the control totals had to be developed and a few records had imputed ages), it was decided to use only two cells, thus simplifying the poststratification process and limiting the opportunities for misclassification which would be a potential source of bias. It was expected that the gains in reducing the potential for bias due to differential nonresponse would be greater than the potential for bias arising from any misclassification that might be incurred.

Upon completion of this poststratification process, the sum of the resulting weights for all responding or ineligible cases within a stratum equaled the frame counts for each stratum, and, if summed within age groups for the States where age adjustments were possible, corresponded to frame counts by age group as well.

B.4.1.3 Adjustment to Account for Multiple Chances of Selection

An extensive effort was undertaken to identify the number of sample frames on which a selected RN appeared, and thus, the number of chances that an RN had to be sampled. To appropriately account for this probability of selection in the development of the sample weight, a factor H_i was established indicating the number of different strata in which RN *i* appeared, whether or not he or she happened to be sampled from a given frame. This factor can be used to implement the "generalized weight share method" described by Lavallee (2007). Lohr (2010) provides an illustrative example of this method in the form of a textbook problem. The weight adjustment is simply the reciprocal of H_i . It is multiplied by the nonresponse adjusted weight associated with each eligible responding or ineligible RN sampled record (license) *i* to produce the "penultimate" weight $w_{p,i}$ (the "final weight at the sampled RN record" level—responding RNs sampled from multiple strata will have a different "final" weight associated with each sampled record, with each weight depending on the stratum from which the record was sampled):

$$W_{p,i} = \left(\frac{1}{H_i}\right) W_{hci}$$

B.4.1.4 Establishing a Sample Weight for Each Registered Nurse

The final sample weight is at the RN level. That is, each eligible responding RN receives a single weight, regardless of the number of times he or she was sampled. Up to this point the weighting process was at the level of (RN) sampled record(s) (licenses). To obtain a single weight w_j for each RN j, the weights $w_{p,i}$ for all sampled license records i associated with the same RN j were summed. This serves as the initial RN level weight w_j (prior to raking) and was calculated as follows:

$$w_j = \sum_{i \in j} w_{p,i}$$

The sum over all of the RN level weights w_j across all States provides an estimate of the number of registered nurses in the nation who appear on at least one sample frame used in sampling. Thus, it represents an estimate of all RNs with currently active licenses in the United States, plus a relatively small number of RNs who previously had active licenses and have had them lapse, but who have not been purged from the State files of licensed RNs. This national estimate of the number of unique

RNs (virtually all with currently active licenses) is represented as the quantity Y in the discussion of Raking in the next section.

B.4.1.5 Raking to State and National Totals

In the development of weights to represent individual RNs rather than sampled licenses, some departures from the numbers licensed in a State were introduced. To restore consistency with State totals while maintaining consistency with the national estimate *Y* described above, the weights were raked to a set of marginal distributions. Raking is an iterative process of adjusting weights that, when convergence is obtained, achieves consistency with a set of dimensions, with each dimension representing a population distribution across one or more variables (see, for example, Brick and Kalton (1996)). It is a standard way of calibrating sample weights, generally serving the same purpose as poststratification but without needing to account for the full cross-classification of all levels of all dimensions. It can be useful when there are many different sets of control totals with which one would wish to achieve consistency, none of which represent extremely detailed interactions of different variables.

For the 2008 NSSRN, 51 sets of raking control totals were established associated with the 50 States and the District of Columbia. For each set, two totals were used as follows:

| Dimension | Total |
|--------------------------|-----------|
| State k (e.g., Ohio) | X_k |
| Balance of United States | $Y - X_k$ |

Here X_k represents the number of unique RNs appearing on the sample frame of a given State (most of whom are currently active, a few of whom may not be but have not yet been purged from the State file of licensed RNs in State k). Y is the estimate of the number of unique RNs appearing on at least one State's listing of RNs with currently active licenses.

The raking process can be described as follows. The State(s) where each RN was licensed was determined, based on an extensive matching effort between the sample and frames. The weights of all RNs determined to be licensed in State 1 (whether sampled from frames associated with State 1 or not) were adjusted so that they summed to the number on the frame for State 1 while the remainder of the weights (those associated with States 2-51) were adjusted to sum to the estimate of the remainder of the unique RNs appearing on frames 2-51. Then, the weights of all RNs

determined to be licensed in State 2 (whether sampled from frames associated with State 2 or not) were adjusted so that they summed to the number on the frame for State 2 while the remainder of the weights (those associated with States 1 or 3-51) were adjusted to sum to the estimate of the remainder of the unique RNs appearing on frames 1 plus 3-51. This process was continued until the targeted consistency with the 51 State totals and the national total was achieved (the weights converged to the targeted totals).

The final weight for RN j is indicated as $w_{f,j}$. After the completion of the raking process, all RNs characterized as ineligible for the survey were removed from further consideration. The sum of the weights $w_{f,j}$ across all eligible respondents to the 2008 NSSRN thus provides an estimate of the total number of RNs with currently active licenses at the time of the NSSRN. Note that there is no State subscript because an RN may be licensed in multiple States. Estimates of the number of RNs employed in a given State, or currently resident in a given State, can be developed based on the data collected from survey respondents (such information was not utilized as part of the weighting process since all sampling frames and corresponding control totals reflect licensure status only).

B.4.2 Estimation, Variance Estimation, and Sampling and Nonsampling Errors

B.4.2.1 Estimation

Final 2008 NSSRN estimates are computed using the final, full-sample weight $W_{f,j}$ for a given RN j along with the variable of interest. For example, an estimate of the total number of RNs working in a particular State is based on the following indicator variable x where:

 $x_i = 1$ if RN *j* worked in State *k*,

= 0 otherwise

The estimated total \hat{X} is computed as:

$$\hat{X} = \sum_{j \in \text{ state } k} W_{f,j} x_j$$

Where

 $W_{f,i}$ = the final full-sample weight for a given RN j

Estimates of ratios and averages are obtained as the ratio of estimated totals, where the estimated totals are defined as given above (for averages the estimated total in the denominator will be determined as a sum of weights).

B.4.2.2 Variance Estimation

The 2008 NSSRN, as with previous administrations of the NSSRN, is considered a complex sample survey. To avoid concerns of bias, estimates of variance must be developed that fully reflect the sample design and sample weighting. One such approach is Jackknife replication (Wolter, 2007), which has been used for the 2008 NSSRN as well as for previous NSSRNs.

Variance estimation using the jackknife approach requires the weights for the full sample and a set of replicate weights using standard procedures for identifying and computing replicate weights. Each replicate weight is based on a subsample of the full NSSRN sample. The variability of an estimate is measured as the variation of the estimates obtained from the replicate subsamples about the estimate based on the full sample weight. For the 2008 NSSRN, 100 replicate weights were constructed. This is a substantial increase over the 20 replicate weights obtained under the previous NSSRN sample design (the alpha segment approach) and thus provides more degrees of freedom for variance estimation.

The variance of estimate y is estimated as follows using the jackknife replication method of variance estimation developed for the 2008 NSSRN:

$$Var(y) = \sum_{m=1}^{100} (y_m - y)^2$$

Where

y = an estimated total obtained using the full sample weight

and

 y_m = an estimated total obtained using replicate weight m

There are many widely available software packages such as SUDAAN, WesVar, and Stata that permit the development of variance estimates using replicate weights.

B.4.2.3 Sampling and Nonsampling Errors

Given reasonably large sample sizes, relatively precise estimates of the licensed RN population of the United States can be made because of the underlying probability sample. However, several sources of error can cause sample estimates to differ from the true population values. These sources of error are commonly classified as sampling errors and nonsampling errors. Each of these sources of error is briefly described.

Sampling Error

Sample survey estimates are subject to variation. This error is known as sampling error and is the consequence of not measuring every unit in the population. Sampling error is measurable, however, and can be reduced with larger sample sizes and more efficient sample designs. Sample design elements that increase variances of estimates are variation in sampling rates and clustering of the sample. In the 2008 NSSRN, in order to increase precision for estimation purposes at the State level, variation in sampling rates arose between States. Sampling rates for smaller States were generally larger than sampling rates for larger States. Rates also varied between strata within a State for the small percentage who were:

- In a State frame and also resident of Alaska, New Mexico, or Hawaii; and
- Members of the groups of States where multi-State strata were developed.

The sampling error of an estimate is measured as its variance or standard error (the square root of the variance.) Given the 100 replicate weights and the roughly 100 degrees of freedom thus available, the probability is approximately 95 percent that the statistic of interest differs from the true population value by not more than 1.96 standard errors.

Specifically, a 95 percent confidence interval for an estimate x can be constructed as:

 $x \pm 1.96\sigma_x$

where

 σ_x = the estimated standard error of x.

Standard errors for a select list of estimates is presented in Table B4-1.

| RN S.E. | | | |
|---|------------|---------------|----------------------|
| | population | Estimated | S.E. of estimated |
| Description | | | |
| Description | estimate | percentage | percentage |
| Initial nursing education | COO E 40 | 00.00 | 0.00 |
| Diploma program | 623,548 | 20.36 | 0.23 |
| Associate degree | 1,388,884 | 45.34 | 0.33 |
| Bachelor's degree | 1,030,026 | 33.63 | 0.33 |
| Master's degree | 13,325 | 0.44 | 0.05 |
| Doctorate | 954 | 0.03 | 0.01 |
| Not reported | 6,425 | 0.21 | 0.03 |
| Employed in nursing | | | |
| Yes | 2,596,599 | 84.77 | 0.23 |
| No | 466,564 | 15.23 | 0.23 |
| Racial/ethnic background | | | |
| White (non-Hispanic) | 2,549,484 | 83.22 | 0.25 |
| Black/African American (non-Hispanic) | 165,433 | 5.40 | 0.16 |
| Asian (non-Hispanic) | 169,454 | 5.53 | 0.16 |
| American Indian/Alaskan Native (non-Hispanic) | 8,571 | 0.28 | 0.03 |
| Native Hawaiian/Pacific Islander (non-Hispanic) | 9,528 | 0.31 | 0.04 |
| Two or more races (non- Hispanic) | 51,568 | 1.68 | 0.09 |
| Hispanic/Latino (White) | 96,122 | 3.14 | 0.13 |
| Hispanic/Latino (Black/African American) | 2,727 | 0.09 | 0.02 |
| Hispanic/Latino (two or more races) | 4,116 | 0.14 | 0.03 |
| Hispanic, other | 6,109 | 0.20 | 0.03 |
| Graduation year | , | | |
| Before 1961 | 85,468 | 2.79 | 0.11 |
| 1961-1965 | 106,810 | 3.49 | 0.11 |
| 1966-1970 | 161,794 | 5.28 | 0.14 |
| 1971-1975 | 258,004 | 8.42 | 0.18 |
| 1976-1980 | 349,421 | 11.41 | 0.21 |
| 1981-1985 | 362,664 | 11.84 | 0.20 |
| 1986-1990 | 318,374 | 10.39 | 0.20 |
| 1991-1995 | 420,254 | 13.72 | 0.22 |
| 1996-2000 | 396,991 | 12.96 | 0.23 |
| 2000-2004 | 289,760 | 9.46 | 0.20 |
| After 2004 | 313,622 | 10.24 | 0.20 |
| Employment setting, if employed in nursing | 515,022 | 10.24 | 0.20 |
| | 1,601,831 | 52.29 | 0.31 |
| Hospital Nursing home extended | 135,514 | 52.29 4.42 | 0.31 |
| - | , | 4.42 3.21 | 0.14 0.11 |
| Nursing education | 98,268 | | |
| Public health/community health | 262,908 | 8.58 | 0.19 |
| School health service | 84,418 | 2.76 | 0.10 |
| Occupation health | 18,840 | 0.62 | 0.05 |
| Ambulatory care | 270,556 | 8.83 | 0.18 |
| Insurance claims/benefits | 49,441 | 1.61 | 0.08 |
| Other | 51,947 | 1.70 | 0.08 |
| Not reported | 22,875 | 0.75 | 0.06 |
| Not employed in nursing | 466,564 | 15.23 | 0.23 |

| | RN | S.E. of | | |
|-----------------------------|------------|------------|------------|--|
| | population | Estimated | estimated | |
| Description | estimate | percentage | percentage | |
| Highest nursing education | | | | |
| Diploma in nursing | 426,279 | 13.92 | 0.22 | |
| Associate degree | 1,103,471 | 36.02 | 0.33 | |
| Bachelor's in nursing | 1,068,250 | 34.87 | 0.31 | |
| Bachelor's in related field | 56,915 | 1.86 | 0.09 | |
| Master's in nursing | 290,084 | 9.47 | 0.19 | |
| Master's in related field | 85,709 | 2.80 | 0.11 | |
| Doctorate in nursing | 13,140 | 0.43 | 0.04 | |
| Doctorate in related field | 15,229 | 0.50 | 0.04 | |
| Not known | 4,084 | 0.13 | 0.03 | |
| Age of nurse | | | | |
| <25 | 79,641 | 2.60 | 0.11 | |
| 25 to 29 | 208,543 | 6.81 | 0.14 | |
| 30 to 34 | 281,999 | 9.21 | 0.13 | |
| 35 to 39 | 331,972 | 10.84 | 0.14 | |
| 40 to 44 | 350,101 | 11.43 | 0.22 | |
| 45 to 49 | 441,831 | 14.42 | 0.20 | |
| 50 to 54 | 495,985 | 16.19 | 0.24 | |
| 55 to 59 | 396,967 | 12.96 | 0.21 | |
| 60 to 64 | 249,533 | 8.15 | 0.16 | |
| 65 to 69 | 140,263 | 4.58 | 0.12 | |
| 70 to 74 | 53,670 | 1.75 | 0.08 | |
| 75+ | 32,657 | 1.07 | 0.06 | |

Table B4-1. Standard errors for select estimates (continued)*

*Estimated percentages may not add to 100 because of rounding.

Nonsampling Error

Other sources of potential error can arise as part of the survey process. These are not related to the sample design or sample size but rather the implementation of the survey. Thus, nonsampling errors can occur even when a complete enumeration, or census, is undertaken. Potential sources of such error include:

- Unit nonresponse (a sampled RN declines to participate or cannot be found);
- Item nonresponse (a participating RN forgets to or chooses not to answer a question);
- Coding errors; and
- Undercoverage (failure to include some RNs on the sample frame).

Attempts to address these nonsampling errors include:

- Efforts to maximize response rates, including refusal conversion;
- Adjustment of the weights for nonresponse;
- Imputation of omitted data items;
- Careful coding of responses; and
- Highly controlled processing of the survey data.

B.5 Nonresponse Bias Analysis

In any survey that includes sampling or achieves less than a 100 percent response rate there is potential for nonresponse bias. This is to say, respondents to the survey may differ in some ways from nonrespondents, and, to the extent that such differences affect estimates, systematic departures may arise between these estimates and the actual population value (unknown) being estimated. In this section we describe the potential sources of nonresponse bias in the 2008 NSSRN, explain the measures that were taken to reduce that bias, and show two sets of comparisons to help assess the degree to which nonresponse bias may exist in the 2008 NSSRN data.

The sample design and weighting processes for the 2008 NSSRN differed in important ways from those employed in previous implementations of the NSSRN dating back to 1977. Departures from historically utilized methodologies reflected, in part, a concern about undercoverage of racial/ethnic minorities and undercoverage of RNs within particular age groups.

B.5.1 Underrepresentation of Racial and Ethnic Minorities

Most notably from the sample design perspective, independent systematic random samples were selected from each stratum for the 2008 NSSRN, while a nested cluster design (clustering RNs based on last names) had been employed previously. The 2008 approach was expected to reduce sample variances to some extent, particularly for estimates of race/ethnicity since clusters based on last name are highly correlated with racial/ethnic background. There had been concern about possible underrepresentation of some minorities using the original design, and efforts were previously made

to supplement the sample in some States to see if minority representation could be improved. This initiative of the NSSRN sample design achieved little success.

Differential nonresponse for some minorities was another possible reason for lower than expected estimated percentages for minority RNs. Race and ethnicity are not generally available on the NSSRN sample frames that are received from the State boards of nursing, and when available race and ethnicity data have substantial amounts of missing data. In fact, other than date of birth (available on 44 of the State listings for the 2008 NSSRN), the only other information uniformly available on the sample frames that was feasible for nonresponse adjustment purposes was address based. In terms of information actually on the frame this would be ZIP Code.

B.5.2 Underrepresentation of Younger Nurses

In addition to underrepresentation of nurses of minority race and ethnicity in surveys prior to 2008, concern had been expressed that the youngest RNs seemed to be underrepresented in the population. In the 2004 NSSRN, the weights of RNs under the age of 26 were poststratified to control figures that were derived from external data reflecting numbers of people passing nursing licensure exams in the recent past. This age group, however, represents a relatively small percentage of the population of RNs with currently active licenses. Moreover, this apparent underrepresentation of younger RNs could also be due to differential nonresponse, so focusing on a single small class of RNs would not necessarily address the issue of concern. The issue of differential response rates by age among nurses over 25 had never been addressed in previous surveys.

Prior to the 2008 NSSRN, nonresponse adjustment of sample weights was accomplished through poststratification of survey respondents' weights to totals representing frame counts of RNs in a hierarchical ordering of States, essentially from smallest to biggest (the frame counts were adjusted to reflect RNs also licensed in a State appearing higher in the hierarchy and thus already accounted for in the weighting process). This weighting procedure thus did not account for differential nonresponse between subgroups within the RN population.

When data collection for the 2008 NSSRN was completed, the ratio of survey respondents to sampled RNs was computed for specified age groupings for each of the 44 States where date of birth was available, as well as for New York, as described earlier. Examination of these ratios showed that older RNs did indeed respond rates at generally higher than the rates for younger RNs. Thus, as part of the weighting process, adjustment cells were constructed to account for these

differential response rates, helping to ensure that the weights of responding RNs reflect the age distribution of the States in which they were licensed.

B.5.3 Comparative Analyses

Two sets of analyses were undertaken to examine the potential for nonresponse bias in the 2008 NSSRN. The first was a comparison of 2008 NSSRN estimates based on two different sets of nonresponse adjusted weights. The two estimates were computed based on weights devised in an identical fashion except that the second set of weights excluded the ratio adjustment for age. The nonage-adjusted estimates thus simulated the types of estimates that had been developed for prior administrations of NSSRN. Comparison of the two sets of estimates is intended to accomplish two purposes. First, to help data users see what the effect of an improved estimates may be due, not to changes in the underlying population, but rather to an expected increase in the accuracy of the survey estimates correlated with age.

The second set of analyses compares the 2008 NSSRN estimates of age and race to corresponding estimates obtained from the American Community Survey (ACS) for the population in nursing occupations. With different survey methodologies, the estimates are not perfectly comparable. Nevertheless, such comparisons may help provide some insight into the degree to which the potential for nonresponse bias may or may not have been reduced for NSSRN estimates.

Potential for Reduction in Bias Due to Nonresponse Adjustments by Age Group: National Estimates

There has been concern in earlier implementations of the NSSRN that younger RNs were underrepresented due to undercoverage on State sampling frames. In the 2000 and 2004 NSSRN administrations, poststratification of the weights of the youngest RNs (those under the age of 26) was undertaken, adjusting the weights to correspond to estimated control totals that were derived based on an external data reflecting numbers of people passing nursing licensure exams in the recent past. While undercoverage of newly licensed RNs may be an issue, another potential source of bias associated with low estimates of younger RNs is differential nonresponse by age. This was examined as part of the 2008 weighting process.

Forty-four States provided date of birth on the sampling frame. Where date of birth was available, six age groups were formed during sample selection for sorting or implicit stratification purposes. Examination of response propensities by age for these groups showed that younger RNs responded at lower rates than older RNs. The weight adjustment processes that had been employed in the past were essentially global in nature. Poststratification was used to achieve consistency between responding RNs associated with a given State of licensure and the corresponding control total, licensed RNs in the State. The control total took into account RNs who were also licensed in what were characterized as higher priority States. Thus, if older RNs had higher response rates in a State as they often did, they would be represented at a disproportionately high rate among the population of RNs licensed in the State. This is a potential source of bias for survey estimates that are correlated with age. In addition to the general age distribution of the RN population, other variables that could be affected include those related to education and employment.

As previously described, the 2008 NSSRN final weights included a poststratification adjustment by age group. Through this process, the sum of the weights of eligible and ineligible respondents was adjusted by age group to corresponding age group totals for each stratum with States where date of birth was available. Poststratification adjustments for New York used the age distribution of the RN population because date of birth was not included on the sample file. This calibration by age served to adjust for both undercoverage and nonresponse. Thus, nonresponse adjustment reflecting differential nonresponse by age was not feasible for only six States.

To assess the extent to which survey estimates may have changed after accounting for differential nonresponse by age group, we developed a separate set of weights using a global approach to nonresponse adjustment. That is, poststratification of all responding eligible and ineligible RNs to a single control total for each stratum within a State. This approach simulated the approach to nonresponse adjustment used in past administrations of NSSRN. This weight is referred to here as the "surrogate" weight.

Table B5-1 provides the estimated percentage population distributions across the variables of interest and the corresponding estimated standard errors for each set of weights. Table B5-1 also presents a ratio designed to serve as a measure of the extent to which there is evidence of practical significance. This ratio is the difference between the two estimates divided by the standard error of the "final weight" estimate. We will refer to it here as the "standardized measure of the difference."

Note that the "final weight" estimate and the "surrogate" weight estimates are highly correlated, and thus one would expect most all the differences to achieve statistical significance, so we are focusing here on differences that appear to be particularly noteworthy. Under these circumstances, we focus here on ratios with an absolute value of 3 or more, taking them to represent evidence of a difference of some consequence.

Consider the variable "initial nursing education" and the estimates based on the final weight and the surrogate weight for those whose initial nursing education was a "diploma," the first line of estimates for Table B5-1. The final weight estimate for those with a diploma is 20.36 percent while the surrogate weight estimate is 21.62 (the ratio of the former to the latter is .94, while the ratio of standard errors is .89). The difference between the final weight estimate and the surrogate weight estimate is -1.26, and the standardize measure of this difference (the ratio of the actual difference to the standard error corresponding to the "Final weight" estimate, a value of 0.23) is -5.43. This standardized difference value, developed for use in assessing evidence of practical significance between the two estimates, is substantially greater than 3. Using the criteria of obtaining an absolute value of 3 or more, a difference this big can reasonably be interpreted as substantial evidence that the difference is of consequence. The expectation is that using the weight incorporating a nonresponse adjustment reflecting differences arise and thus represent evidence of reduced potential include: RNs with a bachelor's degree; RNs employed in nursing; White, non-Hispanic RNs; RNs employed in a hospital setting; and most of the age group and graduation year categories.

It may be of interest to note that the ratio of standard errors varies by analytical variable. For all age groups except those under 25, the ratio was substantially lower than 1, suggesting that using the poststratification cells based on age may have helped reduce the variability of the corresponding estimates. For the education-related variables, generally the ratios were close to 1. For the race and ethnicity categories, almost all of the ratios are greater than 1, suggesting that there may be an interaction between race/ethnicity and age. This could be the result of the larger proportion of older RNs being White, non-Hispanic. The issue of differential nonresponse by race/ethnicity is considered in more detail in the next section.

These findings regarding the effects of nonresponse adjustment by age group should be taken into account when comparing estimates from the 2008 NSSRN to those from previous NSSRNs. Differences between the 2008 NSSRN estimates and those from prior years for estimates correlated with age may be due to an improvement in the weighting process rather than a significant change in the RN population over time.

| Table B5-1. | Evaluating potential for reduction in bias: Comparing final 2008 estimates with 2008 estimates using a weight that does |
|-------------|---|
| | not account for differential response rates by age* |

| | Final Weight | | Surrogate Weight | | Ratios | | Assessing Differences | |
|---|-------------------------|------------------------------------|-------------------------|------------------------------------|--|--|--|--|
| Description | Estimated Percentage | S.E. of Estimated Percentage | Estimated Percentage | S.E. of Estimated Percentage | Ratio of Estimates: Final to Surrogate | Ratio of SEs: Final to Surrogate | Difference between Estimates: Final minus Surrogate | Ratio: (Difference in Estimates) to (SE of "Final Weight" Estimate) |
| Initial nursing education | · · · · | • | | | | • | | |
| Diploma | 20.36 | 0.23 | 21.62 | 0.26 | 0.94 | 0.89 | -1.26 | -5.43 |
| Associate degree | 45.34 | 0.33 | 45.18 | 0.32 | 1.00 | 1.04 | 0.16 | 0.48 |
| Bachelor's degree | 33.63 | 0.33 | 32.54 | 0.32 | 1.03 | 1.01 | 1.09 | 3.32 |
| Master's degree | 0.44 | 0.05 | 0.42 | 0.05 | 1.05 | 1.04 | 0.02 | 0.42 |
| Doctorate | 0.03 | 0.01 | 0.03 | 0.01 | 1.00 | 1.00 | 0.00 | 0.00 |
| Not reported | 0.21 | 0.03 | 0.21 | 0.03 | 1.00 | 0.97 | 0.00 | 0.00 |
| Employed in Nursing | | | | | | | | |
| Yes | 84.77 | 0.23 | 84.06 | 0.25 | 1.01 | 0.94 | 0.71 | 3.05 |
| No | 15.23 | 0.23 | 15.94 | 0.25 | 0.96 | 0.94 | -0.71 | -3.05 |
| Racial/Ethnic background | | | | | | | | |
| White (non-Hispanic) | 83.22 | 0.25 | 83.74 | 0.24 | 0.99 | 1.03 | -0.52 | -2.08 |
| Black/African American (non-Hispanic) | 5.40 | 0.16 | 5.37 | 0.16 | 1.01 | 1.00 | 0.03 | 0.19 |
| Asian (non-Hispanic) | 5.53 | 0.16 | 5.28 | 0.15 | 1.05 | 1.07 | 0.25 | 1.54 |
| American Indian/Alaskan Native (non- Hispanic) | 0.28 | 0.03 | 0.27 | 0.03 | 1.04 | 1.03 | 0.01 | 0.33 |
| Native Hawaiian/Pacific Islander (non- Hispanic) | 0.31 | 0.04 | 0.30 | 0.04 | 1.03 | 1.06 | 0.01 | 0.26 |
| Two or more races (non- Hispanic) | 1.68 | 0.09 | 1.67 | 0.08 | 1.01 | 1.02 | 0.01 | 0.12 |
| Hispanic/Latino (White) | 3.14 | 0.13 | 2.97 | 0.12 | 1.06 | 1.09 | 0.17 | 1.27 |
| Hispanic/Latino (Black/African American) | 0.09 | 0.02 | 0.08 | 0.02 | 1.13 | 1.16 | 0.01 | 0.45 |
| Hispanic/Latino (Two or more races) | 0.14 | 0.03 | 0.13 | 0.03 | 1.08 | 1.08 | 0.01 | 0.37 |
| Hispanic, Other | 0.20 | 0.03 | 0.19 | 0.03 | 1.05 | 1.07 | 0.01 | 0.31 |
| Graduation Year | | | | | - | - | | |
| Before 1961 | 2.79 | 0.11 | 3.08 | 0.12 | 0.91 | 0.89 | -0.29 | -2.64 |
| 1961-1965 | 3.49 | 0.11 | 3.85 | 0.12 | 0.91 | 0.89 | -0.36 | -3.33 |
| 1966-1970 | 5.28 | 0.14 | 5.84 | 0.16 | 0.90 | 0.90 | -0.56 | -3.97 |
| 1971-1975 | 8.42 | 0.18 | 9.32 | 0.21 | 0.90 | 0.89 | -0.90 | -4.95 |
| 1976-1980 | 11.41 | 0.21 | 12.48 | 0.24 | 0.91 | 0.87 | -1.07 | -5.12 |
| 1981-1985 | 11.84 | 0.20 | 12.24 | 0.21 | 0.97 | 0.96 | -0.40 | -2.00 |
| 1986-1990 | 10.39 | 0.20 | 10.52 | 0.21 | 0.99 | 0.96 | -0.13 | -0.64 |
| 1991-1995 | 13.72 | 0.22 | 13.33 | 0.22 | 1.03 | 1.03 | 0.39 | 1.74 |
| 1996-2000 | 12.96 | 0.23 | 12.00 | 0.21 | 1.08 | 1.06 | 0.96 | 4.27 |
| 2000-2004 | 9.46 | 0.20 | 8.41 | 0.19 | 1.12 | 1.07 | 1.05 | 5.17 |
| After 2004 | 10.24 | 0.20 | 8.92 | 0.18 | 1.15 | 1.10 | 1.32 | 6.60 |

| Table B5-1. | Evaluating potential for reduction in bias: Comparing final 2008 estimates with 2008 estimates using a weight that does |
|-------------|---|
| | not account for differential response rates by age (continued)* |

| | Final V | Veight | Surrogat | e Weight | Ratios | | Assessing D | ifferences |
|---|-------------------------|------------------------------------|-------------------------|------------------------------------|--|--|--|--|
| Description | Estimated Percentage | S.E. of Estimated Percentage | Estimated Percentage | S.E. of Estimated Percentage | Ratio of Estimates: Final to Surrogate | Ratio of SEs: Final to Surrogate | Difference between Estimates: Final minus Surrogate | Ratio: (Difference in Estimates) to (SE of "Final Weight" Estimate) |
| Employment Setting, if employed in nursing | | | | | | | | |
| Hospital | 52.29 | 0.31 | 50.90 | 0.32 | 1.03 | 0.98 | 1.39 | 4.44 |
| Nursing Home/Extended Care | 4.42 | 0.14 | 4.52 | 0.13 | 0.98 | 1.02 | -0.10 | -0.73 |
| Nursing Education | 3.21 | 0.11 | 3.32 | 0.12 | 0.97 | 0.96 | -0.11 | -0.99 |
| Public Health/Community Health | 8.58 | 0.19 | 8.77 | 0.20 | 0.98 | 0.97 | -0.19 | -1.00 |
| School Health Service | 2.76 | 0.10 | 2.87 | 0.10 | 0.96 | 0.99 | -0.11 | -1.12 |
| Occupation Health | 0.62 | 0.05 | 0.63 | 0.05 | 0.98 | 1.02 | -0.01 | -0.20 |
| Ambulatory Care | 8.83 | 0.18 | 8.85 | 0.19 | 1.00 | 0.98 | -0.02 | -0.11 |
| Insurance Claims/Benefits | 1.61 | 0.08 | 1.66 | 0.08 | 0.97 | 1.00 | -0.05 | -0.61 |
| Other | 1.70 | 0.08 | 1.76 | 0.08 | 0.97 | 0.99 | -0.06 | -0.74 |
| Not Reported | 0.75 | 0.06 | 0.76 | 0.06 | 0.99 | 1.00 | -0.01 | -0.16 |
| Not employed in nursing | 15.23 | 0.23 | 15.94 | 0.25 | 0.96 | 0.94 | -0.71 | -3.05 |
| Highest Nursing Education | | | | | | | | |
| Diploma in nursing | 13.92 | 0.22 | 14.73 | 0.24 | 0.95 | 0.93 | -0.81 | -3.65 |
| Associate degree in nursing or related field | 36.02 | 0.33 | 35.64 | 0.32 | 1.01 | 1.04 | 0.38 | 1.14 |
| Bachelor's in nursing (or related field for 2008) | 34.87 | 0.31 | 33.87 | 0.30 | 1.03 | 1.01 | 1.00 | 3.25 |
| Bachelor's in related field | 1.86 | 0.09 | 2.00 | 0.09 | 0.93 | 0.93 | -0.14 | -1.63 |
| Master's in nursing (or related field 2008) | 9.47 | 0.19 | 9.66 | 0.20 | 0.98 | 0.98 | -0.19 | -0.98 |
| Master's in related field | 2.80 | 0.11 | 2.98 | 0.12 | 0.94 | 0.97 | -0.18 | -1.59 |
| Doctorate in nursing (or related field 2008) | 0.43 | 0.04 | 0.46 | 0.05 | 0.93 | 0.93 | -0.03 | -0.71 |
| Doctorate in related field | 0.50 | 0.04 | 0.53 | 0.05 | 0.94 | 0.93 | -0.03 | -0.70 |
| Not reported | 0.13 | 0.03 | 0.14 | 0.03 | 0.93 | 0.96 | -0.01 | -0.40 |

| | Final Weight | | Surrogate Weight | | Ratios | | Assessing Differences | |
|--------------|-------------------------|------------------------------------|-------------------------|------------------------------------|--|--|--|--|
| Description | Estimated Percentage | S.E. of Estimated Percentage | Estimated Percentage | S.E. of Estimated Percentage | Ratio of Estimates: Final to Surrogate | Ratio of SEs: Final to Surrogate | Difference between Estimates: Final minus Surrogate | Ratio: (Difference in Estimates) to (SE of "Final Weight" Estimate) |
| Age of nurse | | | | | | | | |
| <25 | 2.60 | 0.11 | 2.15 | 0.09 | 1.21 | 1.19 | 0.45 | 4.21 |
| 25 to 29 | 6.81 | 0.14 | 5.65 | 0.16 | 1.21 | 0.84 | 1.16 | 8.47 |
| 30 to 34 | 9.21 | 0.13 | 7.93 | 0.17 | 1.16 | 0.77 | 1.28 | 10.00 |
| 35 to 39 | 10.84 | 0.14 | 9.50 | 0.21 | 1.14 | 0.66 | 1.34 | 9.57 |
| 40 to 44 | 11.43 | 0.22 | 11.14 | 0.23 | 1.03 | 0.93 | 0.29 | 1.34 |
| 45 to 49 | 14.42 | 0.20 | 14.12 | 0.24 | 1.02 | 0.82 | 0.30 | 1.51 |
| 50 to 54 | 16.19 | 0.24 | 17.97 | 0.30 | 0.90 | 0.79 | -1.78 | -7.54 |
| 55 to 59 | 12.96 | 0.21 | 14.34 | 0.23 | 0.90 | 0.89 | -1.38 | -6.70 |
| 60 to 64 | 8.15 | 0.16 | 9.02 | 0.18 | 0.90 | 0.89 | -0.87 | -5.47 |
| 65+ (65-69) | 4.58 | 0.12 | 5.06 | 0.14 | 0.91 | 0.86 | -0.48 | -4.00 |
| 70 to 74 | 1.75 | 0.08 | 1.94 | 0.09 | 0.90 | 0.93 | -0.19 | -2.29 |
| 75+ | 1.07 | 0.06 | 1.18 | 0.07 | 0.91 | 0.90 | -0.11 | -1.72 |

 Table B5-1.
 Evaluating potential for reduction in bias: Comparing final 2008 estimates with 2008 estimates using a weight that does not account for differential response rates by age (continued)*

* Estimated percentages may not add to 100 because of rounding.

Comparing 2008 NSSRN and 2008 American Community Survey Estimates

The American Community Survey (ACS) is a sizeable survey of the U.S. population carried out by the U.S. Census Bureau and designed primarily to collect data previously obtained from the Census "Long Form." The ACS data however, are collected continually over the course of each decade rather than once every 10 years. A single household respondent is asked to provide data on each member of the household. The ACS is a mail survey that used telephone and in-person follow-up to help reduce nonresponse. The overall response rate reported for the 2008 ACS was 98 percent.

The ACS collects data about occupation, race/ethnicity, and age. It may be of interest to compare the 2008 NSSRN estimates on race/ethnicity and age to corresponding 2008 ACS estimates for the population with an occupation of registered nurse. In making comparisons between the studies, it should be noted that there are some differences in the way the data have been collected that could affect the quality or comparability of the estimates. For instance, the ACS categorizes occupation as the main type of work a person did if the person worked in the previous 5 years. The NSSRN collects data on RNs with currently active licenses regardless of current or past employment status. For the ACS data there is also question about the accuracy of the occupation reporting since a household respondent may be characterizing another household member inaccurately. Points of possible uncertainty on the part of the ACS household respondent include whether: a member is an RN or an licensed practical nurse (LPN) (not a member of the target population for the NSSRN); a member is actually employed as an RN even if he or she has a license; and/or, whether a household member is currently licensed as an RN in at least one State.

Table B5-2 provides estimates from both the 2008 NSSRN and 2008 ACS data and ratios of these estimates. It should be noted that these estimates do not represent precisely the same populations, as the description of the target populations for the various estimates will make clear. Nevertheless, examining these ratios does seem to provide some insight into the potential for nonresponse bias, albeit with necessary caveats.

Table B5-2 shows the 2008 national NSSRN estimates for both those RNs with currently active licenses (3,063,162) and the subset of those employed as an RN (2,596,599). The 2008 ACS estimate for those employed as an RN some time during the previous 5 years is 3,032,470. Thus, this ACS estimate is closer to the figure obtained for all RNs with currently active licenses obtained using 2008 NSSRN data. The 2008 ACS estimate of the number of RNs who were members of the labor force is 2,752,050, closer to the 2008 NSSRN estimate of those employed in nursing.

Table B5-2. Comparing 2008 NSSRN and 2008 American Community Survey (ACS) estimates on racial/ethnic background, and age group

| | | 2008 | NSSRN Estin | nates | | 2008 ACS Estimates | | | |
|---|------------------------------|-------------------------------|---|---|--|--|----------------------------------|--|--|
| | | | Number | F -11 | F-Museta d | Total Estimated Number | | F athanata d | Estimated |
| Racial/ethnic background and age group | Total Estimated Number | Total Estimated Percent | Number in Sample Employed in Nursing | Estimated Population Employed in Nursing | Estimated Percent Employed in Nursing | Employed in Nursing within Last 5 Years | Total Estimated Percentage | Estimated Number in Labor Force | Percentage Among those in Labor Force |
| U.S. Total | 3,063,162 | 100.0 | 28,554 | 2,596,599 | 100.0 | 3,032,470 | 100.0 | 2,752,050 | |
| Racial/ethnic Background | | | - | | | | | | |
| White (non-Hispanic) | 2,549,302 | 83.2 | 24,235 | 2,135,592 | 82.2 | 2,324,060 | 76.6 | 2,093,358 | 76.1 |
| Black/African American (non- Hispanic) | 165,352 | 5.4 | 1,510 | 146,459 | 5.6 | 298,344 | 9.8 | 275,717 | 10.0 |
| Asian (non-Hispanic) | 169,454 | 5.5 | 1,269 | 150,938 | 5.8 | 230,578 | 7.6 | 217,984 | 7.9 |
| Native Hawaiian/Pacific Islander (non-Hispanic) | 9,528 | 0.3 | 84 | 8,792 | 0.3 | 1,611 | 0.1 | | |
| American Indian/Alaskan Native (non-Hispanic) | 8,571 | 0.3 | 122 | 7,838 | 0.3 | 11,434 | 0.4 | | |
| Hispanic/Latino (any Race) | 109,387 | 3.6 | 832 | 101,045 | 3.9 | 133,055 | 4.4 | 123,788 | 4.5 |
| Two or more races (non-Hispanic) Other race (Non-Hispanic) | 51,568 | 1.7 | 502 | 45,935 | 1.8 | 29,160 4,228 | 1.0 0.1 | | |
| Age group | | | | | | | | | |
| Less than 25 | 79,641 | 2.6 | 699 | 77,041 | 3.0 | 105,697 | 3.5 | 101,139 | 3.7 |
| 25-29 | 208,543 | 6.8 | 1,868 | 198,734 | 7.7 | 253,508 | 8.4 | 241,232 | 8.8 |
| 30-34 | 281,999 | 9.2 | 2,467 | 257,472 | 9.9 | 311,285 | 10.3 | 294,137 | 10.7 |
| 35-39 | 331,972 | 10.8 | 2,876 | 302,520 | 11.7 | 367,284 | 12.1 | 345,483 | 12.6 |
| 40-44 | 350,101 | 11.4 | 3,293 | 314,368 | 12.1 | 362,726 | 12.0 | 344,403 | 12.5 |
| 45-49 | 441,831 | 14.4 | 4,333 | 397,552 | 15.3 | 422,303 | 13.9 | 406,798 | 14.8 |
| 50-54 | 495,985 | 16.2 | 5,393 | 433,848 | 16.7 | 458,851 | 15.1 | 434,681 | 15.8 |
| 55-59 | 396,967 | 13.0 | 4,209 | 337,918 | 13.0 | 346,692 | 11.4 | 317,347 | 11.5 |
| 60-64 | 249,533 | 8.1 | 2,234 | 176,932 | 6.8 | 231,271 | 7.6 | 183,088 | 6.7 |
| 65-69 | 140,263 | 4.6 | 809 | 69,595 | 2.7 | 111,668 | 3.7 | 58,367 | 2.1 |
| 70-74 | 53,670 | 1.8 | 251 | 19,752 | 0.8 | 38,178 | 1.3 | 17,824 | 0.6 |
| 75 and over | 32,657 | 1.1 | 122 | 10,867 | 0.4 | 23,007 | 0.8 | 7,551 | 0.3 |

| Table B5-2. | Comparing 2008 NSSRN and 2008 American Community Survey (ACS) estimates on racial/ethnic background, and age |
|-------------|--|
| | group (continued) |

| | | | Ratios: NSS 2008 ACS | | | |
|---|--|--|--|--|---|---|
| Racial/ethnic background and age group | Totals: "NSSRN Full Sample" to "ACS Employed in Nursing in past 5 years" | Percentages: "NSSRN Full Sample" to "ACS Employed in Nursing in past 5 years " | Totals: "NSSRN Employed in Nursing" to "ACS Employed in Nursing in past 5 years " | Percentages: "NSSRN Employed in Nursing" to "ACS Employed in Nursing in past 5 years " | Totals: "NSSRN Employed in Nursing" to "ACS Labor Force " | Percentages: "NSSRN Employed in Nursing" to "ACS Labor Force " |
| U.S. Total | 1.01 | | 0.86 | | 0.94 | |
| Racial/ethnic Background | | | | | | |
| White (non-Hispanic) | 1.05 | 1.04 | 0.88 | 1.03 | 1.02 | 1.08 |
| Black/African American (non-Hispanic) | 0.55 | 0.54 | 0.48 | 0.56 | 0.53 | 0.56 |
| Asian (non-Hispanic) | 0.73 | 0.72 | 0.65 | 0.76 | 0.69 | 0.73 |
| Native Hawaiian/Pacific Islander (non-Hispanic) | | | | | | |
| American Indian/Alaskan Native (non-Hispanic) | | | | | | |
| Hispanic/Latino (any Race) | 0.82 | 0.82 | 0.76 | 0.89 | 0.82 | 0.87 |
| Two or more races (non-Hispanic) | | | | | | |
| Other race (Non-Hispanic) | | | | | | |
| Age group | | | | | | |
| Less than 25 | 0.75 | 0.75 | 0.73 | 0.86 | 0.76 | 0.82 |
| 25-29 | 0.82 | 0.81 | 0.78 | 0.92 | 0.82 | 0.88 |
| 30-34 | 0.91 | 0.90 | 0.83 | 0.96 | 0.88 | 0.93 |
| 35-39 | 0.90 | 0.89 | 0.82 | 0.97 | 0.88 | 0.93 |
| 40-44 | 0.97 | 0.95 | 0.87 | 1.01 | 0.91 | 0.97 |
| 45-49 | 1.05 | 1.03 | 0.94 | 1.10 | 0.98 | 1.04 |
| 50-54 | 1.08 | 1.07 | 0.95 | 1.10 | 1.00 | 1.06 |
| 55-59 | 1.15 | 1.14 | 0.97 | 1.14 | 1.06 | 1.13 |
| 60-64 | 1.08 | 1.06 | 0.77 | 0.89 | 0.97 | 1.02 |
| 65-69 | 1.26 | 1.25 | 0.62 | 0.73 | 1.19 | 1.27 |
| 70-74 | 1.41 | 1.43 | 0.52 | 0.64 | 1.11 | 1.24 |
| 75 and over | 1.42 | 1.45 | 0.47 | 0.53 | 1.44 | 1.46 |

The ratio of the NSSRN national estimate of "RNs with active licenses" to the ACS estimate for "those whose main work within the past 5 years was as an RN" is about 1.01. The ratio of the NSSRN "employed as an RN" estimate to the same ACS figure is 0.86. The ratio of the NSSRN "employed as an RN" estimate to the ACS estimate for those working as an RN in the past 5 years who are still in the labor force is 0.94.

Keeping in mind the methodological differences both in terms of the populations represented and how the data were collected, we examine the ratios for subgroups defined by race/ethnicity and by age for comparison purposes. Here we focus solely on the ratios for Hispanic nurses and for the non-Hispanic component of the racial groups White, Black, and Asian. The other subgroups were left out of this discussion as their numbers were quite small. Most noteworthy is that the various ratios for White are generally slightly more than 1 while the ratios for Black are generally between 0.5 and 0.6, the ratios for Asian are between 0.65 and 0.76, and the ratios for Hispanic are generally between 0.8 and 0.9.

Thus, to the extent that the NSSRN and ACS are comparable, these data suggest that there may be differential nonresponse between race/ethnic groups to the NSSRN with the response rates for minorities somewhat less than that of nurses who are White, non-Hispanic. Differential nonresponse among racial/ethnic groups seems particularly likely for Blacks and Asians. Such differential nonresponse is a potential source of bias.

It should be pointed out that there may be some evidence of the potential for misclassification affecting the identification of Black RNs in the ACS. A November 2004 report done for the Health Resources and Services Administration (HRSA) of the U.S. Department of Health and Human Services (HHS) entitled "Supply, Demand, and Use of Licensed Practical Nurses" provided estimates of the proportion of LPNs classified by race/ethnicity categories based on CPS data from 1984-2001. Blacks represented roughly 20 percent of LPNs over that time period with Hispanics about 5 percent and Asians 2-3 percent. Thus, if ACS respondents were prone to mischaracterizing LPNs as RNs, the minority race/ethnicity group that would seem most likely to be affected is that of Blacks. Such misclassification might have contributed to the low ratios of NSSRN to ACS estimates of Blacks who are RNs.

Examination of the ratios of the NSSRN full sample age group distributions to those of the ACS provides some evidence that the adjustment for differential response rates by age group was effective in reducing bias for estimates correlated with age. For the groups covering the overall age range 30-64, the ratios are roughly from 0.9 to 1.15 with those under 45 having ratios under 1 while

those from 46-64 exceeding 1. The ratios of estimated totals from Table B5-1 indicate that these "age" ratios in Table B5-2 would have been more extreme without the age-based adjustment since the estimates without such an adjustment were lower for the younger age ranges and larger for the older ones.

There do appear to be comparability issues between the NSSRN and the ACS, at least for the three oldest age ranges. For example the ratio of the NSSRN estimate of the proportion of nurses aged 65-69 who were employed in nursing to the ACS estimate of those RNs working in the past 5 years who are still in the labor force was 1.27. The corresponding ratios for 70-74 year olds and for those 75 or older are 1.24 and 1.46. It is not clear why an estimate of those actually working in nursing from one survey would exceed by a considerable margin the estimate from another survey of those remaining in the labor force. The sample sizes for these age groups are relatively small, so the corresponding estimates are somewhat unstable which may be one contributing factor to the relative size of the estimates.

References

- Krenzke, T., Judkins, D., and Fan, Z. (2005). *Vector imputation at Westat*. Presented at Statistical Society of Canada Meetings, Saskatoon, Saskatchewan.
- Lee, H., Carlson, E., Lo, A., Norman, G., and Lopdell, J. (2006). *Pre-elementary education longitudinal study, final wave3 update to methodology report.* Prepared under contract to the Office of Special Education Programs, U.S. Department of Education. Rockville, MD: Westat.
- Little, R.J. (1988). Missing-data adjustments in large surveys. *Journal of Business and Economic Statistics*, 6, 287-296.
- Piesse, A., Judkins, D., and Fan, Z. (2005). Item imputation made eEasy. Proceedings of the American Statistical Association Survey Research Methods Section.
- Lavallée, P. (2007). Indirect Sampling. Springer, New York
- Lohr, S. (2010). Sampling: Design and Analysis. Second edition. Brooks/Cole, Cengage Learning.
- Wolter, K.(2007). Introduction to Variance Estimation. Second Edition. Springer.
- Brick, J.M., and Kalton, G. (1996). Handling missing data in survey research. *Statistical Methods in Medical Research*, 5(3), 215-238.
- Sample Survey for the National Survey of Registered Nurses, Technical Report (Volume I), and Appendices (Volume II). Westat Inc. and the American Nurses Association, 1976 (unpublished).

- U.S. Department of Health and Human Services, Health Resources and Services Administration (2001). The Registered Nurse Population: March 2000 Findings from the National Sample Survey of Registered Nurses.
- U.S. Department of Health and Human Services, Health Resources and Services Administration (2006). The Registered Nurse Population: Findings from the March 2004 National Sample Survey of Registered Nurses.
- U.S. Department of Health and Human Services, Health Resources and Services Administration (2006). 2004 National Sample Survey of Registered Nurses: Technical Report (unpublished).

Appendix C

NSSRN Questionnaire



2008 National Sample Survey of Registered Nurses

The **2008** *National Sample Survey of Registered Nurses (NSSRN)* is being conducted by the Health Resources and Services Administration of the U.S. Department of Health and Human Services and is the ninth cycle of the survey.

Instructions

How do I complete the survey electronically?

On your Web browser, log onto www.nssrn.org and type in your unique Access Code and PIN that is printed in the box below. If you complete the survey online, you do not need to return this paper questionnaire.

What if I received more than one questionnaire?

We may not have been able to eliminate all of the duplicates in our list of nurses. Please complete only one questionnaire but return any extra copies you receive, preferably in the same envelope as your completed survey. Please write "DUPLICATE" at the top of these blank surveys. By returning extra surveys, we can avoid unnecessary follow-up mailings to you.

What if I have questions about this survey?

If you have any questions about this survey or about how to complete it electronically, please call (toll-free) 1-888-371-9725, or send an e-mail to nssrn2008@westat.com.

Please correct any errors in the name/address information

shown below to the right.

| Corrections to First Name Corrections to M.I. | | |
|--|--|-----------------|
| | | |
| Corrections to Last Name | | |
| | | |
| Corrections to Number and Street | First Name M.I. Last Name | |
| | Street Address City, State Zip Code | |
| Corrections to City/Town | | |
| | | |
| Corrections to State Corrections to ZIP Code | | |
| If there are <u>any</u> corrections to the "State(s) Where Actively Licensed", please relist <u>ALL</u> of the States where you are actively licensed. | State(s) Where Actively Licensed: State 1, State 2, State 3 | |
| | Web site URL: www.nssrn.org Access Code: [XXXXXXX] | PIN# [x] |
| Corrections to the State(s) Where Actively Licensed | OMB No.: 0915-0276 Expiration | Date: 5/31/2011 |

According to the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0915-0276. The time required to complete this information collection is estimated to average 20 minutes per response, including the time to review instructions, search existing data resources, gather the data needed, and complete and review the information collection. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to the HRSA Reports Clearance Officer, 5600 Fishers Lane, Room 10-33, Rockville, Maryland, 20857.

| Please mark an [X] in the box corresponding to your answer | r for each question, | or supply the requested information. | Use blue or black ink. |
|--|----------------------|--------------------------------------|------------------------|
| | EXAMPLE | | |

| RIGI | HT WAY | 7 |
|------|--------|---|
| | | |

| WRO | NG | WAY |
|-----|----|-----|
| X | 9 | 8 |

| | WRONG WAY |
|--|--|
| Section A. Eligibility and Education | |
| 1. On March 10, 2008, were you <u>actively licensed</u> to practice as a registered nurse (RN) in any U.S. State or the District of Columbia (whether or not you were employed in nursing at that time)? Yes → Go to Question 2 No → If No, you do not need to complete this questionnaire. Please mark "no" and return this questionnaire so we know you are not eligible. 2. In what U.S. State were you issued your first RN license? State: Year: Year: 3. Which type of nursing degree or nursing credential qualified you for your first U.S. RN license? <i>Mark one box only</i>. Diploma Program Associate Degree Bachelor's Degree Doctorate Other Specify 4. In what month and year did you graduate from this nursing program? | 6. Please indicate all post-high-school degrees you received before starting your initial RN educational program. Mark all that apply. None -> Go to Question 8 Associate Degree Bachelor's Degree Master's Degree Doctorate Other Specify 7. What was the field of study for your highest degree identified in Question 6? Mark one box only. Health-related field or Non-health related field Biological or Physical Science Business or Management Education Liberal Arts, Social Science, or Humanities Law Computer Science Social Work Other non-health related field Specify |
| Month: Year: | 8. Have you ever been <u>licensed</u> as a licensed practical nurse (LPN) or licensed vocational nurse (LVN) in the U.S.? Yes No |
| 5. In which U.S. State (including the District of Columbia), U.S. Territory, or foreign country was this program located? State: Philippines Canada United Kingdom Nigeria Other Specify | Continued on next page |

| h | Vere you ever employed in any of the following ealth-related jobs <u>before</u> completing your initial RN ducation? <i>Mark all that apply</i> . | 11: | a. Within the past year, have you received or provided emergency preparedness training, in any of the following areas? <i>Mark all that apply</i> . |
|---|--|-----|--|
| | No health-related position before RN education Nursing Aide or Nursing Assistant Home health aide or assistant Licensed Practical or Vocational Nurse Emergency Medical Technician (EMT) or Paramedic Medical assistant Dental assistant Allied Health technician or technologist (radiological technician, laboratory technician) Manager in health care setting Clerk in health care setting Military medical corps Medical doctor Midwife Another type of health-related position | 111 | None → Go to Question 12 Chemical accident or attack Nuclear/radiological accident or attack Infectious disease epidemics Biological accident or attack Natural disaster Other public health emergencies b. Please specify the total number of hours spent in the above training(s) within the past year. <i>(enter 0 if none)</i> Hours of training received Hours of training provided |
| | | 110 | c. Thinking about the area in which you are <u>best</u> prepared for an emergency, are you? |
| | Iow did you finance your <u>initial</u> RN education? <i>Iark all that apply.</i> Earnings from your health-care-related employment Earnings from your non-health-care-related | | Very prepared Adequately prepared Somewhat prepared Not at all prepared |
| | employment Earnings from other household members Personal household savings Other family resources (parents or other relatives) Employer tuition reimbursement plan (including Veterans Administration employer tuition plan) Federal traineeship, scholarship, or grant Federally-assisted loan Other type of loan State/local government scholarship or grant Non-government scholarship or grant | 12 | How well do you know the disaster/emergency plan at your place of employment? Full understanding Some understanding Little or no understanding No plan exists at my place of employment Do not have a place of employment |
| | Other resources | | Continued on next page |

13. Did you earn any additional academic degrees <u>after</u> graduating from your initial registered nurse education program that you described in Question 3? Do not include degrees you are currently working towards.

Yes \rightarrow Please complete all columns of the following table for each degree you earned. No \rightarrow Go to Question 14 on page 5

| | 18 | | | | | |
|--|--|--|---|---|---|--|
| | Α | В | С | D | Е | F |
| Type of Degree Nursing Degrees | Did you receive this degree? Mark all that apply. | What was the primary focus of this degree? Enter two-digit code from table below. | Has this degree been related to your career in nursing? | In what year did you receive the degree? | In what state or country was this educational program located? | Was this degree program undertaken through a distance-based learning program? (with more than 50% of coursework through correspondence or online) |
| Nursing Degrees | | | | | | |
| a. Associate degree in nursing | | | | | | Yes No |
| Bachelor's degree in nursing | | | | | | Yes No |
| c. Master's in nursing | | | | | | Yes No |
| d. Another Master's in nursing | | | | | | Yes No |
| e. Doctorate in nursing (PhD, ScD, DNS, ND, DNP) | | | | | | Yes No |
| Non-nursing Degrees | | | | ······································ | | |
| f. Associate degree in non-nursing field | | | Yes No | | | Yes No |
| g. Bachelor's degree in non-nursing field | | | Yes No | | | Yes No |
| Master's in non-nursing field | | | Yes No | | | Yes No |
| i. Another Master's in non-nursing field | | | Yes No | | | Yes No |
| j. Doctorate in non-nursing field (PhD, JD, MD, EdD) | | | Yes No | | | Yes No |

For Column B, enter the two-digit code for your Bachelor's, Master's, or Doctorate degree above.

Primary Focus of Degree

| 01 Clinical Practice | 07 Humanities, Liberal Arts, or Social Sciences |
|---------------------------------------|---|
| 02 Administration/Business/Management | 08 Informatics |
| 03 Education | 09 Computer Science |
| 04 Public health/community health | 10 Research |
| 05 Law | 11 Social Work |
| 06 Biological or Physical Sciences | 12 Other health field |
| | 13 Other non-health field |

14. <u>Since</u> graduating from the initial nursing program you described in Question 3, have you completed a formal educational program preparing you as a Nurse practitioner, Clinical nurse specialist, Nurse-midwife, or Nurse anesthetist?

| | Yes → Go to Question 14a No → Go to Question 15 on page 6 | | | | |
|-----|---|-------------------------------|--|-------------------------------|-------------------------------|
| | | Α | В | С | D |
| In | formation on preparation and credentials | Nurse Practitioner (NP) | Clinical Nurse Specialist (CNS) | Nurse- Midwife (NM) | Nurse Anesthetist (NA) |
| 14a | Did you receive preparation as a? Mark each column if yes. | | | | |
| 14b | What was the length of the program? 1. Less than 8 months 2. 8-12 months 3. 13-36 months 4. 37 months or more | (Mark one) | (Mark one) | (Mark one) | (Mark one) |
| 14c | What was the highest credential you received in that program? 1. Certificate/Award 2. Bachelor's degree 3. Master's degree 4. Post-Master's Certificate 5. Doctorate | (Mark one) | (Mark one) | (Mark one) | (Mark one) |
| 14d | In what year did you receive this credential? | | | | |
| 14e | Do you have certification from a <u>national</u> <u>certifying organization</u> for this specialty? <i>IF YES:</i> Is this certification required by your employer for your job? | Yes No Yes No N/A | Yes No Yes No N/A | Yes No Yes No N/A | Yes No Yes No N/A |
| 14f | • Do you have licensure, certification, or recognition from a <u>State Board of Nursing</u> for this specialty? <i>IF YES:</i> | Yes No | Yes No | Yes No | Yes No Yes |
| | Is this license, certification, or recognition required by your employer for your job? | No N/A | No N/A | No N/A | No N/A |

14g. Which specialties were the focus of your NP, CNS, NM, or NA studies? Mark all that apply.

| Acute Care | Geriatrics or Gerontology Palliative Care | ; |
|--|--|-----------------|
| Adult Health | Home Health Pediatrics | |
| Anesthesia | Maternal-Child Health Psychiatrics of | r Mental Health |
| Cardiac Care | Neonatal Rehabilitation | |
| Community Health | Nurse-Midwifery School Health | |
| Critical Care | Obstetrics or Gynecology Women's Heal | th |
| Family Care | Occupational Health Other | |
| General Medical Surgical | Oncology Specify | |
| Cardiac Care Community Health Critical Care Family Care | NeonatalRehabilitationNurse-MidwiferySchool HealthObstetrics or GynecologyWomen's HealOccupational HealthOther | |

| 15. | On March 10, 2008, were you enrolled in a formal education program leading to an academic degree or certificate? | |
|------|---|--|
| | Yes No → Go to Section B | |
| 16. | Was this formal education program? Mark one box only. | |
| | In nursing In a non-nursing field to enhance your career/employment in nursing In a non-nursing field to allow you to pursue career/employment opportunities <u>outside of nursing</u> In an area of personal interest without regard to future employment | |
| 17a. | Were you a full-time or part-time student? | |
| | Full-time studentPart-time student | |
| 17b. | What percent of your coursework was distance-based (online or correspondence)? | |
| | 0% 1-25% 26-50% 51-75% 76-100% | |
| 18. | What type of degree or certificate have you been working toward in this program? <i>Mark one box only</i> . | |
| | Associate Degree Bachelor's Degree Master's Degree Doctorate Post-Master's Certificate Other Certificate | |
| | | |
| | | |
| | | |

Section B. Principal Nursing Employment

19. On March 10, 2008, were you employed or self-employed in nursing? (Employed in nursing includes working for pay in nursing, even if on temporary leave.)

No → Go to Section D on page 10

For all the questions in this section (Questions 20 - 32), your principal nursing position is the nursing position, on March 10, 2008, in which you spent the largest share of your working hours.

- **20.** Are you required to maintain an active RN license in order to hold your principal nursing position?
 - Yes No

Yes

21. Where was the location of the <u>principal</u> nursing position you held on March 10, 2008? (If you are not employed in a fixed location, enter the location that best reflects where you practice.)

| City/Town: |
|-------------------------------------|
| County: |
| State (or country if not U.S.A.) |
| ZIP+4 code (if available) |

22. In the principal nursing position you held on March 10, 2008, were you...

Mark one box only.

- An employee of the organization or facility where you were working?
- Employed through an employment agency, but <u>not</u> as a traveling nurse?
- Employed through an employment agency as a traveling nurse?
- Self-employed, per diem, or working as-needed?

| Hospital (including all types of care at a hospital location) | Academic Education Program |
|--|---|
| Community hospital or medical center, Non-Federal, short stay | Nursing aide and/or home health aide program |
| Inpatient unit | LPN/LVN program |
| Nursing home unit in hospital | Diploma program (RN) |
| Outpatient clinic/medical practice owned by a hospital | Associate degree RN program |
| Outpatient clinic/medical practice located at a hospital but | Bachelor's and/or higher degree RN program |
| not owned by the hospital | Associate degree RN and LPN/LVN program |
| Other administrative or functional area | Associate degree RN and BSN program |
| Specialty hospital, Non-Federal (children's, heart, burns, cancer) Inpatient unit | Other education program, not patient education |
| Outpatient clinic/medical practice owned by a hospital | Specify |
| Outpatient clinic/medical practice located at a hospital but | |
| not owned by the hospital | Public or Community Health Setting |
| Other administrative or functional area | State Health or Mental Health Agency |
| Long-term hospital, Non-psychiatric, Non-Federal | City or County Health Department |
| Inpatient unit | Correctional Facility (non-hospital) |
| Nursing home unit in hospital | Community mental-health organization or clinic |
| Other administrative or functional area | Substance abuse center/clinic |
| Psychiatric hospital, Non-Federal | Other community setting |
| Inpatient unit | |
| Nursing home unit in hospital | Specify |
| Outpatient clinic/medical practice owned by a hospital | School Health Service |
| Outpatient clinic/medical practice located at a hospital but | School or school system (K-12) |
| not owned by the hospital | College or university |
| Other administrative or functional area | Other school health setting |
| Federal Government hospital (Military, VA, NIH | Occupational Health (Employee Health Service) |
| or IHS-supported) | Private industry |
| Inpatient unit | Government occupational health services |
| Nursing home unit in hospital | Other occupational health setting |
| Outpatient clinic/medical practice located at a hospital | Ambulatory Care Setting, not located in a hospital |
| Other administrative or functional area | Medical/physician practice |
| Hospital unit in an institution (infirmary, correctional facility) | Nurse practice |
| All types | In-store or retail clinic |
| Other Type of hospital | Community health center |
| Inpatient unit | Federal clinic (Military, VA, NIH or IHS-supported) |

| | |
|--------------|-------|
| npatien | tunit |
| inpation | i uni |
| 1 | |

Nursing home unit in hospital

Outpatient clinic/medical practice owned by a hospital

| Outpatient clinic/medical | l practice | located | at a | hospital | but |
|---------------------------|------------|---------|------|----------|-----|
| not owned by the hospita | .1 | | | | |

Other administrative or functional area

Specify

Nursing Home/Extended Care Facility

- Nursing home/extended care facility (not in a hospital)
- Facility for mentally retarded or developmentally disabled
- Residential care/assisted living facility
- Other type of extended care facility

Specify

Home Health Setting

- Visiting nurse service (VNS/VNA)
- Home health service unit (hospital-based)
- Home health agency (non-hospital based)
- Private duty in a home setting
- Hospice
 - Other home health setting

Insurance Claims/Benefits/Utilization Review

Hospital-owned off-site clinic or surgery center

Ambulatory surgical center, not hospital-owned

Dialysis center or clinic, not in a hospital

Federally-supported clinic (not a community health center)

Government insurer/benefits department: federal, state, or local Insurance company or other private

claims/benefits/utilization review organization **Other**

- Policy, planning, regulatory, or licensing agency
- Consulting organization or self-employed
- Home-based self-employment

Urgent care

Specify

Specify

Other ambulatory setting

- Telehealth, telenursing, or call center
- Pharmaceutical/medical device/medical software Other

| Which one of the following <u>best</u> corresponds to the job title for the principal nursing position you held on March 10, 2008? <i>Mark one box only</i>. Staff nurse or direct care nurse Charge nurse or team leader First-line management (head nurse, floor supervisor) Middle management/administration (assistant director, house supervisor, associate dean, department head) Senior management/administration (CEO, vice president, nursing executive, dean) Certified Registered Nurse Anesthetist (CRNA) Clinical Nurse Specialist (CNS) Certified Nurse-Midwife (CNM) | | | | |
|--|--|--|--|--|
| Charge nurse or team leader First-line management (head nurse, floor supervisor) Middle management/administration (assistant director, house supervisor, associate dean, department head) Senior management/administration (CEO, vice president, nursing executive, dean) Certified Registered Nurse Anesthetist (CRNA) Clinical Nurse Specialist (CNS) Certified Nurse-Midwife (CNM) Nurse practitioner (NP) School nurse Public health nurse Community health nurse Patient educator Staff educator or instructor in clinical setting Staff development director Instructor/lecturer Professor Patient care coordinator, case manager, discharge planner Quality improvement nurse, utilization review nurse Infection control Advice/triage nurse Consultant Legal nurse | | | | |
| Patient educator Staff educator or instructor in clinical setting Staff development director Instructor/lecturer Professor Patient care coordinator, case manager, discharge planner Quality improvement nurse, utilization review nurse Infection control Advice/triage nurse Informatics nurse Consultant Legal nurse Researcher Surveyor/auditor/regulator No position title Other | | | | |
| Specify | | | | |
| Senior management/administration (CEO, vice president, nursing executive, dean) Certified Registered Nurse Anesthetist (CRNA) Clinical Nurse Specialist (CNS) Certified Nurse-Midwife (CNM) Nurse practitioner (NP) School nurse Public health nurse Community health nurse Patient educator Staff educator or instructor in clinical setting Staff development director Instructor/lecturer Professor Patient care coordinator, case manager, discharge planner Quality improvement nurse, utilization review nurse Infection control Advice/triage nurse Informatics nurse Consultant Legal nurse Researcher Surveyor/auditor/regulator No position title Other Specify Specify For the principal nursing position you held on March 10, 2008, did you work? Mark one box only. Full-time (including full-time for an academic year) Part-time (including working only part of the calendar or academic year) | | | | |
| Staff development director Instructor/lecturer Professor Patient care coordinator, case manager, discharge planner Quality improvement nurse, utilization review nurse Infection control Advice/triage nurse Informatics nurse Consultant Legal nurse Researcher Surveyor/auditor/regulator No position title Other Specify For the principal nursing position you held on March 10, 2008, did you work? Full-time (including full-time for an academic year) Part-time (including working only part of the calendar or academic year) For the principal position you held on March 10, 2008, | | | | |
| 26. For the principal position you held on March 10, 2008, how many months would you normally work per year? | | | | |
| months | | | | |
| | | | | |

27. For the principal nursing position you held on March 10, 2008, please provide information about the number of hours you work in a typical week.

Hours (enter 0 if none)

a. Number of hours worked, including all overtime and on-call hours, <u>except</u> on-call hours that were stand-by only



- b. Number of hours you stated above in "a" that were worked from <u>on-call</u> <u>duty</u>. Do not include stand-by hours
- Number of hours you stated above in "a" that were <u>paid</u> as overtime
- **d.** Number of paid overtime hours you stated above in "c" that were <u>mandatory</u> overtime
- e. Number of paid or unpaid on-call hours that were stand-by <u>only</u>

- **f.** Number of stand-by hours you stated above in "e" that were <u>paid</u> at an on-call stand-by rate
- **28.** For the principal nursing position you held on March 10, 2008, please estimate the percentage of your time spent in the following activities during a usual workweek. *Do not use decimal places*.

| | The total should equal 100% | 100% | |
|----|--|------|---|
| 1. | Other | | % |
| з. | Teaching, precepting or orienting students or new hires (include preparation time) | | % |
| f. | Research | | % |
| e. | Administration | | % |
| 1. | Supervision and management | | % |
| c. | Consultation with agencies and/or professionals | | % |
| b. | Non-nursing tasks (housekeeping, locating supplies) | | % |
| a. | Patient care and charting | | % |

| 29a. For the principal nursing position you held on March 10, 2008, in what <u>level of care</u> or <u>type of work</u> did you spend the majority of your time? <i>Mark one or more boxes</i> . | 29c. For the principal nursing position you held on March 10, 2008, in what type of <u>clinical specialty</u> did you spend most of your patient care time? <i>Mark one or more boxes</i> . |
|---|---|
| General or specialty inpatient Critical/intensive care Step-down, transitional, progressive, telemetry Sub-acute care Emergency Urgent care Rehabilitation Long-term care/nursing home Surgery (including ambulatory, pre-operative, post-operative, post-anesthesia) Ambulatory care (including primary care, outpatient settings, except surgical) Ancillary care (radiology, laboratory) Home health Public health/community health Education Business, administration, review, case management Research Other Specify 29b. For the principal nursing position you held on March 10, 2008, with what patient population did you spend at least 50% of your patient care time? <i>Mark only one box.</i> | No patient care General medical surgical Critical care Cardiac or cardiovascular care Chronic care Dermatology Emergency or trauma care Gastrointestinal Gynecology (women's health) Hospice Infectious/communicable disease Labor and delivery Neurological Obstetrics Occupational health Oncology Primary care Psychiatric or mental health (substance abuse and counseling) Pulmonary/respiratory Radiology (diagnostic or therapeutic) Renal/dialysis No specific area Other specialty for a majority of my time |
| > No patient care → Go to Question 30 Adult Geriatric Pre-natal Newborn or neonatal Pediatric and/or Adolescent Multiple age groups (less than 50% time spent with any of the above) Specify | 30. Please estimate your 2008 pre-tax <u>annual</u> earnings from your principal nursing position. Include overtime and bonuses, but exclude sign-on bonuses. \$,,00 per year 31. Were you represented by a labor union or collective bargaining unit in the principal nursing position you held on March 10, 2008? Yes Yes No |
| | 32a. Do you plan to leave or have you left the principal nursing position you held on March 10, 2008? Yes, have left or will leave within the next 12 months Yes, in 1 year to 3 years No plans to leave within next 3 years Undecided → Go to Question 33 on page 10 32b. Do you plan to work in nursing after you leave that position? Yes No Unsure |


Section C. Secondary Employment in Nursing

33. Aside from the principal nursing position you just described, did you hold any other positions in nursing for pay on March 10, 2008?

Yes No \rightarrow Go to Section E on page 11

34. In your <u>other nursing position(s)</u>, are you...? *Mark all that apply.*

- An employee of the organization or facility for which you are working?
- Employed through an employment agency, but <u>not</u> as a traveling nurse?
- Employed through an employment agency as a traveling nurse?
- Self-employed, per diem, or working as needed?



| Hospital | |
|--|--|
| Nursing home/Extended care facility | |
| Academic education program | |
| Home health setting | |
| Public or community health setting | |
| School health service | |
| Occupational health | |
| Ambulatory care setting | |
| Insurance claims/benefits | |
| Telehealth, telenursing or call center | |
| Other | |
| Specify | |

36. In your additional nursing position(s), please indicate how much you work, and where the job is located:

| | Weeks per year | Average hours per week, during weeks of work | Locations of where most of work is done (state, or country) |
|----------------------|----------------------|---|---|
| Additional job #1 | | | |
| Additional job #2 | | | |
| All other jobs | | | N/A |

37. Please estimate your 2008 pre-tax annual earnings from all your <u>other nursing position(s)</u>. Do not include earnings from your principal nursing position.



All go to Section E on page 11

Section D. Nurses Not Working in Nursing

If you were working for pay in nursing on March 10, 2008, please go to Section E on page 11.

38. What are your intentions regarding paid work in registered nursing? *Mark one box only.*

| Have returned to nursing since March 10, 2008 | | |
|--|--|--|
| → Go to Section E on page 11 | | |
| Actively looking for work in nursing | | |
| Plan to return to nursing in the future, not looking | | |
| for work now | | |
| → Go to Question 41 | | |
| No future intention to work for pay in nursing | | |
| \rightarrow Go to Question 42 on page 11 | | |
| Undecided at this time | | |
| ightarrow Go to Question 42 on page 11 | | |

- **39.** How long have you been actively looking for paid work in nursing?
 - Months (if one or more)
 - Less than one month
- **40.** Are you looking for a position that is ...?

| | Full-time Part-time Either | → Go to Question 42 on page 11 |
|--|----------------------------------|--------------------------------|
|--|----------------------------------|--------------------------------|

- **41.** When do you plan to return to paid work in nursing?
 - Years (if one or more)
 - Less than one year

| 42. How long has it been since you last were employed or self-employed as a registered nurse? | 46a. Please select from the list below the item that best describes the <u>field</u> of your principal position <u>outside</u> of nursing. <i>Mark one box only</i> . |
|---|---|
| Years (if one or more) Less than one year Never worked as a Registered Nurse | Computer services Consulting organization Emergency response (ambulance, fire, police) Financial, accounting, and insurance services Legal Education, elementary and secondary |
| 43. What are the primary reasons you are not working in a nursing position for pay? <i>Mark all that apply</i>. Retired Taking care of home and family Burnout Stressful work environment Scheduling/inconvenient hours/too many hours Physical demands of job Disability Ilness Inadequate staffing Salaries too low/better pay elsewhere Skills are out-of-date Liability concerns Lack of collaboration/communication between health care professionals Inability to practice nursing on a professional level Lack of good management or leadership Career change Difficult to find a nursing position Travel Volunteering in nursing Went back to school Other | Food services Government Health-related services, outside nursing Pharmaceutical, biotechnology, or medical equipment Real estate Retail sales and services Other Specify 46b. Which of the following best describes your job title for your principal position <u>outside</u> of nursing? Mark one box only. Business owner or proprietor Management Sales Instructor or professor Administrative or clerical support Consultant Other type of employee Specify 47. How many months would you normally work per year in this principal position <u>outside</u> of nursing? |
| Section E. Employment Outside Nursing | 48. What is the average number of hours you work per week in your principal position <u>outside</u> of nursing? |
| 44. On March 10, 2008, were you employed for pay in an occupation other than nursing? | hours per week |
| Yes No → Go to Section F on page 12 | 49. Please estimate your 2008, pre-tax annual earnings from your principal position <u>outside</u> of nursing. |
| 45. On March 10, 2008, was this non-nursing employment with a health-related organization or in a health-related position? | \$, |
| Yes No | |

Section F. Prior Nursing Employment

| 5 | 1. For this question count only the years you worked at least 50% of the calendar year in nursing. Since receiving your first U.S. RN license, how many years have you worked in nursing? | 55 |
|----|--|----|
| | Years (if one or more) | |
| 5 | • Have you left work in nursing for one or more years since becoming an RN? | |
| | Yes Total years (if one or more) No | |
| 5′ | Have not worked in nursing more than one yearWere you employed in nursing one year ago | |
| 5. | (March 10, 2007)? | 56 |
| | Yes No → Go to Section G on page 14 | |
| 5. | For the principal nursing position you held on <u>March 10</u>, <u>2007</u>, did you work? <i>Mark one box</i>. | |
| | Full-time (including full-time for an academic year) Part-time (including working only part of the calendar or academic year) | |
| 54 | How would you describe the principal nursing position you held on <u>March 10, 2007</u>? | |
| | Same position/same employer as principal nursing position on March 10, 2008 Go to Section G on page 14 Different position/same employer as current one | |
| | Different employer than current one | |
| | | |
| | | |
| | | |
| | | |

55. What was the location of the principal nursing position you held on <u>March 10, 2007</u>? (If you were <u>not</u> employed in a fixed location enter the location that best reflects where you practice.)

| City/Town: |
|-------------------------------------|
| County: |
| State (or country if not U.S.A.) |
| ZIP+4 code (if available) |

56. Were any of the following the <u>primary</u> reason(s) for your employment change? *Mark all that apply.*

| Burnout |
|---|
| Stressful work environment |
| Interested in another position/job |
| Lack of advancement opportunities |
| Lack of collaboration/communication between |
| health care professionals |
| Lack of good management or leadership |
| Career advancement/promotion |
| Inadequate staffing |
| Interpersonal differences with colleagues or |
| supervisors |
| Physical demands of job |
| Opportunity to do the kind of nursing that I like |
| Pay/benefits better |
| Scheduling/inconvenient hours/too many hours |
| Relocated to different geographic area |
| Reorganization that shifted positions |
| Laid off/downsizing of staff |
| Sign-on bonus offered |
| Personal/family |
| Went back to school |
| Retired |
| Disability |
| Illness |
| Other |
| Specify |

| 57. Whi | 7. Which one of the following best describes the employment setting of the principal nursing position you held on March 10, | | | | |
|---------|---|--|--|--|--|
| 2007 | 7? Mark one box only. | | | | |
| H | lospital (including all types of care at a hospital location) | | Academic Education Program | | |
| | Community hospital or medical center, Non-Federal, short stay | | Nursing aide and/or home health aide program | | |
| | Inpatient unit | | LPN/LVN program | | |
| | Nursing home unit in hospital | | Diploma program (RN) | | |
| | Outpatient clinic/medical practice owned by a hospital | | Associate degree RN program | | |
| | Outpatient clinic/medical practice located at a hospital but | | Bachelor's and/or higher degree RN program | | |
| | not owned by the hospital | | Associate degree RN and LPN/LVN program | | |
| | Other administrative or functional area | | Associate degree RN and BSN program | | |
| | Specialty hospital, Non-Federal (children's, heart, burns, cancer) | | Other education program, not patient education | | |
| | Inpatient unit | | G · · C | | |
| | Outpatient clinic/medical practice owned by a hospital | | Specify | | |
| | Outpatient clinic/medical practice located at a hospital but | | | | |
| | not owned by the hospital | | Public or Community Health Setting | | |
| | Other administrative or functional area | | State Health or Mental Health Agency | | |
| | Long-term hospital, Non-psychiatric, Non-Federal | | City or County Health Department | | |

Specify

Inpatient unit

Inpatient unit

or IHS-supported) Inpatient unit

All types

Specify

Inpatient unit

Other Type of hospital

Nursing home unit in hospital

Psychiatric hospital, Non-Federal

Nursing home unit in hospital

not owned by the hospital

Nursing home unit in hospital

Nursing home unit in hospital

not owned by the hospital

Other administrative or functional area

Nursing Home/Extended Care Facility

Residential care/assisted living facility Other type of extended care facility

Outpatient clinic/medical practice owned by a hospital

Outpatient clinic/medical practice located at a hospital

Outpatient clinic/medical practice owned by a hospital

Nursing home/extended care facility (not in a hospital) Facility for mentally retarded or developmentally disabled

Outpatient clinic/medical practice located at a hospital but

Hospital unit in an institution (infirmary, correctional facility)

Federal Government hospital (Military, VA, NIH

Outpatient clinic/medical practice located at a hospital but

Home Health Setting

- Visiting nurse service (VNS/VNA)
- Home health service unit (hospital-based)
- Home health agency (non-hospital based)
- Private duty in a home setting
- Hospice
 - Other home health setting

Nurse practice In-store or retail clinic Community health center Federal clinic (Military, VA, NIH or IHS-supported)

Ambulatory Care Setting, not located in a hospital

Federally-supported clinic (not a community health center)

Hospital-owned off-site clinic or surgery center

- Ambulatory surgical center, not hospital-owned
- Urgent care
- Dialysis center or clinic, not in a hospital Other ambulatory setting

Correctional Facility (non-hospital)

Substance abuse center/clinic

School or school system (K-12)

Other occupational health setting

Other community setting

School Health Service

College or university

Private industry

Other school health setting

Medical/physician practice

Specify

Community mental-health organization or clinic

Occupational Health (Employee Health Service)

Government occupational health services

Specify

Specify

| | Insurance Claims/Benefits/Utilization Review |
|----|---|
| | Government insurer/benefits department: federal, |
| | state, or local |
| | Insurance company or other private |
| | claims/benefits/utilization review organization |
| | <u>Other</u> |
| | Policy, planning, regulatory, or licensing agency |
| | Consulting organization or self-employed |
| | Home-based self-employment |
| | Telehealth, telenursing, or call center |
| | Pharmaceutical/medical device/medical software |
| Ē. | Other |

-

| Section G. General Information | 64a. Are you of Latino or Hispanic ethnicity? |
|--|---|
| 58. How satisfied are you with your principal job, or most recent job if you are not now working? <i>Mark one box only</i>. Extremely satisfied Moderately satisfied Neither satisfied nor dissatisfied Moderately dissatisfied Extremely dissatisfied Neither currently nor previously employed 59. Where do you currently reside? This information is critical for producing State estimates of the nursing workforce. City/Town: ZIP+4 code (if available) 60. Did you reside in the same city/town a year ago (March 10, 2007)? Yes → Go to Question 62 No 61. Where did you reside a year ago? This information is critical for producing State estimates. | Yes No 64b. Which one or more of the following would you use to describe your race? <i>Please see page 16 for definitions. Mark all that apply.</i> White Black or African American Asian American Indian or Alaska Native Native Hawaiian or Other Pacific Islander 65. What languages do you speak fluently, other than English? <i>Mark all that apply.</i> No other languages Spanish Filipino language (Tagalog, other Filipino dialec Chinese language) French German American Sign Language Other Other Other Other Married or in domestic partnership Widowed, divorced, separated Never married 67. Describe the children/parents/dependents who either live at home with you or for whom you provide a significant amount of care. <i>Mark all that apply.</i> |
| County: State (or country if not U.S.A.) ZIP+4 code | No children/parents/dependents at home Child(ren) less than 6 years old at home Child(ren) 6 to 18 years old at home Other adults at home (i.e., parents or dependents) Others living elsewhere (i.e., children, parents or dependents) 68. Including employment earnings, investment earnings, and other income of all household members, what is your current, pre-tax annual total household income? <i>Pick one appropriate category</i>. \$15,000 or less \$15,001 to \$25,000 \$25,001 to \$35,000 \$50,001 to \$50,000 \$100,001 to \$150,000 \$100,001 to \$150,000 \$150,001 to \$200,000 More than \$200,000 |

Section H. License and Certification Detail

|)a. Do | you currently have any National nursing certifications? |
|---------------|--|
| | Yes No → Go to Section I on page 16 |
| • | Fig |
| | nich of the following <u>skill-based</u> certifications do a currently have? <i>Mark all that apply</i> . |
| <i>y</i> 0 | No current skill-based certifications |
| | Life Support (BLS, ALS, BCLS, and others) |
| | Resuscitation (CPR, NRP, and others) |
| | Emergency Medicine/Nursing (EMT, ENPC, and |
| | others) Trauma Nursing (TNCC, ATCN, ATN, and others) |
| | Other |
| | Specify |
| | Other |
| | Specify |
| | |
|)c. W | nich of the following Nurse Practitioner, Clinical Nurse |
| | ecialist, Nurse Midwife, or Nurse Anesthetist |
| cei | tifications do you currently have? Mark all that apply. |
| | No current Nurse Practitioner, Clinical Nurse |
| | Specialist, Nurse Midwife, or Nurse Anesthetist |
| | certifications |
| | |
| Nu | <u>rse Practitioner (NP)</u> |
| Nu | rse Practitioner (NP) Acute Care NP |
| Nu | Acute Care NP Adult NP |
| | Acute Care NP Adult NP Diabetes Management NP |
| | Acute Care NP Adult NP Diabetes Management NP Family NP |
| | Acute Care NP Adult NP Diabetes Management NP |
| | Acute Care NP Adult NP Diabetes Management NP Family NP Gerontological NP Hospice and Palliative Care NP Neonatal NP |
| | Acute Care NP Adult NP Diabetes Management NP Family NP Gerontological NP Hospice and Palliative Care NP Neonatal NP Oncology NP |
| | Acute Care NP Adult NP Diabetes Management NP Family NP Gerontological NP Hospice and Palliative Care NP Neonatal NP Oncology NP Pediatric NP (CPNP/CPNP-PC/CPNP-AC) |
| | Acute Care NP Adult NP Diabetes Management NP Family NP Gerontological NP Hospice and Palliative Care NP Neonatal NP Oncology NP |
| | Acute Care NP Adult NP Diabetes Management NP Family NP Gerontological NP Hospice and Palliative Care NP Neonatal NP Oncology NP Pediatric NP (CPNP/CPNP-PC/CPNP-AC) Psychiatric & Mental Health NP - Adult |

| <u>Clini</u> | <u>cal Nurse S</u> | Specialist (CNS) |
|--------------|--|--|
| | Adult Heat Communit Diabetes M Gerontolog Home Heat Hospice ar Medical-S Oncology Pediatric C Psychiatric | y Health/Public Health CNS Aanagement CNS gical CNS lth CNS ad Palliative Care CNS urgical CNS CNS CNS 2 & Mental Health CNS - Adult 2 & Mental Health CNS - Child/Adolescent 2 & Mental Health CNS - Family |
| | Specify | |
| <u>Nurs</u> | <u>e Midwife</u> | |

Nurse Midwife (CNM)

Nurse Anesthetist

- Nurse Anesthetist (CRNA)
- **70d.** Other than those previously listed, what other National nursing certifications do you currently have? *Specify name or acronym and organization below.*

■ No other current National nursing certifications → Go to Section I on page 16

Certification Name:

Certifying Organization:

Certification Name:

Certifying Organization:

Certification Name:

Certifying Organization:

Continued on next page

Specify

Other

Women's Health Care NP

Section I. Contact Information/Comments

| E-mail address: | | |
|---|---|--|
| Telephone No.: | Home Work | Cell (Area Code Telephone Number |
| Time of day/week be | st to contact you by phone: | |
| 72. Do you have any re | ecommendations for how this survey co | uld be improved? Please print clearly. |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | Tha | nk You. |
| | e return this surve | y and any duplicate surveys |
| | e return this surve | |
| Definitions for Question 6 White: origins in any of the | e return this survey in the enclosed, po | y and any duplicate surveys |
| Definitions for Question 6 White: origins in any of the East, or North Africa | e return this survey in the enclosed, po | y and any duplicate surveys |
| Definitions for Question 6 White: origins in any of the East, or North Africa Black or African America groups of Africa Asian: origins in any of the Southeast Asia, or the India | e return this survey in the enclosed, po 4b e original peoples of Europe, the Middle an: origins in any of the black racial e original peoples of the Far East, in subcontinent including, for example, pan, Korea, Malaysia, Pakistan, the | y and any duplicate surveys |
| Definitions for Question 6 White: origins in any of the East, or North Africa Black or African America groups of Africa Asian: origins in any of the Southeast Asia, or the India Cambodia, China, India, Ja Philippine Islands, Thailand American Indian or Alask | e return this survey in the enclosed, po 44b e original peoples of Europe, the Middle an: origins in any of the black racial e original peoples of the Far East, in subcontinent including, for example, pan, Korea, Malaysia, Pakistan, the d, and Vietnam ca Native: origins in any original peoples America and who maintains tribal | y and any duplicate surveys ostage-paid envelope. FOR OFFICE |

